

# Screening of Process Variables for the Production of Extracellular Lipase from Palm Oil by *Trichoderma Viride* using Plackett-Burman Design

R. Rajendiran, S. Gayathri devi, B.T. SureshKumar and V. Arul Priya

**Abstract**—Plackett-Burman statistical screening of media constituents and operational conditions for extracellular lipase production from isolate *Trichoderma viride* has been carried out in submerged fermentation. This statistical design is used in the early stages of experimentation to screen out unimportant factors from a large number of possible factors. This design involves screening of up to 'n-1' variables in just 'n' number of experiments. Regression coefficients and t-values were calculated by subjecting the experimental data to statistical analysis using Minitab version 15. The effects of nine process variables were studied in twelve experimental trials. Maximum lipase activity of 7.83  $\mu\text{mol/ml/min}$  was obtained in the 6th trial. Pareto chart illustrates the order of significance of the variables affecting the lipase production. The present study concludes that the most significant variables affecting lipase production were found to be palm oil, yeast extract,  $\text{K}_2\text{HPO}_4$ ,  $\text{MgSO}_4$  and  $\text{CaCl}_2$ .

**Keywords**—lipase, submerged fermentation, statistical optimization, *Trichoderma viride*

## I. INTRODUCTION

LIPASES (Triacylglycerol acylhydrolases E.C.3.1.13) are found in animals, plants and microorganisms [1], but only microbial lipases are commercially significant [2]. Lipases have been extensively studied because of their actual and potential applications in the detergent, oil, food, pharmaceutical and chemical industries [3].

Among the lipase producing microorganisms, fungi are preferable lipase sources because fungal enzymes are usually excreted extracellularly, facilitating extraction from fermentation media [4]. Many fungi as *Rhizopus oryzae* [5], *Geotrichum* sp., [6], *Aspergillus terreus*, [7], *Trichoderma viride* [8], *Trichoderma reeshi* [9] and *Fusarium solani* [10]

R.Rajendiran is with the Poultech Agro Research centre Bioinput Production Unit, Namakkal, TamilNadu, India ( Phone: +91-9994262289; Fax: 91-04286-275171; e-mail: rrmicro@gmail.com)

S. Gayathri devi is with department of Microbiology, Vivekanandha College of Arts and Sciences for Women, Tiruchengode, Namakkal, Tamilnadu, India (e-mail: gayumicro@yahoo.co.in).

B.T.Sureshkumar is with department of Microbiology, Vivekanandha College of Arts and Sciences for Women, Tiruchengode, Namakkal, Tamilnadu, India (e-mail: btsmicro@gmail.com).

V. Arulpriya is with the Poultech Agro Research centre Bioinput Production Unit, Namakkal, TamilNadu, India (e-mail: arulblue@yahoo.co.in).

have been noted as producers of lipases using vegetable oil for carbon and energy source for lipase production. But, by considering the rate, all vegetable oils were highly expensive except palm oil. So, in industries increasing the contents of palm oil, the lipase yields also become increasing with low cost, because they were the cheapest substrate and available in higher volume. A cheap and effective medium can decrease the production cost, so it is crucial to the industrialization of the enzyme method [11].

The application of statistically based experimental designs to optimize fermentation media is an efficient approach to studying the effects of several factors and to improve product yields. The Plackett-Burman design provides an efficient way of a large number of variables and identifying the most important ones [12]. Numerous reports have proved the applicability of Plackett-Burman design in the optimization of media components for various enzyme production such as the optimization of Xylanase by *Aspergillus niger* LPB 326 [13] and lipase production by *Yarrowia lipolytica* NCIM 3589 [14].

Hence an attempt is made in this paper to utilize the palm oil as a substrate for the production of lipase using indigenous isolate of *Trichoderma viride* by submerged fermentation. It was under taken to optimize the key process variables, including incubation time, pH, carbon level, nitrogen level and salt level of the medium for the production of lipase using Plackett- Burman design.

## II. MATERIALS AND METHODS

### Isolation of lipolytic fungi

Lipase producing fungal isolate of *Trichoderma viride* was isolated from oil mill waste which was collected from December 2010 to February 2011 at Namakkal, Tamil Nadu, and India by serial dilution technique [15]. After isolation, the fungus was subcultured and the pure culture was stored in PDA at 4°C.

### Lipase activity on Rhodamine-B agar plates

The fungal strain was tested for lipase production on agar plates that contained 15g, olive oil 10 ml, Rhodamine-B 0.01g and Tween 80- 0.001% in 1000 ml of distilled water. After 7 days of incubation at 28°C, the plate was irradiated with UV light (350 nm), with a bright pink fluorescent halo confirming lipolytic activity [16].

### A. Development of Inoculums

The culture was grown on Potato Dextrose Agar (PDA) plates for 5 days, which were wetted by adding 10 ml of sterilized distilled water. With the help of inoculating needle, conidia were scratched and the tubes were shaken vigorously to break the clumps of conidia and to obtain homogeneous suspension. Conidial suspension was used as inoculums and the number of spores was counted in a Neubauer hemocytometer and the concentration of the spores was adjusted to  $1 \times 10^9$  with sterile distilled water.

### B. Lipase production

Medium optimization for maximum lipase production was done in the basal liquid medium (Table 1 and 2). The nutritional components and operating conditions such as palm oil, Yeast extract,  $\text{KH}_2\text{PO}_4$ ,  $\text{K}_2\text{HPO}_4$ ,  $\text{MgSO}_4$ ,  $\text{CaCl}_2$ ,  $\text{NaNO}_3$ , pH and incubation periods were varied according to the statistical designs and the optimum components were incorporated in the basal medium. Erlenmeyer flask with 100 ml liquid medium was inoculated with 2 ml spore suspension and incubated at  $30 \pm 2^\circ\text{C}$  and 120 rpm. After the fermentation periods, the culture filtrate was collected aseptically by passing through Whatman No. 1 filter paper and filtrate was assayed for lipase productivity.

### C. Assay for lipase activity

Lipase activity was determined in an emulsifier free system using olive oil as substrate [17]. Reactions were carried out in 100 ml conical flasks at  $40^\circ\text{C}$  by immersion in a water bath. The reaction mixture, consisting of 2 ml of 0.1 M potassium phosphate buffer, pH 7.0, 1 ml of olive oil and 1 ml of culture supernatant was incubated at  $40^\circ\text{C}$  for 30 min. The enzyme reaction was terminated by the addition of 5 ml of 96% ethanol and followed by titration with 0.01 N NaOH solution using phenolphthalein as the indicator. One unit of lipase activity was defined as the amount, which liberated  $1 \mu\text{mol}$  of fatty acid per min at  $40^\circ\text{C}$ .

### D. Plackett-Burman design

The Plackett-Burman statistical experimental design is used for experiments confronted with two-factor interaction. This is a very economical design with the run number a multiple of four and comprises of two level screening designs. This design is extremely useful in finding importance of the factors affecting the production of the enzyme. PB design mathematically computes the significance of a large number of factors in one experiment, which is time saving and gives the effect of change in more than one factors in single experiment [18]. To evaluate the effect of 9 factors of medium components and operating conditions on lipase activity PB factorial design in 12 experimental run was carried out. Nine assigned variables and two unassigned variables (commonly referred as dummy variables) were screened in PB design of 12 experiments. Dummy variable are used to estimate experimental errors in data analysis [19]. Nine factors consisting of medium components and operating conditions prepared at two levels  $-1$  for low level and  $+1$  for high level. The factors (in g or ml /1000ml) such as palm oil, yeast

extract,  $\text{KH}_2\text{PO}_4$ ,  $\text{K}_2\text{HPO}_4$ ,  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ ,  $\text{CaCl}_2$ ,  $\text{NaNO}_3$ , pH and Incubation periods and 2 unassigned variables at same level were studied. Each variable is represented at two levels, high and low denoted by  $(+1)$  and  $(-1)$  respectively (Table 1).

TABLE I  
ACTUAL VALUES OF THE PROCESS VARIABLES FOR 1000ML OF MEDIUM

Process variables	Low levels (-1)	High Level (+1)
Palm oil (ml)	5	20
Yeast extract (g)	15	40
$\text{KH}_2\text{PO}_4$ (g)	0.5	3.0
$\text{K}_2\text{HPO}_4$ (g)	0.5	3.0
$\text{MgSO}_4$ (g)	0.1	0.5
$\text{CaCl}_2$ (g)	0.1	0.5
$\text{NaNO}_3$ (g)	1	5
pH	4	8
Incubation periods	3	7

### Statistical analysis

The statistical software package Minitab version 15 (Minitab Ltd., Coventry CV3 2TE, UK) was used for analyzing the experimental data. Table 2 shows the factors considered for investigation and the PB design for 12 experimental run and the experimental lipase activity. PB experimental design is based on the first order model as given in equation 1.

$$Y = \beta_0 + \sum \beta_i x_i \quad (1)$$

Where, Y is the response (enzyme activity),  $\beta_0$  is the model intercept, i is variable number,  $\beta_i$  is variable estimates and  $X_i$  are independent variables. This model describes no interaction among the factors that influence lipase production and enzyme activity. All experiments were carried out in duplicate and the averages of lipase activity were taken as responses. The variables whose confidence levels were higher than 90 % were considered that significantly influences the lipase activity.

### III. RESULTS AND DISCUSSION

*Trichoderma viride* isolated and identified based on macroscopic and microscopic observation with descriptions and illustrations in mycological literature of Domsch et al. [20] and it showed a zone of fluorescence which indicated lipolytic activity using Rhodamine B method [16]. The data on enzyme activity in table 2 was subjected to multiple linear regression analysis to estimate t-value, p-value and confidence level of each component. The results (Table 2) indicated that there was a variation of lipase production in the twelve trials in the range from 5.67 to 7.83  $\mu\text{mol}/\text{ml}/\text{min}$ . These variations reflected the importance of medium optimization to obtain higher lipase yield. On analysis of regression coefficients and t-value of 9 factors such as palm oil, yeast extract,  $\text{K}_2\text{HPO}_4$ ,  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  and  $\text{CaCl}_2$  showed a positive effect on lipase production, all other factors shown a negative effect on lipase production. When the sign of the concentration effect of the tested variable was positive, the influence of the variable upon the lipase production was greater at a high concentration, and when it was negative, the influence of the variable was greater

TABLE II  
PLACKETT-BURMAN DESIGN OF 12 RUNS FOR 11 VARIABLES ALONG WITH OBSERVED  
CONCENTRATION OF LIPASE PRODUCTION IN FERMENTATION BROTH

Run No.	Palm oil	Yeast extract	KH <sub>2</sub> PO <sub>4</sub>	K <sub>2</sub> HPO <sub>4</sub>	MgSO <sub>4</sub>	CaCl <sub>2</sub>	NaNO <sub>3</sub>	pH	Incubation periods	DV-1	DV-2	Experimental Lipase activity (μmol/ml/min)	Predicted Lipase activity (μmol/ml/min)
1	+1	-1	+1	-1	-1	-1	+1	+1	+1	-1	+1	5.67	5.558
2	+1	+1	-1	+1	-1	-1	-1	+1	+1	+1	-1	6.50	6.612
3	-1	+1	+1	-1	1	-1	-1	-1	+1	+1	+1	5.83	5.863
4	+1	-1	+1	+1	-1	+1	-1	-1	-1	+1	+1	7.30	7.333
5	+1	+1	-1	+1	+1	-1	+1	-1	-1	-1	+1	7.67	7.558
6	+1	+1	+1	-1	+1	+1	-1	+1	-1	-1	-1	7.83	7.797
7	-1	+1	+1	+1	-1	+1	+1	-1	+1	-1	-1	6.17	6.137
8	-1	-1	+1	+1	+1	-1	+1	+1	-1	+1	-1	5.67	5.782
9	-1	-1	-1	+1	+1	+1	-1	+1	+1	-1	+1	6.33	6.218
10	+1	-1	-1	-1	+1	+1	+1	-1	+1	+1	-1	6.83	6.942
11	-1	+1	-1	-1	-1	+1	+1	+1	-1	+1	+1	6.67	6.703
12	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	-1	6.00	5.967

Sign '+1' is for high concentration of variables and '-1' is for low concentration of variables; DV- Dummy variables

TABLE III  
STATISTICAL ANALYSIS OF PLACKETT-BURMAN DESIGN ON LIPASE ACTIVITY BY *Trichoderma viride*

Variable/ Term	Main Effect	Coefficients	SE Coefficients	t-value	p-value	Confidence level, %
Constant		6.5392	0.05827	112.23	0.000	100
Palm oil	0.8550	0.4275	0.05827	7.34	0.018*	98.2
Yeast Extract	0.4783	0.2392	0.05827	4.10	0.055*	94.5
KH <sub>2</sub> PO <sub>4</sub>	-0.2550	-0.1275	0.05827	-2.19	0.160	84
K <sub>2</sub> HPO <sub>4</sub>	0.1350	0.0675	0.05827	1.16	0.366	63.4
MgSO <sub>4</sub>	0.3083	0.1542	0.05827	2.65	0.118	88.2
CaCl <sub>2</sub>	0.6317	0.3158	0.05827	5.42	0.032*	96.8
NaNO <sub>3</sub>	-0.1850	-0.0925	0.05827	-1.59	0.253	74.7
pH	-0.1883	-0.0942	0.05827	-1.62	0.247	75.3
Incubation Period	-0.6350	-0.3175	0.05827	-5.45	0.032*	96.8

R-Sq = 98.67%, R-Sq(adj) = 92.67%

SE- Standard error; t – student's test; p – corresponding level of significance; \* Significant.

at a low concentration [21].

In the present work, variables with confidence levels greater than 90% were considered as significant. Palm oil, incubation period, CaCl<sub>2</sub> and yeast extract were found to be highly significant factor for lipase production (Table 3). Palm oil and yeast extract acted as carbon source and nitrogen source respectively for lipase enzyme production was reported by Treichel et al. [22]. The low incubation periods in this experiment model coincide with the data presented for Nadia et al. [23] in which the maximum lipase production was observed at 3 days of incubation periods after that it was reduced. This may be attributed to the fact that submerged technique allowed better emulsification of the lipids and also good aeration. The enzyme production was moderately enhanced in the presence of Ca<sup>2+</sup> which is important in maintaining cell wall rigidity, stabilizing oligomeric proteins and covalently bounding protein peptidoglycan complexes in the outer membrane [24]. The model in equation 2 gives the effect of significant and most important variables on lipase activity.

$$Y = 6.5392 + 0.4275 * \text{Palm oil} + 0.2392 * \text{Yeast extract} - 0.1275 * \text{KH}_2\text{PO}_4 \dots \dots \quad (2)$$

The goodness of fit model was checked by the determination of coefficient (R<sup>2</sup>) which indicated that the

TABLE IV  
ANALYSIS OF VARIANCE (ANOVA) FOR LINEAR MODEL ON  
EFFECT OF INDEPENDENT VARIABLES ON LIPASE PRODUCTION

Source	DF	SS	ASS	MSS	F	P
Main Effects	9	6.03021	6.03021	0.67002	16.45	0.059
Residual Error	2	0.08148	0.08148	0.04074		
Total	11	6.11169				

DF- Degree of freedom; SS- Sum of squares; ASS- Adjusted sum of squares; MSS- Mean sum of squares; F – Fishers's function; p – Corresponding level of significance

model could explain up to 98.67% variation of the data.

Table 4 shows the Analysis of Variance (ANOVA) for linear model on effect of independent variables on lipase production. A similar result was obtained by Baskar and Renganathan [25] in the optimization of L-asparaginase by *Aspergillus terreus* MTCC 1782.

Hymavathi et al. [26] reported that the Pareto chart of effects was plotted for identifying the factors that are important for enzyme production and shows the factors main effect estimates on the horizontal axis. The selected factors main effects are rank ordered according to their significance. The chart also shows a vertical line indicating statistical

significance ( $p = 0.10$ ). It is evident from the Pareto chart of process variables (Fig. 1) that the most important for lipase production is palm oil, followed by incubation period,  $\text{CaCl}_2$  and yeast extract.

Fig. I Effect of medium components and operating conditions on lipase activity

#### IV. CONCLUSION

The evaluation of the medium components and operating conditions for lipase production was done using the Plackett-Burman statistical method which offers efficient methodology to identify the significant variables with minimum number of experiments for extracellular lipase production by *Trichoderma viride* using palm oil as a substrate. The effect of nine medium components and operating conditions were studied and among them palm oil, incubation period,  $\text{CaCl}_2$  and yeast extract were found to be the significant variables for lipase production by *Trichoderma viride* as the percentage confidence level was more than 90%. The significant factors identified by Plackett-Burman design are considered for the next stage of the medium optimization using response surface optimization technique for the future study.

#### REFERENCES

- [1] F.Burkert, F. Maugeri, and M.I. Rodrigues, "Optimization of extracellular lipase production by *Geotrichum* sp. using factorial design." *Biores. Technol.*, vol.91, pp. 77-84, 2004.
- [2] M. Chaturvedi, M. Singh, M.R. Chugh, and S. Pandey, "Lipase production from *Bacillus subtilis* MTCC 6808 by solid state fermentation using ground nut oil cake as substrate," *Res. J. Microbiol.*, vol.5, pp. 725-730, 2010.
- [3] K. Ohnishi, Y. Yoshida, and J. Sekiguchi, "Lipase production of *Aspergillus oryzae*," *J. Ferment. Bioeng.*, vol. 77, pp.490-495, 1994.
- [4] H. Abbas, A. Hiol, V. Deyris, and L. Comeau, "Isolation and characterization of an extracellular lipase from *Mucor* sp. strain isolated from palm fruit," *Enzyme Microbial Technol.*, vol 31, pp. 968-975, 2002.
- [5] A. Hiol, M.D. Jonzo, N. Rugani, D. Druet, L. Sarda, and L.C. Comeau, "Purification and characterization of an extracellular lipase from a thermophilic *Rhizopus oryzae* strain isolated from palm fruit," *Enz. Microb. Technol.*, vol. 26, pp.421-430, 2000.
- [6] E.S. Kamimura, O. Medietta, M.I. Rodrigues, and F. Maugeri, "Studies on lipase affinity adsorption using response-surface analysis," *Biotechnol. Appl. Biochem.*, vol.33, pp.153-159, 2001.
- [7] R.Gulati, R.K. Saxena, R. Gupta, R.P. Yadav, and W. Sheba Davidson, "Parametric optimisation of *Aspergillus terreus* lipase production and its potential in ester synthesis," *Process Biochem.*, vol.35, pp. 459-464, 1999.
- [8] M.A.Kashmiri, A. Adnan, and B.W. Butt, "Production, purification and partial characterization of lipase from *Trichoderma viride*," *Afr. J. Biotechnol.*, vol. 5, pp. 878-882, 2006.
- [9] E.M. Rajesh, R. Arthe, R. Rajendran, C. Balakumar, N. Pradeepa and S. Anitha, "Investigation of lipase production by *Trichoderma reesei* and optimization of production parameters," *EJEA Che.*, vol. 9, pp. 1177-1189, 2010.
- [10] M.M.D. Maia, A. Heasley, M.M. Camargo de Morais, E.H.M. Melo, M.A. Jr. Morais, W.M. Ledingham, and J.L. Lima Filho, "Effect of culture conditions on lipase production by *Fusarium solani* in batch fermentation," *Biores. Technol.*, vol. 76, pp. 23-27, 2001.
- [11] J.A. Rodriguez, T. Bhagnagar, S. Roussos, J. Cordova, and J. Baratti, "Improving lipase production by nutrient source modification using *Rhizopus homothallicus* cultured in solid state fermentation," *Process Biochem.*, vol.41, pp. 2264-2269, 2006.
- [12] D.V. Desmukh, and P.R. Puranik, 2010, "Application of Plackett-Burman design to evaluate media components affecting antibacterial activity of alkaliphilic cyanobacteria isolated from Lonar Lake," *Turk. J. Biochem.*, vol. 35, no. 2, pp. 114-120, 2010.
- [13] G.M. Maciel, L.P.S. Vandenberghe, C.W.I. Haminiuk, and C.R. Soccol. "Xylanase production by *Aspergillus niger* LPB 326 in solid-State fermentation using statistical experimental designs. Xylanase production by *A. niger* in SSF," *Food Technol. Biotechnol.*, vol. 46, pp. 183-189, 2008.
- [14] S.B. Imandi, S.K. Karanam, and H.R. Garapati, "Optimization of media constituents for the production of lipase in solid state fermentation by *Yarrowia lipolytica* from palm Kernal cake (*Elaeis guineensis*)," *Adv. Biosci. Biotechnol.*, vol.1, pp. 115-121, 2010.
- [15] D.A. Akano, and O.O. Atanda, "The present level of aflatoxin in Nigerian groundnut cake (kulikuli)," *Lett. Applied Microbiol.*, vol. 10, pp.187-189, 1990.
- [16] M.G. Valeria, K. Lima-Nadia, M. Maria-Inez, A. Sarquis-David, P. Mitchell-Luiz, and D. Ramos-Jose, "Production of Lipase by *Penicillium aurantiogriseum*," *Food Technol. Biotechnol.*, vol. 41, pp. 105-110, 2003.
- [17] S. Fadiloglu, and Z. Soylemez, "Kinetics of lipase-catalyzed hydrolysis of olive oil," *Food Res. Intern.*, vol. 30, pp. 171-175, 1997.
- [18] F. Pedreschi, K. Kaack, and K. Granby, "The effect of asparaginase on acrylamide formation in French fries," *Food Chem.*, vol.109, pp.386-392, 2008.
- [19] N.A. Soliman, M.B. Mahmoud, and Y.R. Abdel-Fattah, "Polyglutamic acid production by *Bacillus* sp. SAB-26: Application of Plackett-Burman experimental design to evaluate culture requirements," *Appl. Microbiol. Biotechnol.*, vol. 69, pp. 259-267, 2005.
- [20] K.W. Domsch, W. Gams and T.H. Anderson, "Compendium of Soil Fungi," Academic Press, London, pp: 859. 1980.
- [21] G. Dhanya, S. Swetha, N.K. Madhavan, S. Rajeev and P. Ashok, "Response surface methodology for the optimization of alpha amylase production by *Bacillus amyloliquefaciens*," *Biores. Technol.*, vol. 99, pp. 4597-4602, 2008.
- [22] H. Treichel, D. de Oliveira, M. A. Mazutti, M. D. Luccio, and J. Vladimir Oliveira, "A review on microbial lipases production food," *Bioprocess Technol.*, vol.3, pp.182-196, 2010.
- [23] N. Nadia, Z.A. Nehad, A.E. Elsayed, M.A. Essam, and M.A. Hanan, "Optimization of lipase synthesis by *Mucor racemosus* production in a triple impeller bioreactor," *Malaysian J. Microbiol.*, vol. 6, pp. 7-15, 2010.
- [24] D. Maccio, A. Fabra, and S. Castro, "Acidity and calcium interaction affect the growth of *Bradyrhizobium* sp. and attachment to peanut roots," *Soil Biol. Biochem.*, vol. 34, pp.201-208, 2002.
- [25] G. Baskar, and S. Renganathan, "Statistical screening of process variables for the production of L-asparaginase from cornflour by *Aspergillus terreus* MTCC 1782 in submerged fermentation," *Indian J. Sci. Technol.*, vol. 2, pp. 45-48, 2009.
- [26] M. Hymavathi, T. Sathish, P. Brahmaiah, and R. S. Prakasham, "Impact of carbon and nitrogen sources on L-asparaginase production by isolated *Bacillus circulans* (MTCC 8574): Application of saturated Plackett-Burman design," *Chem. Biochem. Eng. Q.*, vol. 24, no. 4, pp. 473-480, 2010.