

# Complexity of Multivalued Maps

David Sherwell and Vivien Visaya

**Abstract**—We consider the topological entropy of maps that in general, cannot be described by one-dimensional dynamics. In particular, we show that for a multivalued map  $F$  generated by single-valued maps, the topological entropy of any of the single-value map bounds the topological entropy of  $F$  from below.

**Keywords**—Multivalued maps, Topological entropy, Selectors

## I. INTRODUCTION

One measure of the complexity of a system is the positivity of its topological entropy. The topological entropy for single-value maps is well-known [1], while the case for multivalued maps, in general, is not widely studied. However, it is worth noting the works of [2] and [3] on lower bounds for topological entropy using the Conley index theory, and works of [4] and [5] on branch image entropy defined from the backward iterates of a non-invertible map.

Because many systems encountered in practice cannot be described by one-dimensional dynamics (e.g. in population dynamics, where many observed units can occupy a state, and are mapped to different states [6]), it is useful to consider multivalued maps. We define in Section II the topological entropy of such maps, and give conditions in bounding their topological entropy from below in Section III. We give two examples in Section IV, particularly focusing on the map studied in [6].

## II. PRELIMINARIES

We give the standard definition of the topological entropy due to Bowen [8]. Let  $(X, d)$  be a compact metric space with distance  $d$  and let  $f : X \rightarrow X$  be continuous. For any integer  $\ell \geq 1$ , define the distance function  $d_\ell : X \times X \rightarrow \mathbb{R}_{\geq 0}$  by

$$d_\ell(x, y) = \max_{0 \leq j < \ell} d(f^j(x), f^j(y)).$$

**Definition 1:** A finite set  $E \subset X$  is called  $(\ell, \delta)$ -separated if  $d_\ell(x, y) \geq \delta$  for all  $x, y \in E$ . Moreover, if  $E$  has the maximal cardinality among all the  $(\ell, \delta)$ -separated sets, then  $E$  is called a *maximal  $(\ell, \delta)$ -separated set*.

**Definition 2:** The topological entropy of  $f$  is given by

$$h_{\text{top}}(f) = \lim_{\delta \rightarrow 0} \limsup_{\ell \rightarrow \infty} \frac{\log s_f(\ell, \delta)}{\ell},$$

where  $s_f(\ell, \delta)$  is the cardinality of the maximal  $(\ell, \delta)$ -separated set for  $f$ .

**Definition 3:** [9] Let  $X$  and  $Y$  be arbitrary sets. A *multivalued map  $F$  from  $X$  to  $Y$* , denoted by  $F : X \rightrightarrows Y$ , is such that  $F(x)$  is assigned a set  $Y_x \subset Y$  for all  $x \in X$ . Let

D. Sherwell and V. Visaya are with the School of Computational and Applied Mathematics, University of the Witwatersrand, Johannesburg, South Africa

1.  $\text{dom}(F) = \{x \in X \mid F(x) \neq \emptyset\}$ ,
2.  $F(X') = \bigcup \{F(x) \mid x \in X'\}$  for  $X' \subset X$ .

A single-valued map  $f : X'(\subset X) \rightarrow Y$  is called a *selector for  $F$  over  $X'$*  if  $g(x) \in F(x)$  for all  $x \in X'$ .

Let  $(X, d)$  be a compact metric space. Let  $F : X \rightrightarrows X$  be a multivalued map,  $g : X \rightarrow X$  be a continuous map,  $\text{Sel}^0(F)$  be the set of all continuous selectors for  $F$ , and  $\{u_m\}_{m=0}^M$  be a sequence of  $M+1$  points such that

$$u_{m+1} \in F(u_m) \quad \text{for } m = \{0, 1, \dots, M-1\}.$$

Assume that the graph of  $F$  can be viewed as a union of its continuous selectors, i.e.

$$F(x) = \bigcup \{g(x) \mid g \in \text{Sel}^0(F)\} \quad \forall x \in \text{dom}(F).$$

**Definition 4:** Let  $(X, d)$  be a compact metric space and let  $F : X \rightrightarrows X$  be a multivalued map. Denote the *set of partial orbits of  $F$  of length  $\ell$*  by

$$U_\ell = \{u = \{u_i, u_{i+1}, \dots, u_{i+\ell}\}\}_{i=0}^{M-\ell} \subset X^{\ell+1},$$

where  $u_{i+j} \in F(u_{i+j-1})$  for all  $1 \leq j \leq \ell$ . A set  $S \subset U_\ell$  is called a  $(\ell, \delta)$ -separated set for  $F$  if for any  $u, u' \in S$ ,  $d(u, u') \geq \delta$ .

We extend the notion of topological entropy to multivalued maps.

**Definition 5:** Let  $S_F(\ell, \delta)$  be the maximal  $(\ell, \delta)$ -separated set for  $F$  with cardinality  $s_F(\ell, \delta)$ . We define the topological entropy of  $F$  by

$$h_{\text{top}}(F) = \lim_{\delta \rightarrow 0} \limsup_{\ell \rightarrow \infty} \frac{\log s_F(\ell, \delta)}{\ell}.$$

**Definition 6:** [10] Let  $X$  be any set and let  $F : X \rightrightarrows X$  be a multivalued map. Suppose  $\mathcal{P} = \{P_0, P_1, \dots, P_{L-1}\}$  is a partition of  $X$  into  $L$  disjoint regions and suppose that the intersection of any element of  $\mathcal{P}$  with the image under  $F$  of another is either itself or is empty. The structure of  $\mathcal{P}$  can be described a transition matrix  $T = (T_{ij})$  defined by

$$T_{ij} = \begin{cases} 1 & \text{if } P_j \cap F(P_i) \neq \emptyset \\ 0 & \text{otherwise.} \end{cases}$$

$T$  is called the transition matrix for  $F$ .

From the definition above, if  $x \in P_i$ , then  $F(x) \in P_j$ .

**Definition 7:** [11] Given a transition matrix  $T = (T_{ij})$  with  $T_{ij} \in \{0, 1\}$ , let

$$\Sigma_T^+ = \{w = (w_0 w_1 \dots) \mid T_{w_m, w_{m+1}} = 1 \ (\forall m \geq 0)\}.$$

The shift map  $\sigma_T : \Sigma_T^+ \rightarrow \Sigma_T^+$  is such that  $(\sigma_T(w))_m = w_{m+1}$  ( $\forall m \geq 0$ ). The pair  $(\sigma_T, \Sigma_T^+)$  is called the *one-sided subshift of finite type (SFT) for the matrix  $T$* .

**Theorem 1:** [11] Let  $T$  be a transition matrix and let  $\sigma_T : \Sigma_T^+ \rightarrow \Sigma_T^+$  be the associated SFT. Then

$$h_{\text{top}}(\sigma_T) = \ln(\lambda_{\max})$$

where  $\lambda_{\max}$  is the maximum eigenvalue of  $T$ .

### III. RESULTS

**Theorem 2:** Let  $(X, d)$  be a compact metric space and let  $F : X \rightrightarrows X$  be a multivalued map. For any continuous selector  $g : X \rightarrow X$  of  $F$ , the following inequality holds

$$h_{\text{top}}(F) \geq \sup\{h_{\text{top}}(g) | g \in \text{Sel}^0(F)\}.$$

**Proof.** Take an arbitrary  $g \in \text{Sel}^0(F)$ . Let  $S_g$  be the maximal  $(k, \delta)$ -separated set for  $g$ . We show that the maximal  $(\ell, \delta)$ -separated set for any  $g \in \text{Sel}^0(F)$  is a  $(\ell, \delta)$ -separated set for  $F$ . For all  $x \in S_g$ , denote by

$$S_g^\ell = \{x_g^\ell = (x, g(x), \dots, g^\ell(x))\} \subset U_\ell$$

the subset of the partial orbits of  $g$  of length  $\ell$ . Clearly,  $\#S_g = \#S_g^\ell$ . Since

$$d_\ell(u, v) = \max_{0 \leq j < \ell} (u_j, v_j)$$

for  $u, v \in X^{\ell+1}$  and  $u_j, v_j \in X$ , then for any two distinct  $x, y \in S_g$ ,

$$d(x, y) \geq \delta \Rightarrow d_\ell(x_g^\ell, y_g^\ell) \geq \delta,$$

where  $x_g^\ell, y_g^\ell \in S_g^\ell$ . Thus,  $S_g^\ell$  is a  $(\ell, \delta)$ -separated set for  $F$ . Note however that  $S_g$  may not be maximal for  $F$  so

$$s_F(\ell, \delta) \geq s_g(\ell, \delta).$$

Passing to the limits, we establish the claim.

**Corollary 1:** Let  $(X, d)$  and  $F$  be as in Theorem 2. If  $g_1, g_2 : X \rightarrow X$  are continuous selectors of  $F$ , then  $g = g_1 \circ g_2$  is also a continuous selector of  $F$ , and that

$$h_{\text{top}}(F) \geq h_{\text{top}}(g).$$

### IV. EXAMPLES

**Example A.** Consider the case on the interval  $X = [0, 1]$ , where the multivalued map  $F : X \rightrightarrows X$  is generated by two self-maps

$$g_1 = \begin{cases} 2x & 0 \leq x \leq \frac{1}{2} \\ 2-2x & \frac{1}{2} \leq x \leq 1 \end{cases}$$

and

$$g_2 = \begin{cases} 2ax & 0 \leq x \leq \frac{1}{2} \\ 2a-2ax & \frac{1}{2} \leq x \leq 1 \end{cases}, \quad 1/2 \leq a < 1.$$

It is well-known that the topological entropy of  $g_1$  and  $g_2$  are  $\log 2$  and  $\log 2a$  respectively. Hence,

$$h_{\text{top}}(F) \geq \sup\{\log 2, \log 2a\}.$$

**Example B.** We consider the method in analysing longitudinal data, studied in [6]. Longitudinal data is simply a repeated measurement of the same variables (observed units) in time. Let  $\mathbb{N}_0$  be the set of non-negative integers, and let the integer




Fig. 1. The graph of the multivalued map  $F$  as a union of  $g_1$  and  $g_2$ .

$n \geq 1$ . Let  $I_n = \{0, 1, \dots, n-1\}$ ,  $\mathcal{C}_n = \mathcal{P}(I_n)$  be the power set of  $I_n$ , and  $x_i^*$  be such that

$$x_i^* = \begin{cases} 1 & \text{if } x_i = 0 \\ 0 & \text{if } x_i = 1. \end{cases}$$

Let

$$\Gamma_2^n = \{(x_j)_{j=0}^{n-1} : x_j \in \{0, 1\}\}$$

and let

$$\Gamma_n^n = \{(i_j)_{j=0}^{n-1} : i_j \in I_n, i_j' \text{ s distinct}\},$$

where the subscripts 2 and  $n$  are the cardinalities of the sets  $\{0, 1\}$  and  $I_n$ , respectively. Let

$$\begin{aligned} \mathcal{S}_n &= \{p = (x, y) : x \in \Gamma_2^n, y \in \Gamma_n^n\} \\ &= \Gamma_2^n \times \Gamma_n^n. \end{aligned}$$

Given  $n$ , we have  $|\Gamma_2^n| = 2^n$ ,  $|\Gamma_n^n| = n!$ , and so  $\mathcal{S}_n$  is a finite space composed of  $L = 2^n \times n!$  states.

**Definition 8:** Let  $\Delta \in \mathcal{C}_n$  and let  $j \in \Delta$ . The *change map*  $\phi_\Delta : \mathcal{S}_n \rightarrow \mathcal{S}_n$  is defined by

$$\phi_\Delta(x_0 x_1 \dots x_j \dots x_{n-1}, y) = (x_0 x_1 \dots x_j^* \dots x_{n-1}, y).$$

The *jump map*  $\phi_j : \mathcal{S}_n \rightarrow \mathcal{S}_n$  is defined by

$$\phi_j(x_0 x_1 \dots x_j \dots x_{n-1}, i_0 i_1 \dots i_j \dots i_{n-1}) = (x_0 x_1 \dots x_{j-1} x_{j+1} \dots x_{n-1} x_j, i_0 i_1 \dots i_{j-1} i_{j+1} \dots i_{n-1} i_j).$$

Let  $j, j' \in \Delta$ . Then  $x_j$  and  $x_{j'}$  both change values under  $\phi_\Delta$ . If  $j < j'$ , then  $\phi_j$  is first applied to  $j'$ . That is,

$$\begin{aligned} \phi_j(x_0 x_1 \dots x_j \dots x_{j'} \dots x_{n-1}, i_0 i_1 \dots i_j \dots i_{j'} \dots i_{n-1}) &= \\ (x_0 x_1 \dots x_{j-1} x_{j+1} \dots x_{j'-1} x_{j'+1} \dots x_{n-1} x_j x_{j'}, & \\ i_0 i_1 \dots i_{j-1} i_{j+1} \dots i_{n-1} i_j i_{j'}) &. \end{aligned}$$

Consider an observed unit (variable)  $k$  in the longitudinal data, and questionnaire  $\mathcal{Q}$  of  $n \geq 1$  questions. Denote by

$$\begin{aligned} \mathcal{Q}_t^k &= \{Q_{i_0 t}^k, Q_{i_1 t}^k, \dots, Q_{i_{(n-1)t}^k}^k\} \quad (*) \\ &:= \{Q_{i_0}, Q_{i_1}, \dots, Q_{i_{n-1}}\}_t^k, \quad i_j, i_{jt} \in I_n \end{aligned}$$

any reordering of questions of unit  $k$  at time  $t$ ,

$$\mathcal{A}_t^k = \{x_{0t}^k, x_{1t}^k, \dots, x_{(n-1)t}^k\} := \{x_0, x_1, \dots, x_{n-1}\}_t^k$$

the set of coded answers to  $\mathcal{Q}_t^k$ ,

$$x_t^k = x_{0t}^k x_{1t}^k \dots x_{(n-1)t}^k := (x_0 x_1 \dots x_{n-1})_t^k = (x_{jt}^k)_{j=0}^{n-1}$$

the concatenation of elements of  $\mathcal{A}_t^k$ , and

$$\begin{aligned} y_t^k &= i_{0t}^k i_{1t}^k \dots i_{(n-1)t}^k := (i_0 i_1 \dots i_{n-1})_t^k = (i_{jt}^k)_{j=0}^{n-1} \\ &\text{the concatenation of indices in } \mathcal{Q}_t^k. \end{aligned}$$

**Definition 9:** For each observed unit  $k$ , define  $\Delta_t^k \in \mathcal{C}_n$  by  $\Delta_t^k = \{j : i_j \text{ is a question index of unit } k \text{ at time } t \text{ that changes answer value at } t+1, \text{ ordered in ascending order, as in } (*)\}$ .

**Definition 10:** Fix observed unit  $k$  and let  $p_0^k$  be the initial state of  $k$ . Define the map  $\varphi : (\mathbb{N}_0, \mathcal{C}_n, \mathcal{S}_n) \rightarrow \mathcal{S}_n$  such that

$$\begin{aligned}\varphi(t, \Delta_t^k, p_t^k) &= \varphi_{[\Delta_t^k]}(p_t^k) \\ &= (\phi_j \circ \phi_{\Delta_t^k})(p_t^k) \\ &= p_{t+1}^k.\end{aligned}$$

The set  $\Delta_t^k \in \mathcal{C}_n$  is given by the longitudinal data for all  $t$ . The action of  $\phi_{\Delta_t^k}$  is data dependent while  $\phi_j$  is a strictly deterministic reordering of the position of the question indices and answer values in  $\Delta_t^k$ . At any time  $t$ , we can always trace the answer to its corresponding question such that the coded answer value  $x_j$  corresponds to question  $i_j$ . For each  $k$ , the nonautonomous map  $\varphi_{[\Delta_t^k]}$  displaces the most frequently changing answers to the right, while slowly changing answers displace to the left.

Given  $k$ , if  $p_t^k \in p$  and  $p_{t+1}^k \in p'$ , then we say that there is a transition from  $p$  to  $p'$  under  $\Delta_t \in \mathcal{C}_n$ . A *self-transition* is under the empty set  $\Delta = \emptyset$  (i.e. there is no change in answer). A transition from  $p$  to  $p'$ , and from  $p'$  to  $p$ , under the same set  $\Delta$  is *reversible*. A way to visualize the state transitions of unit  $k$  in  $\mathcal{S}_n$  is by a *directed graph* (digraph)  $\mathcal{G}$  whose vertices are points in  $\mathcal{S}_n$ , with an edge from  $p$  to  $p'$  if there is a transition from  $p$  to  $p'$ . A *path of length  $m$*  in  $\mathcal{G}$  is a sequence of vertices  $v_0, v_1, \dots, v_m$  such that there is a directed edge from  $v_j$  to  $v_{j+1}$ .




Fig. 2. All possible transitions between states in (a)  $\mathcal{S}_1$  and (b)  $\mathcal{S}_2$ .

Fig. 2 illustrates all possible transitions between states in  $\mathcal{S}_1$  and  $\mathcal{S}_2$ . Transitions alternating between states 1 and 2 in Fig. 2(a) denote alternating answer between 0 and 1, to question 0. On the other hand, transitions alternating between states 1 and 2 in Fig. 2(b) denote constant answer=0 to question 1, and alternating answer between 0 and 1, to question 0. Because question 1 has constant answer, it is positioned on the left of question order '10'.

Consider our state space  $\mathcal{S}_n$ . In [6], the map  $\varphi_{[\Delta_t^k]}$  is constructed for each observed unit  $k$ . By defining a multivalued map  $F : \mathcal{S}_n \rightrightarrows \mathcal{S}_n$ , we can describe all possible paths of observed units in  $\mathcal{S}_n$ . Denote by  $\varphi_{[\Delta]} : \mathcal{S}_n \rightarrow \mathcal{S}_n$  the case where  $\Delta_t$  in Definition 9 is constant. Let

$$G = \{\varphi_{[\Delta]} : \Delta \in \mathcal{C}_n\}.$$

It is clear that  $\varphi_{[\Delta]}$  is a selector for  $F$  over  $\mathcal{S}_n$ , and is continuous if  $\mathcal{S}_n$  is endowed with the discrete metric. Define

$$F(p) = \{p' : g(p) = p', g \in G\}.$$

Observe that  $|F(p)| = 2^n$ , i.e.  $F$  is a 1 to  $2^n$  map. Define

$$G_t = \{g_{t-1} \circ \dots \circ g_{t_0} : \mathcal{S}_n \rightarrow \mathcal{S}_n | g_{t_i} \in G\}.$$

For  $g_{t_i} = \varphi_{[\Delta_t^k]}$  for all  $t \geq 1$ , we define the orbit of  $p \in \mathcal{S}_n$  under  $F$  by

$$\begin{aligned}\mathcal{O}_F(p_0^k) &= \{p_t^k\}_{t \geq 0} \\ &= \{G_t(p_0^k)\}_{t \geq 1},\end{aligned}$$

where under  $F$ , the state  $p_{t+1}^k \in F(p_t^k)$ . The multivalued map  $F$  can be interpreted as a digraph whose  $N = n!2^n$  vertices are the points in  $\mathcal{S}_n$ , and edge  $p \rightarrow p'$  if  $p' \in F(p)$ . Equivalently,  $F$  can be defined as a square matrix of size  $N$ . Orbits of the nonautonomous map  $\varphi_{[\Delta_t^k]}$  respect paths in the digraph  $F$ .

Consider  $\mathcal{S}_n$  as a union of  $L$  disjoint states labeled  $i = 1, 2, \dots, N = 2^n n!$ , i.e.,

$$\mathcal{S}_n = \bigcup_{i=1}^N s_i.$$

A way of labeling  $s_i$  is via the map

$$\psi : \mathcal{S}_n \rightarrow \{1, 2, \dots, N\}, \quad s_i \mapsto \psi(s_i) = i.$$

**Definition 11:** Let  $n \geq 1$ ,  $V = \{1, 2, \dots, N\}$ , and  $i, j \in V$ . For  $p, p' \in \mathcal{S}_n$ , let  $\psi(p) = i$ , and  $\psi(p') = j$ . The transition matrix for  $F$  is denoted by  $T_n^{(F)} = (T_{ij}^{(F)})$ , where

$$T_{ij}^{(F)} = \begin{cases} 1 & \text{if } p' \in F(p) \\ 0 & \text{otherwise.} \end{cases}$$

**Remarks.**

- (i) From Fig. 2, the transition matrices for  $n = 1$  and  $n = 2$  are

$$T_1^{(F)} = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \quad T_2^{(F)} = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$

In [6], the analysis of longitudinal data from household units is studied in  $\mathcal{S}_3$ . We do not give the matrix here, however, we illustrate in the Appendix A all possible transitions in  $\mathcal{S}_3$ .

- (ii) Because  $F(p)$  is a 1 to  $2^n$  map, then by properties of non-negative matrices,  $T_n^{(F)}$  has  $\lambda_{max} = 2^n$ . By Theorem 1,  $h_{\text{top}}(\sigma_{T_n^{(F)}}) = \ln(2^n)$ .
- (iii) For constant  $\Delta$ , we have

$$h_{\text{top}}(\varphi_{[\Delta]}) = 0.$$

We illustrate in Appendix B the case where  $\Delta = I_3$ , i.e. where all three answers are constantly changing. The transition from any point  $p$  to  $p' = \varphi_{[\Delta]}(p)$  is reversible. For example,  $\varphi_{[I_3]}$  takes state 1 to state 48, and state 48 back to state 1. Note that the strict inequality in Theorem 2 is satisfied.

## V. CONCLUSIONS


We have presented a definition of the topological entropy of multivalued maps. For the case where the multivalued map  $F$  is generated by single-valued maps, then we are able to give a lower bound for the complexity of  $F$ . If we can find a selector  $g$  for a multivalued map  $F$ , and find that the topological entropy of  $g$  is positive, then we can say that  $F$  is at least as complicated as  $g$ . In the case that we can define a partition for the state space of  $F$  (as in Example B), then we can encode  $F$  as a transition matrix  $T$ , and analysis of  $F$  is through the subshift associated to  $T$ . Then any selector, or composition of selectors of  $F$ , has complexity bounded by the complexity of  $F$ . We note that this paper does not suggest a general techniques for computing selectors, but will be persued in a following paper.

## ACKNOWLEDGMENT

This work was supported by a University of the Witwatersrand URC Grant. V. Visaya would also like to thank Prof. Hiroshi Kokubu for useful discussion on multivalued map during her stay in Japan.

## REFERENCES

- [1] Alsedà, J. Llibre, & M. Misiurewicz, *Combinatorial Dynamics and Entropy in Dimension One* (2nd. ed.), World Scientific, 2000.
- [2] A. Baker, Lower bounds on entropy via the Conley index with applications to time series, *Topology and Its Applications* 120 (2000) 333-354.
- [3] S. Day, R. Frongillo, and R. Trevino, Algorithms for Rigorous bound and symbolic dynamics
- [4] M. Hurley, On topological entropy of maps, *Ergodic Th. & Dynam. Sys.* 15 (1995) 557-568.
- [5] Z. Nitecki, Preimage entropy for mappings, *International Journal of Bifurcation and Chaos* 9 (1999) 1815-1843.
- [6] C.L. Mberi Kimpolo, *Deterministic Dynamics in Questionnaires in the Social Sciences*, Ph.D. Thesis, University of the Witwatersrand (2010), Supervisor D. Sherwell.
- [7] KS2 C. L. Mberi Kimpolo, D. Sherwell, et al., *Orbit Theory: Analysis of Longitudinal Data by Visualization of Fitness States*, submitted.
- [8] R. Bowen, Entropy for group endomorphisms and homogeneous spaces, *Trans. AMS* **153** (1971) 509-510.
- [9] M. Mrozek, Topological invariants, multivalued maps and computer assisted proofs in dynamics, *Computers Math. Applic.* 32 (1996) 83-104.
- [10] R. Gilmore and M. Lefranc, *The Topology of Chaos*, John Wiley & Sons, Inc. (2002).
- [11] C. Robinson, *Dynamical Systems: Stability, Symbolic Dynamics, & Chaos*, C.R.C. Press, 1994.

APPENDIX A  
TRANSITIONS IN  $\mathcal{S}_3$ APPENDIX B  
REVERSIBLE TRANSITIONS UNDER CONSTANT  $\Delta = I_3$ .