Cereals' Products with Red Grape and Walnut Extracts as Functional Foods for Prevention of Kidney Dysfunction

Sahar Y. Al-Okbi, Doha A. Mohamed, Thanaa E. Hamed, and Ahmed Ms Hussein

Abstract-In the present research, two nutraceuticals made from red grape and walnut that showed previously to improve kidney dysfunction were incorporated separately into functional foods' bread made from barley and rice bran. The functional foods were evaluated in rats in which chronic renal failure was induced through feeding diet rich in adenine and phosphate (APD). The evaluation based on assessing kidney function, oxidative stress, inflammatory biomarkers and body weight gain. Results showed induction of chronic kidney failure reflected in significant increase in plasma urea, creatinine, malondialdehyde, tumor necrosis factor- α and low density lipoprotein cholesterol along with significant reduction of plasma albumin, and total antioxidant and creatinine clearance and body weight gain on feeding APD compared to control healthy group. Feeding the functional foods produced amelioration in the different biochemical parameters and body weight gain indicating improvement in kidney function.

Keywords-Functional food, kidney dysfunction, rats.

I. INTRODUCTION

XIDATIVE stress and inflammation are involved as risk factors of chronic non communicable diseases including renal dysfunction [1], [2]. Increased level of malondialdehyde was reported during renal damage [3]. Both native and oxidized forms of low density lipoprotein-cholesterol (LDL-Ch) may be involved in the glomerular damage mediated by oxidative stress [4]. Inflammatory lesions caused by oxidative stress participate in the glomerular damage [2]. So, antioxidant and anti-inflammatory nutraceuticals could play an important role in the protection from incidence of chronic renal failure. On the other hand some beverages made from barley have been used in Egypt as Folk medicine to alleviate kidney dysfunction. Phytate, β -glucan, tocopherols and tocotrienols were reported to present in barley seeds [5], referring to its potential hypocholesterolemic, antiinflammatory and antioxidant effect. Also, previously rice bran was shown to contain different bioactive constituents that possess antioxidant and anti-inflammatory activity and LDL-Ch lowering effect [6]. Policosanol, gamma oryzanol,

Sahar Y. Al-Okbi is with the Food Sciences and Nutrition Department, National Research Centre, Dokki, 12622, Cairo, Egypt (corresponding author phone: 00201003785152; e-mail: S_Y_alokbi@hotmail.com).

Doha A. Mohamed and Thanaa E. Hamed are with the Food Sciences and Nutrition Department, National Research Centre, Dokki, 12622, Cairo, Egypt.

Ahmed MS Hussein is with the Food Technology Department, National Research Centre, Dokki, 12622, Cairo, Egypt.

phytosterols, unsaturated fatty acids, phenolic compounds and dietary fibers [6], [7] are examples of such active ingredients that present in rice bran. In a previous work, nutraceuticals prepared from red grape and walnut were proved to improve kidney dysfunction [8]. Red grape with seeds and walnut were reported to contain total phenolic as 6.4 and 97.42g gallic acid equivalent/100g alcohol extract, linolenic as 3.71 and 6.1% of total fatty acids and 15.868 and 11.03 phytosterol % of unsaponifiable portion [8]. Phytosterol, phenolic compounds and omega 3 fatty acids such as linolenic are reported to possess antioxidant, anti-inflammatory and hypocholesterolemic effect [9]-[11]. So, walnut and red grape are important sources of nutraceuticals. Incorporation of barley or rice bran together with such nutraceuticals into functional foods may lead to highly efficient products that might have health effect towards chronic renal dysfunction.

The aim of the present research was preparation and chemical, sensory and biological evaluation of functional foods for protection from chronic renal failure utilizing efficient nutraceuticals from previous research and cereals rich in antioxidant and anti-inflammatory active constituents.

II. MATERIALS AND METHODS

A. Materials

- 1. Flours: Wheat grains (Giza 168) and naked barley (Hordeum vulgare variety Giza 129) were obtained from Agric. Res. Center, Giza, Egypt. Stabilized rice bran (RB) was supplemented by International Trade and Marketing, Egypt.
- 2. Nutraceuticals' Sources: Red grape (with seeds) and walnut were purchased from local markets, Cairo, Egypt.
- 3. Main Chemical: Adenine was obtained from Sigma (Sigma-Aldrich Co., St. Louis, Missouri, USA).
- 4. Animals: Male Sprague–Dawley rats of 215.8g average body weight were used in the present study. Animals were obtained from animal house of National Research Centre, Cairo, Egypt. They were kept individually in metabolic stainless steel cages; water and food were given ad-libtium all over the experiment.

B.Methods

1. Preparation of Flour: Barley and wheat grain were separately cleaned, tempered (15% moisture) and milled (Quadrumat Junior flour mill) to 100 % extraction flour.

Whole meal wheat flour (WWF) was well blended with rice bran in the ratio of 70: 30. The samples were stored in airtight containers and kept at 5-7°C until used.

- 2. Preparation of Plant Materials: Fresh red grape with seed was washed by tap water and cut into small pieces and dried in an air-circulated oven at 40°C. Both dried grape and walnut were separately reduced into powder.
- 3. Preparation of Plant Extracts: The dried powder of grape and walnut were separately extracted by continuous extraction apparatus for extraction by petroleum ether (40-60°C) followed by ethanol. The solvent of each extract was removed by evaporation under reduced pressure.
- 4. Preparation of Extracts Mixtures (Nutraceuticals): Ethanol and petroleum ether extracts of each plant was mixed together in the ratio of their presence in the whole plant.
- 5. Preparation of Bread: Flat bread was prepared from 100% whole barley flour (WBF) with or without incorporation of grape nutraceutical as functional food I and control, respectively. Balady bread was prepared from 70% WWF and 30% RB with or without walnut nutraceutical as functional food II and control, respectively.
- 6. Sensory Evaluation of Bread: Each functional food along with its corresponding control was sensory evaluated by 10 trained panelists. Texture, color, odor, taste and appearance were the criteria for evaluation of flat bread. Sensory attributes used for organoleptic assessment of balady bread were appearance, separation of layers, roundness, distribution of crumb, crust color, taste, and odor. Sensory attribute that verify > 50% of the score was deemed as accepted.
- 7. Determination of Proximate Composition of Food Products: Moisture, crude fiber, ash, protein and fat contents of the breads were determined according to AOAC [12]. Total carbohydrate was calculated by difference.
- 8. Preparation of Experimental Diets: In the present study, renal dysfunction was induced in rats by feeding them high adenine (0.75%) high phosphate (2.4%) diet for 10 days followed by high adenine (0.75%) diet for another consecutive 11 days (APD). The different Types of bread were dried and reduced into powder for preparation of experimental diets as shown in Table I.
- 9. Design of Animal Experiment: Forty-two rats were divided into two groups. Group one served as normal control (n= 6 rats) and fed on balanced diet for three weeks. Group two (n= 36 rats) and fed on high phosphate-high adenine diet for 10 days followed by high adenine diet for another 11 days to induce chronic renal failure (APD). During the experiment, body weight and food intake were recorded weekly. At the end of the three weeks (first stage), total food intake, body weight gain and food efficiency ratio (Body weight gain/total food intake) were calculated. Also, 24 hour urine was collected from the control normal rats and six rats from APD group. Blood samples were obtained from the same fasted animals. Heparin was used as an anticoagulant and

plasma was separated by centrifugation at 3500rpm for 10 min. Plasma malondialdehyde (MDA) was determined according to Satoh [13] as indicator of lipid peroxidation and oxidative stress. Plasma total antioxidant capacity (TAC) as an antioxidant biomarker was assessed according to Koracevic et al. [14]. Plasma tumor necrosis factor- α (TNF- α), as inflammatory biomarker, was determined according to Stepaniak et al. [15]. Plasma creatinine [16] and urea [17] were estimated as indicator of kidney function. Plasma albumin [18], plasma total cholesterol (T.Ch) [19] and LDL-Ch [20] were also determined which may serve as extra indicators of kidney function. Creatinine was determined in the collected 24h urine [16] for calculation of creatinine clearance. In the second stage, residual APD rats (n 30) were divided into five groups. Rats of group one were fed on balanced diet, rats of group two and three were fed on balanced diet containing barley bread and rice bran bread, respectively. Rats of groups four and five were fed on balanced diet containing functional food I and II, respectively. This stage of experiment continued for a month. Nutritional parameters, biochemical analysis and creatinine clearance were again assessed as mentioned previously in the end of the first stage.

10. Statistical Analysis: All the results were expressed as mean \pm SE. Sensory attributes and data of nutritional parameters of the 1st stage of animal experiment were analyzed statistically using t-student test. One-way analysis of variance ANOVA followed by Duncan's test were used for comparison of all other parameters of the different experimental groups. In all cases p<0.05 was used as the criterion of statistical significance.

III. RESULTS

Proximate composition of the prepared functional foods (Table II) showed that the values of both functional food I and II were more or less similar to their respective control bread except for slight change in ash and carbohydrate. In functional food II, fat was slightly more than that of the respective control.

Sensory parameters of functional foods seen in Table III and IV showed no significant changes in all sensory attributes when compared with their respective control. All bread samples showed to be accepted by panelists.

The determined biochemical parameters of the different experimental groups are shown in Table V. APD group showed a significant elevation in plasma creatinine and urea levels as indicator of kidney dysfunction compared with normal control. There were no significant differences in plasma total cholesterol level, while LDL-Ch was elevated significantly in APD group compared to normal group. Plasma MDA and TNF- α showed significant increase in APD group compared to normal group. Compared to normal group compared to normal group compared to normal group compared to normal group. Plasma total antioxidant capacity and creatinine clearance were significantly reduced in APD group compared to control normal group.

International Journal of Biological, Life and Agricultural Sciences ISSN: 2415-6612 Vol:7, No:7, 2013

TABLE I Composition of Different Diets (g/100 g)									
a)Ingredients	Balanced diet	High phosphate-high adenine diet	High adenine diet	Balanced diet containing barley bread.	Balanced diet containing functional food I	Balanced diet containing rice bran bread	Balanced diet containing functional food II		
Casein	12	12	12	1.8	1.8	1.44	1.44		
Corn oil	10	10	10	8.2	8.2	7.88	7.5		
^a Flat barley bread	-	-	-	70	-	-	-		
^b Functional food I	-	-	-	-	70	-	-		
^c Rice bran bread	-	-	-	-	-	70	-		
^d Functional food II	-	-	-	-	-	-	70		
Sucrose	23.5	22.8	22.8	13.8	13.8	16.18	16.56		
Starch	47	45.56	46.95	-	-	-	-		
Salt mix.	3.5	-	3.5	3.5	3.5	3.5	3.5		
High phosphate salt mix.	-	4.89	-	-	-	-	-		
Vitamin mixture	1	1	1	1	1	1	1		
Cellulose	3	3	3	1.7	1.7	-	-		
Adenine	-	0.75	0.75	-	-	-	-		

12 g casein contains 10 g protein.
^a 70 g of flat barley bread contains 8.47 g protein, 1.75 g fat and 1.302 crude fibers.
^b 70 g of functional food I contains 8.47 g protein, 1.75 fat and 1.302 crude fibers
^c 70 g of rice bran bread contains 8.8 g protein, 2.12 g fat and 3.143 crude fibers.
^d 70 g of functional food II contains 8.8 g protein, 2.5 g fat and 3.143 crude fibers

TABLE II Proximate Composition of Different Breads										
Parameter g/100g dry sample 100%WBF Flat bread Functional Food I WWF (70%) + (30%) RB Functional Food I										
Protein	12.1±0.01	12.1±0.02	12.57±0.18	12.57 ±0.19						
Fat	2.5±0.03	2.5±0.03	3.03±0.12	3.57±0.13						
Ash	2.64 ± 0.07	2.7±0.06	2.95±0.16	3.0±0.17						
Crude fibers	1.86±0.05	1.86 ± 0.06	4.49±0.05	4.49±0.05						
Carbohydrate	80.9±0.11	80.84±0.12	76.96±1.12	76.37±1.20						
Moisture (g/100g fresh sample)	14.01±0.12	14.01±0.13	42.0±0.62	42.0±0.50						

TABLE III

ORGANOLEPTIC PROPERTIES OF FLAT BREAD									
Samples	Texture (10)	Color (10)	Odor (10)	Taste (10)	Appearance (10)	Overall acceptability (10)			
100%WBF	7.4±0.84	7.1±1.73	6.9±1.73	7.0±1.56	6.8±1.62	7.04±1.69			
100%WBF+ red grape nutraceutical (Functional food I)	6.9±0.32	6.5±1.01	6.7±0.86	6.6±0.46	6.5±1.13	6.64±1.25			
	NS	NS	NS	NS	NS	NS			

WBF: Whole meal barley flour

NS: Non significant

TABLE IV Organoleptic Properties of Balady Bread									
Samples	General appearance (20)	Separation of layers (20)	Roundness (15)	Distribution of crumb (15)	Crust colour (10)	Taste (10)	Odour (10)		
70%WWF + 30% RB	15.79±1.85	20±0.85	12.85±1.23	11.64±2.02	8.07b±0.47	7.65b±0.46	8.07b±0.79		
70%WWF + 30% RB+ walnut nutraceutical (Functional food II)	15.25±0.72	20±0.46	12.82±1.06	11.42±1.75	7.91b±0.22	7.50b±0.22	8.15b±0.35		
	NS	NS	NS	NS	NS	NS	NS		

WWF: Whole meal wheat flour

RB: rice bran

NS: Non significant

International Journal of Biological, Life and Agricultural Sciences ISSN: 2415-6612 Vol:7, No:7, 2013

BIOCHEMICAL PARAMETERS OF DIFFERENT EXPERIMENTAL GROUPS									
Groups	Creatinine	Urea	Albumin	TNF-α	TAC	MDA	T-Ch	LDL-Ch	Creatinine Clearance
	(mg/dl)	(mg/dl)	(g/dl)	(pg/ml)	(mM/l)	(nmol/l)	(mg/dl)	(mg/dl)	(ml/min)
Normal control	0.738 ^a	25.3ª	4.1 ^a	17.5 ^a	0.734 ^a	7.8 ^a	91.2 ^a	18.5 ^a	2.4 ^a
Normal control	±0.038	±0.997	±0.115	±0.359	±0.026	±0.215	±1.402	±0.344	±0.069
A DD as metrical	3.2 ^b	92.8 ^b	3.3 ^b	29.7 ^b	0.141 ^b	12.2 ^b	94.7 ^a	24.5 ^b	1.45 ^b
APD control	±0.174	±1.872	± 0.086	±0.592	±0.007	±0.269	±1.195	±0.764	±0.118
ADD halamand dist	2.5°	77.6 ^c	3.6 ^c	26.3°	0.155 ^b	11.1°	92.9 ^a	23.5 ^b	1.8 ^c
APD + balanced diet	±0.092	±1.953	±0.094	±0.321	±0.012	±0.347	± 0.803	±0.428	±0.168
A DD De sleve have a	1.8 ^d	68.4 ^d	4.0 ^{ac}	24.7 ^d	0.216 ^d	9.1 ^d	91.4 ^a	21.7 ^d	2.24 ^{ac}
APD+ Barley bread	±0.173	±2.192	±0.290	±0.336	±0.016	±0.149	± 0.852	±0.422	±0.106
APD+ Barley bread	1 5 ^d	67.2 ^d	4 1 ^{ad}	24 2 ^d	0 227 ^d	8 9 ^d	93.2ª	20 3 ^d	2 4 ^{ac}
with grape extract	+0.237	+3 022	+0.143	+0.331	+0.0227	+0.351	+1 215	+0.422	+0.135
(Functional food I)	-0.257	±3.022	40.145	20.551	+0.022	±0.551	=1.210	-0.122	20.155
ADD D broad	2.1 ^e	69.5 ^d	3.7 ^{ac}	25.2 ^d	0.182 ^d	9.1 ^d	92.5ª	22 ^d	2.2^{ac}
APD+KB bread	±0.173	±1.599	±0.189	±0.312	±0.016	±0.235	± 1.231	±0.365	±0.065
APD+ RB bread with	2 1°	67.8 ^d	3 Qac	24 Qd	0 199 ^d	9 0 ^d	01 5 ^a	21 ^d	2 26 ^{ac}
walnut extract	+0.227	+2 780	10 192	2 4 .9	+0.002	+0.244	± 2.171	± 0.265	10.060
(Functional food II)	±0.237	±2.789	±0.182	±0.310	±0.003	±0.344	-2.1/1	±0.303	±0.009

TABLE V

In column different letters means significant difference at 0.05 probabilities

TABLE VI NUTRITIONAL PARAMETERS OF DIFFERENT EXPERIMENTAL GROUPS

	First stage			Second stage				
Parameters	Normal APD control control (n=6) (n=36)		APD fed balanced diet	APD fed Barley bread	APD fed Barley + grape bread (Functional food I)	APD fed RB bread	APD fed RB + Walnut bread (Functional food II)	
Initial BW(g)	215.7	215.9	141.7 ^a	141.2^{a}	141.3 ^a	141.2 ^a	141.2ª	
Final BW (g)	± 2.990 272.5 ± 7.176	± 3.402 156.3* ± 3.322	± 2.882 206.5 ° ± 5.777	± 4.501 191.8 ° ± 5.042	± 4.909 196.2 ° ± 6.568	± 5.973 192.8 ^a ± 7.267	± 5.323 192.3 ^a ± 5.529	
Body weight gain (g)	59.8 ± 3.496	-59.7* ± 3.301	64.8^{a} ± 5.635	50.7^{a} ± 3.738	54.8^{a} ± 3.806	51.6^{a} ± 3.062	55.5^{a} ± 2.029	
Total food intake (g)	520.5 ± 6.232	209.9 * ± 2.329	517.5 ° ± 5.438	514.8 ^a ±10.446	503.3 ° ± 7.030	511 ^a ± 8.739	518.3 ª ± 6.665	
Food efficiency ratio	0.115 ± 0.006	-0.284* ± 0.015	0.126 ^a ± 0.012	$0.098^{a} \pm 0.007$	0.109 ^a ± 0.006	$0.101^{a} \pm 0.005$	0.107 * ± 0.004	

First stage: Values significantly differ from normal control: *: p< 0.001

Second stage: In each row same letters means nonsignificant difference; different letter means the significance among the tested groups at 0.05 probabilities

Feeding balanced diet after induction of kidney dysfunction produced significant reduction of plasma creatinine, urea, TNF- α and MDA with significant increase in plasma albumin and creatinine clearance which were still not matching the control normal. However no significant change was noticed in plasma TAC, T.Ch and LDL-Ch. Feeding balanced diet mixed with any type of bread after induction of kidney dysfunction produced significant improvement of the different biochemical parameters reflecting improvement in kidney dysfunction compared to APD group. Also, feeding APD rats balanced diet mixed with different bread produced significant improvement in all biochemical parameters compared to those fed the balanced diet except for creatinine clearance that showed an improvement that was nonsignificant. Also, albumin demonstrated only significant improvement in the group fed functional food I compared to those fed on balanced diet. There was only significant reduction of plasma creatinine in APD rats fed on either barley bread or functional food I compared to those fed on rice bran bread or functional food II, but there was no significant change in all other biochemical parameters among the groups fed on the different bread.

However, functional food I seem to be the most efficient in improving kidney dysfunction.

Table VI showed the nutritional parameters of different experimental groups. Nutritional parameters (body weight gain, total food intake, final body weight and food efficiency ratio) were significantly reduced in APD group compared to normal control. Rats regained weight after stopping APD and feeding either balanced diet or balanced diet containing any of the studied bread. All nutritional parameters among APD fed rats after feeding on balanced diet or balanced diets containing different breads showed non-significant changes when compared with each other.

IV. DISCUSSION

Kidney is a vital organ for the body, therefore renal impairment results in high mortality and morbidity among population. Different animal models of renal dysfunction have been developed for evaluation of new therapy; high adenine or high adenine high phosphate diet were fed to rats for different periods of time for induction of such models [21], [22]. On feeding adenine it is metabolized to 2, 8-dihydroxyadenine that induced severe chronic renal failure due to degeneration of the proximal and distal tubules and interstitial fibrosis [23]. Hyperphosphatemia, is a nearly universal complication of kidney failure [24], therefore high phosphate diet contributed in the induction of renal dysfunction in rats. So in the present research, rats were fed high adenine-high phosphate diet for 10 days followed by feeding high adenine diet for 11 days for induction of chronic renal failure in rats. Results emphasized the induction of chronic renal failure through the changes noticed in the different biochemical parameters.

In the present study, APD produced significant reduction in body weight and food efficiency which might be due to reduction in food intake as shown from the present results. Reduction in body weight may also be ascribed to loss of muscle and visceral protein stores as reported previously [25]. Loss of muscle protein may be related to stimulation of protein degradation and/or reduction of protein synthesis reported in kidney disease [26]. Loss of renal function induced a condition of protein energy wasting as demonstrated previously [25]. Gastrointestinal function and protein metabolism could also be affected by uremia which may share in the resulted poor nutritional status [26].

APD rats showed significant reduction in plasma level of albumin. This reduction in albumin level may be due to loss of albumin in urine due to kidney dysfunction and/ or loss of muscle protein. Hypoalbuminemia is a common feature in protein energy wasting and has a strong association with increased mortality [27] and morbidity [28]. Albumin level is a negative acute phase reactant and its serum level is profoundly affected by the presence of an inflammatory response [29] that could be emphasized by the increased TNF- α in APD group in the present study. Significant increase in the inflammatory cytokines (IL-1, IL-6, TNF- α) have been reported in dialyzed patients suggesting that a systemic inflammatory response is common in dialysis-treated patients [30], [31]. Inflammation has been accused for serologic and anthropometric evidence of malnutrition [30].

Malondialdehyde as indicator of lipid peroxidation increased while total antioxidant capacity reduced, in APD rats which accounted for increased oxidative stress. Increased levels of malondialdehyde and F₂-isoprostanes, two products of lipid peroxidation, were reported to be associated with renal damage [3]. In kidney diseases, oxidative stress results from an imbalance between oxidant generation and anti-oxidant defense mechanisms, leads to cell and tissue injury. Nguyen-Khoa et al. [32] demonstrated a direct correlation between lipid and protein oxidation markers in patient with renal dysfunction.

In the present study, LDL-Ch increased significantly in APD group which together with its oxidized form that may result from elevated oxidative stress might be involved in the glomerular damage of endothelial and mesangial cells. Oxidation of LDL is induced by infiltrating leukocytes. Native LDL can stimulate fibronectin secretion by mesangial cells while oxidized LDL may stimulate the genetic expression of fibronectin [33].

In a previous study, nutraceuticals prepared from red grape and walnut showed to be efficient in prevention of cisplatin induced kidney dysfunction [8]. These nutraceuticals were incorporated in the present study into functional foods (made from barley or rich in rice bran) to test their efficiency in improving chronic renal failure when acting synergistically with the active constituents of barley and rice bran.

Rats fed on balanced diet containing rice bran bread, barley bread, functional food I and functional food II showed significant improvement in the majority of biochemical parameters compared to APD and APD fed balance diet. Also these treatments showed improvement in all the nutritional parameters compared to APD group.

The activity of the prepared functional foods in the reduction of kidney failure may be attributed to the presence of biologically active constituents. In a previous work, walnut and grape nutraceuticals were shown to contain phenolic compounds and unsaturated fatty acids which were of extremely higher level in walnut than in grape while phytosterol content in grape nutraceutical was higher [8].

Red grape was reported to have antioxidant and antiinflammatory effect [34] due to presence of resveratrol (polyphenolic compound), beta-carotene, tocopherols and dietary fiber [35], [36]. Walnut is a good source of essential fatty acids (linoleic acid), tocopherols, tocotrienols, phytosterols, tannins and other polyphenols [37]. Most phenolic compounds commonly identified in walnut seeds are phenolic acids, namely gallic, ellagic, syringic, 5-Ocaffeoylquinic, caffeic, p-coumaric, ferulic and sinapic acids, and tannins, such as glansrins A, B and C, casuarinin and stenophyllarin [38], [39], therefore it is expected to possess antioxidant and anti-inflammatory effect [40]. So, it was proposed that both nutraceuticals could improve kidney dysfunction through alleviating oxidative stress and inflammation in addition of reducing LDL-Ch.

Rice bran contains an array of bio-active phytochemicals such as γ -oryzanols, phytosterols, tocopherols, tocotrienols, squalene, policosanols, phytic acid and ferulic acid [41], [42]. Also, rice bran is a good source of insoluble dietary fiber, protein and phytic acid [43] that possesses antioxidant, antiinflammatory and LDL-Ch lowering effect [43]. On the other hand barley contains phytate, beta-glucan, tocopherols and tocotrienols [5] that may reflect its antioxidant and antiinflammatory activity. This may explain the improvement in kidney dysfunction on feeding the bread made from barley and that rich in rice bran in the present study.

Reported diuretic activity of bioactive constituents that present in functional foods in the current study such as phytosterols and flavonoids [44], [45] may participate in the renal protective effect through increasing creatinine clearance. Reduction of plasma creatinine and increased creatinine clearance on feeding the different bread reflected the improvement of glomerular function and proximal tubules [46].

Acceptance of different bread from sensorial point of view may encourage the possible marketing of such bread for preventing renal dysfunction or as dietary intervention during chronic kidney diseases.

V. CONCLUSION

Both barley and rice bran bread with or without the studied nutraceuticals produced improvement in kidney function that might be ascribed to the presence of antioxidant and antiinflammatory active constituents. It could be noticed that functional food made from barley and red grape nutraceutical was the most promising in improving kidney dysfunction.

ACKNOWLEDGMENT

This research is a part of a project (No: 9130103) financed by National Research Centre, Egypt.

REFERENCES

- N. A. G. Santos, C. S. Catão, N. M. Martins, C. Curti, M. L. P. Bianchi, and A. C. Santos, "Cisplatin-induced nephrotoxicity is associated with oxidative stress, redox state unbalance, impairment of energetic metabolism and apoptosis in rat kidney mitochondria," *Archives of Toxicology*, 81 (7), 495-504, 2007.
- [2] N. D. Vaziri, and K. C. Norris, "Reasons for the lack of salutary effects of cholesterol-lowering interventions in end-stage renal disease populations" *Blood Purif.*, 35(1-3), 31-6, 2013.
- [3] M.C. Martin-Mateo, M. Sanchez-Portugal, S. Iglesias, A. de Paula, and J. Bustamante, "Oxidative stress in chronic renal failure," *Ren. Fail.*, 21, 155–167, 1999.
- [4] J. Galle, K. Heermeier, and C. Wanner, "Atherogenic lipoproteins, oxidative stress and cell death," *Kidney Int.*, 56 (Suppl. 71), S62–S65, 1999.
- [5] R. A. Moreau, P. Bregitzer, K. Liu, and K. B. Hicks, "Compositional equivalence of barleys differing only in low- and normal-phytate levels," *J Agric Food Chem.*, 60(26), 6493-8, 2012.
- [6] M. S.Islam, R. Nagasaka, K. Ohara, T. Hosoya, H. Ozaki, H. Ushio, and M. Hori, "Biological abilities of rice bran-derived antioxidant phytochemicals for medical therapy," *Curr Top Med Chem.*, 11(14), 1847-53, 2011.
- [7] S. Y. Al-Okbi, N. M. Ammar, D. A. Mohamed, I. M. Hamed, A. H. Desoky, H. F. El-Bakry, and A. M. Helal, "Egyptian rice bran oil: chemical analysis of the main phytochemicals (Periodical style— Accepted for publication)," *La Rivista Italiana delle Sostanze Grasse*, to be published.
- [8] S. Y. Al-Okbi, Prevention and Management of Some Chronic Diseases (Renal and hepatic) by Nutraceuticals and Functional Foods (2010-2013), a project in National Research Centre, Egypt.
- [9] N. Erdinest, O.Shmueli, Y. Grossman, H. Ovadia, and A. Solomon, "Anti-inflammatory effects of alpha linolenic acid on human corneal epithelial cells,". *Invest Ophthalmol Vis Sci.*, 53(8), 4396-406. 2012.
- [10] T. E. Sialvera, G. D. Pounis, A. E. Koutelidakis, D. J. Richter, G. Yfanti, M. Kapsokefalou, G. Goumas, N. Chiotinis, E. Diamantopoulos, and A. Zampelas, "Phytosterols supplementation decreases plasma small and dense LDL levels in metabolic syndrome patients on a westernized type diet." *Nutr Metab Cardiovasc Dis.*, 22(10), 843-8, 2012.
- [11] M. Lutz, K. Jorquera , B. Cancino B, R. Ruby, and C. Henriquez, "Phenolics and antioxidant capacity of table grape (*Vitis vinifera* L.) Cultivars grown in Chile." *J Food Sci.*, 76(7), C1088-93, 2011.
- [12] A.O.A.C., "Official Methods of Analysis of the Association of Official Analytical Chemists," 17th ed., Association of Official Analytical Chemists, Arlington, Virginia, USA, 2000.
- [13] K. Satoh, "Serum lipid peroxide in cerebrovascular disorders determined by a new colorimetric method," *Clinica Chimica Acta*, 20, 37-43, 1978.
- [14] T. Koeda, K. Wakaki, F. Koizumi, T. Yokozawa, and H. Oura, "Early changes of proximal tubules in the kidney of adenine-ingesting rats, with special reference to biochemical and electron microscopic studies," *Nippon Jinzo Gakkai Shi*, 30: 239–46, 1998.

- [15] J. A. Stepaniak, K. E. Gould, D. Sun, and R. H. Swanborg, "A comparative study of experimental autoimmune encephalomyelitis in Lewis and DA rats, "*J Immunol.*, 155, 2762-2769, 1995.
- [16] Houot O. 1985. "Interpretation of clinical laboratory tests," G. Siest, J. Henny, F. Schiele, D. S. Young, edits., Biomedical publications, 1985.
 [17] J. K. Fawcett, J. E. Scott, "A rapid and precise method for the
- [17] J. K. Fawcett, J. E. Scott, "A rapid and precise method for the determination of urea," *J. Clin. Pathol.*, 13, 156-159, 1960.
- [18] B. T. Doumas, W. A.Watson, and H. G. Biggs, "Albumin standards and the measurement of serum albumin with bromocresol green," Clin. *Chem. Acta.*, 1972, 31, 87, 1972.
 [19] D. A. Watson, "Simple method for the determination of serum
- [19] D. A. Watson, "Simple method for the determination of serum cholesterol," *Clin. Chem. Acta.*, 5, 637-642, 1960.
- [20] H. Schriewer, U. Kohnert, G. Assmann, "Determination of LDL cholesterol and LDL apolipoprotein B following precipitation of VLDL in blood serum with phosphotungstic acid/MgCl₂," J. Clin. Chem. Clin. Biochem., 22, 35-40, 1984.
- [21] K. Katsumata, K. Kenichiro, H. Michinori, T. Kunihiko, N. Nobuo, K. B. Steven, "Sevelamer hydrochloride prevents ectopic calcification and renal osteodystro- phy in chronic renal failure rats," *Kidney Int.*, 64, 441–50, 2003.
- [22] M. Shuvy, S. Abedat, R. Beeri, H. D. Danenberg, D. Planer, I. Z. Ben-Dov, K. Meir, J. Sosna, and C. Lotan, "Uraemic hyperparathyroidism causes a reversible inflammatory process of aortic valve calcification in rats," *Cardiovascular Research*, 79, 492–499, 2008.
- [23] H. Okada, Y. Kaneko, T. Yawata, et al: "Reversibility of adenineinduced renal failure in rats," *Clin Exp Nephrol.*, 3, 82–88, 1999.
- [24] M. Tonelli, N. Pannu and B. Manns, "Oral Phosphate Binders in Patients with Kidney Failure," N Engl J Med., 362, 1312-1324, 2010.
- [25] D. Foque, K. Kalantar-Zadeh, J. D. Kopple, N. Canau, P. Chauveau, L. Cuppari, H. Franch, G. Guarnieri, T. A. Ikizler, and G. A. Kaysen, "Proposed nomenclature and diagnostic criteria for protein-energy wasting in acute and chronic kidney disease," *Kidney Int.*, 73, 391-398, 2008.
- [26] A. Bonanni, I. Mannucci, D. Verzola, A. Sofia, S. Saffioti, E. Gianetta, and G. Garibotto, "Protein-Energy Wasting and Mortality in Chronic Kidney Disease," *Int. J. Environ. Res. Public Health*, 8, 1631-1654, 2011.
- [27] K. Kalantar-Zadeh, R. D. Kilpatrick, and N. Kuwae, "Revisiting mortality predictability of serum albumin in the dialysis population: time dependency, longitudinal changes and population-attributable fraction," *Nephrol. Dial. Transpl.*, 20, 1880-1888, 2005.
- [28] T. A. Ikizler, R. I. Wingard, J. Harvell, Y. Shyr, and R. Hakim, "Association of morbidity with markers of nutrition and inflammation in chronic hemodialysis patients: A prospective study," *Kidney Int.*, 55, 1945-1951, 1999.
- [29] G. M. Chertow, D. J. Goldstein-Fuchs, J. M. Lazarus, G. A. Kaysen, "Prealbumin, mortality, and cause-specific hospitalization in hemodialysis patients. Kidney Int., 68, 2794-2800, 2005.
- [30] P. Stenvinkel, O. Haimburger, F. Paultre, U. Diczfalusy, T. Wang, L. Berglund, and T. Jogestrand, "Strong association between malnutrition, inflammation, and atherosclerosis in chronic renal failure," *Kidney Int.*, 55, 1899-1911, 1999.
- [31] V. Panichi, M. Migliori, S. De Pietro, D. Taccola, A. M. Bianchi, L. Giovannini, M. Norpoth, M. R. Metelli, R. Cristofani, A. A. Bertelli, "C-reactive protein and interleukin-6 levels are related to renal function in predialytic chronic renal failure," *Nephron*, 91, 594-600, 2002.
- [32] T. Nguyen-Khoa, Z. A. Massay, J. P. De Bandt, M. Kebede, L. Salama, G. Lambrey, V. Witko-Sarsat, T. B. Drüeke, B. Lacour, and M. Thévenin, "Oxidative stress and haemodialysis: Role of inflammation and duration of dialysis treatment," *Nephrol. Dial. Transpl.*, 16, 335-340, 2001.
- [33] G. Ding, H. Van Goor, S. D. Ricardo, J. M. Orlowski, and J. R. Diamond, "Oxidized LDL stimulates the expression of TGF and fibronectin in human glomerular epithelial cells," *Kidney Int.*, 51, 147–154, 1997.
- [34] E. F. Gris, F. Mattivi, E. A. Ferreira, U. Vrhovsek, D. W. Filho, R. C. Pedrosa, M. T. Bordignon-Luiz, "Stilbenes and tyrosol as target compounds in the assessment of antioxidant and hypolipidemic activity of Vitis vinifera red wines from southern Brazil." J Agric Food Chem., 59(14), 7954-61, 2011.
- [35] G. M. Cole, G. P. Lim, and F. Yang, "Prevention of Alzheimer's disease: Omega-3 fatty acid and phenolic antioxidant interventions," *Neurobiol. Aging*, 26S, S133 - S136, 2005.

International Journal of Biological, Life and Agricultural Sciences ISSN: 2415-6612 Vol:7, No:7, 2013

- [36] Rho K.A., Kim M.K. Effects of different grape formulations on antioxidative capacity, lipid peroxidation and oxidative DNA damage in aged rats. J. Nutr. Sci. Vitaminol., Tokyo 2006; 52 (1): 33 - 46.
- [37] J. A. Pereira, I. Oliveira, A. Sousa, I. Ferreira, A. Bento, and L. Estevinho, "Bioactive properties and chemical composition of six walnut (*Juglans regia* L.) cultivars," *Food Chem. Toxicol.*, 46, 2103–2111, 2008.
- [38] T. Fukuda, H. Ito, and T. Yoshida, "Antioxidative polyphenols from walnuts (Juglans regia L.)," *Phytochemistry*, 63, 795–801, 2003.
- [39] M. Colaric, R. Veberic, A. Solar, M. Hudina, and F. Stampar, "Phenolic acids, syringaldehyde, and juglone in fruits of different cultivars of Juglans regia L," J. Agr. Food Chem., 53, 6390–6396, 2005.
- [40] B. Muthaiyah, M. M. Essa, V. Chauhan, A. Chauhan, "Protective effects of walnut extract against amyloid beta peptide-induced cell death and oxidative stress in PC12 cells," *Neurochem Res.*, 36(11), 2096-103, 2011.
- [41] S. Khatoon, and A. G. Gopalakrishna, "Fat-soluble nutraceuticals and fatty acid composition of selected Indian rice varieties," *Journal of the American Oil Chemists' Society*, 81, 939–943, 2004.
- [42] H. Ardiansyah Shirakawa, T. Koseki, K. Ohinata, K. Hazhizume, and M. Komai, "Rice bran fractions improve blood pressure, lipid profile, and glucose metabolism in stroke-prone spontaneously hypertensive rats," *Journal of Agricultural and Food Chemistry*, 54, 1914–1920, 2006.
- [43] Y. Saito, K. Ikushima, S. Hatakeda, T. Ito, and T. Goto, "Fractional extraction of rice bran oil and its esters with supercritical carbon dioxide," *Int. Chem.Eng.*, 33, 307 – 314, 1993.
- [44] S. Christie, A. F. Walker, and G. T. Lewith, "Flavonoids a new direction for the treatment of fluid retention," *Phytother Res*, 15:, 467– 475, 2001.
- [45] K. K. Hullatti, U. V. Gopikrishna, I.J.J Kuppast, "Phytochemical investigation and diuretic activity of Cyclea peltata leaf extracts", . Adv Pharm Technol Res., 2(4), 241-4, 2011.
- [46] F. Van Lente, and P. Sult, "Assessment of renal function by serum creatinine and creatinine clearance: Glomerular filteration rate estimated by four procedures," *Clin.Chem.*, 35 (12): 2326-2330, 1989.