

Traveling wave solutions for the (3+1)-dimensional breaking soliton equation by $(\frac{G'}{G})$ -expansion method and modified F-expansion method

Mohammad Taghi Darvishi, Maliheh Najafi and Mohammad Najafi

Abstract—In this paper, using $(\frac{G'}{G})$ -expansion method and modified F-expansion method, we give some explicit formulas of exact traveling wave solutions for the (3+1)-dimensional breaking soliton equation. A modified F-expansion method is proposed by taking full advantages of F-expansion method and Riccati equation in seeking exact solutions of the equation.

Keywords—Exact solution, The (3+1)-dimensional breaking soliton equation, $(\frac{G'}{G})$ -expansion method, Riccati equation, Modified F-expansion method.

I. INTRODUCTION

IN many different fields of science and engineering, it is very important to obtain exact or numerical solutions of nonlinear partial differential equations. It is well known that nonlinear phenomena are very important in a variety of scientific fields, especially in fluid mechanics, solid state physics, plasma physics, plasma waves and chemical physics. Searching for exact and numerical solutions, especially for traveling wave solutions, of nonlinear equations in mathematical physics plays an important role in soliton theory [1], [2]. Recently many new approaches to nonlinear equations were proposed, such as the homotopy perturbation method [3], [4], [5], [6], [7], the variational iteration method [8], [9], [10], parameter expansion method [11], [12], [13], [14], spectral collocation method [15], [16], [17], [18], [19], homotopy analysis method [20], [21], [22], [23], [24], [25], and the Exp-function method [26], [27], [28], [29], [30], [31]. In this paper, we solve a (3+1)-dimensional breaking soliton equation by the $(\frac{G'}{G})$ -expansion method and modified F-expansion method, and obtain some exact and new solutions for it.

The(2+1)-dimensional breaking soliton equation has the following form

$$u_{xt} - 4u_{xy}u_x - 2u_{xx}u_y - u_{xxx} = 0, \quad (1)$$

this equation describes the (2+1)-dimensional interaction of the Riemann wave propagated along the y -axis with a long wave propagated along the x -axis [32]. Wazwaz [33] presented an extension to equation (1) by adding the last three terms with y replaced by z . By his work, one enables to establish

M.T. Darvishi, Maliheh Najafi and Mohammad Najafi: Department of Mathematics, Faculty of Science, Razi University, Kermanshah 67149, Iran. **e-mail:** darvishimt@yahoo.com, malihe_math87@yahoo.com and m_najafi82@yahoo.com.

the following (3+1)-dimensional breaking soliton equation

$$u_{xt} - 4u_x(u_{xy} + u_{xz}) - 2u_{xx}(u_y + u_z) - \quad (2)$$

$$(u_{xxy} + u_{xxz}) = 0,$$

where $u = u(x, y, z, t) : \mathbb{R}_x \times \mathbb{R}_y \times \mathbb{R}_z \times \mathbb{R}_t \rightarrow \mathbb{R}$.

In this paper, by means of the $(\frac{G'}{G})$ -expansion method and modified F-expansion method, we obtain some exact traveling wave solutions for equation (2).

The outline of this paper is as follows. In the following section we have a brief review on the $(\frac{G'}{G})$ -expansion. In Section III we apply the $(\frac{G'}{G})$ -expansion method on equation (2) to obtain some traveling wave solution for the equation. In Section IV a review on the modified F-expansion method is presented. We obtain some traveling wave solutions for equation (2) by the modified F-expansion method in Section V. The paper is concluded in Section VI.

II. THE $(\frac{G'}{G})$ -EXPANSION METHOD

Wang et al. [34] proposed the $(\frac{G'}{G})$ -expansion method to solve nonlinear partial differential equations, where $G = G(\xi)$ satisfies a second order linear ordinary differential equation. In this section we describe the $(\frac{G'}{G})$ -expansion method to find traveling wave solutions of nonlinear evolution equations. Suppose that a nonlinear equation, say in two independent variables x, t , is given by

$$P(u, u_t, u_x, u_{tx}, u_{xx}, \dots) = 0, \quad (3)$$

where $u = u(x, t)$ and P is a polynomial of u and its derivatives in which the highest order derivatives and nonlinear terms are involved. The main steps of the $(\frac{G'}{G})$ -expansion method are as follows:

- **First.** Suppose that

$$u(x, t) = u(\xi), \quad \xi = x + wt \quad (4)$$

the traveling wave variable (4) permits us reducing (3) to an ordinary differential equation (shortly ODE) for $u = u(\xi)$ such as

$$P(u, u', u'', u''', \dots) = 0, \quad (5)$$

- **Second.** Now, we suppose that the solution of (5) can be expressed by a polynomial in $(\frac{G'}{G})$ as

$$u(\xi) = \alpha_m \left(\frac{G'}{G}\right)^m + \dots \quad (6)$$

where $G = G(\xi)$ satisfies the second order linear ODE in the following form

$$G'' + \lambda G' + \mu G = 0 \tag{7}$$

α_m, \dots, λ and μ are constants to be determined later, $\alpha_m \neq 0$. The unwritten part in (6) is also a polynomial in $(\frac{G'}{G})$, the degree of which is generally equal to or less than $m - 1$. The positive integer m can be determined by considering the homogeneous balance between the highest order derivatives and nonlinear terms appearing in (5).

- **Third.** Substituting (6) into (5) and using the second order linear ODE (7), collecting all terms with the same order of $(\frac{G'}{G})$ together, the left-hand side of (5) is converted into another polynomial in $(\frac{G'}{G})$. Equating each coefficients of this polynomial to zero, yields a set of algebraic equations for α_m, \dots, λ and μ .
- **Fourth.** Assuming that the constants α_m, \dots, λ and μ can be obtained by solving the algebraic equations in item **Third**. Since we know the general solutions of the second order linear ODE (7), then substituting α_m, \dots, w and the general solutions of (7) into (6) yields the traveling wave solutions of the nonlinear equation (3).

III. APPLICATION OF THE $(\frac{G'}{G})$ -EXPANSION METHOD FOR THE (3+1)-DIMENSIONAL BREAKING SOLITON EQUATION

In this section, we apply the $(\frac{G'}{G})$ -expansion method to construct the traveling wave solutions for (3+1)-dimensional breaking soliton equation

$$u_{xt} - 4u_x(u_{xy} + u_{xz}) - 2u_{xx}(u_y + u_z) - (u_{xxxy} + u_{xxxz}) = 0. \tag{8}$$

Some exact solutions for (8) were presented in [35] by the three-wave method. To apply the $(\frac{G'}{G})$ -expansion method on this equation, we suppose that

$$u(x, y, z, t) = u(\xi), \quad \xi = kx + my + nz + wt \tag{9}$$

k, m, n, w are constants that to be determined later.

By substitute eq. (9) into eq. (8), we obtain

$$w u'' - 6k(m+n)u' u'' - k^2(m+n)u^{(4)} = 0. \tag{10}$$

Integrating (10) once, we have

$$w u' - 3k(m+n)(u')^2 - k^2(m+n)u^{(3)} = c \tag{11}$$

where c is the integration constant that can be determined later.

Suppose that the solutions of the ODE (11) can be expressed by a polynomial in $(\frac{G'}{G})$ as follows:

$$u(\xi) = \sum_{i=0}^m a_i \left(\frac{G'}{G}\right)^i \tag{12}$$

where a_i are constants, $G = G(\xi)$ satisfies the following second order linear ODE

$$G'' + \lambda G' + \mu G = 0, \tag{13}$$

where λ and μ are constants.

By balancing the order of $(u')^2$ and $u^{(3)}$ in eq. (11), we have $2m + 2 = m + 3$ then $m = 1$. So we can write

$$u(\xi) = a_1 \left(\frac{G'}{G}\right) + a_0, \quad a_1 \neq 0 \tag{14}$$

a_1, a_0 are constants to be determined later.

Then it follows:

$$u'(\xi) = -a_1 \left(\frac{G'}{G}\right)^2 - a_1 \lambda \left(\frac{G'}{G}\right) - a_1 \mu \tag{15}$$

$$u'' = 2a_1 \left(\frac{G'}{G}\right)^3 + 3a_1 \lambda \left(\frac{G'}{G}\right)^2 + a_1 (\lambda^2 - 2\mu) \left(\frac{G'}{G}\right) + \lambda \mu a_1 \tag{16}$$

and

$$u''' = -6a_1 \left(\frac{G'}{G}\right)^4 - 12a_1 \lambda \left(\frac{G'}{G}\right)^3 - a_1 (8\mu + 7\lambda^2) \left(\frac{G'}{G}\right)^2 - a_1 (\lambda^3 + 8\lambda\mu) \left(\frac{G'}{G}\right) - a_1 (\lambda^2 \mu + 2\mu^2). \tag{17}$$

Substituting eq. (14) into eq. (11) and collecting all terms with the same power of $(\frac{G'}{G})$ together, equating each coefficient to zero, yields a set of simultaneous algebraic equations as follows:

$$\left(\frac{G'}{G}\right)^4 : 6k^2m - 3ka_1m - 3ka_1n + 6k^2n = 0$$

$$\left(\frac{G'}{G}\right)^3 : 12k^2n\lambda - 6ka_1n\lambda - 6ka_1m\lambda + 12k^2m\lambda = 0$$

$$\left(\frac{G'}{G}\right)^2 : -6ka_1m\mu - 3ka_1n\lambda^2 - 3ka_1m\lambda^2 - w + 7k^2m\lambda^2 + 8k^2m\mu - 6ka_1n\mu + 8k^2n\mu + 7k^2n\lambda^2 = 0$$

$$\left(\frac{G'}{G}\right)^1 : 8k^2m\lambda\mu - 6ka_1m\lambda\mu - 6ka_1n\lambda\mu + k^2m\lambda^3 + k^2n\lambda^3 - w\lambda + 8k^2n\lambda\mu = 0$$

$$\left(\frac{G'}{G}\right)^0 : a_1(-3ka_1m\mu^2 + k^2m\lambda^2\mu + k^2n\lambda^2\mu) - a_1(w\mu - 3ka_1n\mu^2 + 2k^2m\mu^2 + 2k^2n\mu^2) - c = 0.$$

Solving the above algebraic equations by using Maple, we get:

$$a_0 = a_0, a_1 = 2k, c = 0, w = -(4k^2\mu - k^2\lambda^2)(m+n) \tag{18}$$

where k, n, m, a_0 are arbitrary constants.

Substituting(18) into eq. (14), we have

$$u(\xi) = 2k \left(\frac{G'}{G}\right) + a_0 \tag{19}$$

where $\xi = kx + my + nz - (4k^2\mu - k^2\lambda^2)(m + n)t$. Substituting the general solutions of eq. (13) into eq. (19), we obtain:

Case i: When $\lambda^2 - 4\mu > 0$

$$u_1(\xi) = -k\lambda + k\sqrt{\lambda^2 - 4\mu} \times \left(\frac{C_1 \sinh \frac{1}{2}\sqrt{\lambda^2 - 4\mu}\xi + C_2 \cosh \frac{1}{2}\sqrt{\lambda^2 - 4\mu}\xi}{C_1 \cosh \frac{1}{2}\sqrt{\lambda^2 - 4\mu}\xi + C_2 \sinh \frac{1}{2}\sqrt{\lambda^2 - 4\mu}\xi} \right) + a_0$$

where

$$\xi = kx + my + nz - (4k^2\mu - k^2\lambda^2)(m + n)t,$$

and C_1, C_2, k, m, n and a_0 are arbitrary constants. In particular, if $C_1 = 1, C_2 = 0, k = m = n = 1, \lambda = 2, \mu = 0$, then we have

$$u(x, y, z, t) = -2 + 2 \tanh(x + y + z + 8t) + a_0.$$

Case ii: When $\lambda^2 - 4\mu < 0$

$$u_2(\xi) = -k\lambda + k\sqrt{4\mu - \lambda^2} \times \left(\frac{-C_1 \sin \frac{1}{2}\sqrt{4\mu - \lambda^2}\xi + C_2 \cos \frac{1}{2}\sqrt{4\mu - \lambda^2}\xi}{C_1 \cos \frac{1}{2}\sqrt{4\mu - \lambda^2}\xi + C_2 \sin \frac{1}{2}\sqrt{4\mu - \lambda^2}\xi} \right) + a_0$$

where

$$\xi = kx + my + nz - (4k^2\mu - k^2\lambda^2)(m + n)t,$$

and C_1, C_2, k, m, n, a_0 are arbitrary constants. In particular, if

$$C_1 = 1, C_2 = 0, k = m = n = 1, \lambda = 0, \mu = 1,$$

then we have

$$u(x, y, z, t) = -2 \tanh(x + y + z - 8t) + a_0.$$

Case iii: When $\lambda^2 - 4\mu = 0$

$$u_3(\xi) = \frac{k(2C_2 - C_1\lambda - C_2\lambda\xi)}{C_1 + C_2\xi} + a_0$$

where

$$\xi = kx + my + nz - (4k^2\mu - k^2\lambda^2)(m + n)t,$$

and C_1, C_2, k, m, n, a_0 are arbitrary constants. In particular, if

$$C_1 = 1, C_2 = 1, k = m = n = 1, \lambda = 2, \mu = 1,$$

then we have

$$u(x, y, z, t) = \frac{2}{1 + x + y + z} + a_0$$

IV. THE MODIFIED F-EXPANSION METHOD

We simply describe the modified F -expansion method. To do this we follow descriptions which has presented in [36]. Consider a given nonlinear partial differential equation with independent variables x_1, x_2, \dots, x_l, t and dependent variable u as:

$$P(u, u_t, u_{x_1}, u_{x_2}, \dots, u_{tt}, \dots) = 0, \tag{20}$$

in general, the function P is a polynomial in u and its various partial derivatives, which we seek its traveling wave solutions by taking

$$u(x_1, x_2, \dots, x_l, t) = u(\xi), \xi = k_1(x_1 + k_2 x_2 + \dots + k_l x_l + w t) \tag{21}$$

where k_1, k_2, \dots, k_l and w are unknown constants to be determined. Inserting (21) into (20) yields an ODE for $u(\xi)$ as

$$P(u, u', u'', u''', \dots) = 0 \tag{22}$$

conversely, we suppose that $u(\xi)$ can be expressed as

$$u(\xi) = a_0 + \sum_{i=-N}^N a_i F^i(\xi), \quad (a_N \neq 0) \tag{23}$$

where a_0 and a_i s are constants to be determined. $F(\xi)$ satisfies Riccati equation

$$F'(\xi) = A + B F(\xi) + C F^2(\xi) \tag{24}$$

where A, B, C are constants to be determined. Integer N can be determined by considering the homogeneous balance between the governing nonlinear term(s) and highest order derivatives of $u(\xi)$ in (22).

Substituting (23) into (22), and using (24), the left-hand side of (22) can be converted into a finite series in $F^p(\xi)$ for $p = -N, \dots, -1, 0, 1, \dots, N$. Equating each coefficient of $F^p(\xi)$ to zero yields a system of algebraic equations for $a_i, i = -N, \dots, -1, 0, 1, \dots, N, k_j, j = 1, \dots, l$ and w .

Then solving the obtained system of algebraic equations, with the aid of a symbolic computations like Mathematica or Maple, a_i, k_j, w can be expressed by A, B, C or the coefficients of (22). Finally, by substituting these results into (23), we can obtain the general form of traveling wave solutions to (22).

From the general form of traveling wave solutions of equation (24) listed in Table I, we can give a series of soliton-like solutions, trigonometric function solutions, and exponential function solutions to (20); cf. [36].

TABLE I
SOLUTIONS OF EQUATION (24) GIVEN IN [36].

A	B	C	$F(\xi)$
0	1	-1	$\frac{1}{2} + \frac{1}{2} \tanh\left(\frac{\xi}{2}\right)$
0	-1	1	$\frac{1}{2} - \frac{1}{2} \coth\left(\frac{\xi}{2}\right)$
$\frac{1}{2}$	0	$-\frac{1}{2}$	$\coth(\xi) \pm \operatorname{csch}(\xi), \tanh(\xi) \pm \operatorname{sech}(\xi)$
1	0	-1	$\tanh(\xi), \coth(\xi)$
$\frac{1}{2}$	0	$\frac{1}{2}$	$\sec(\xi) + \tan(\xi), \csc(\xi) - \cot(\xi)$
$-\frac{1}{2}$	0	$-\frac{1}{2}$	$\sec(\xi) - \tan(\xi), \csc(\xi) + \cot(\xi)$
1(-1)	0	1(-1)	$\tan(\xi), \cot(\xi)$
0	$\neq 0$	$\neq 0$	$-\frac{1}{C} \frac{1}{\xi + m}$ (m is arbitrary constant)
arbitrary constant	0	0	$A \xi$
arbitrary constant	$\neq 0$	0	$\frac{\exp(B\xi) - A}{B}$

V. NEW EXACT SOLUTIONS TO THE (3+1)-DIMENSIONAL
BREAKING SOLITON EQUATION BY THE MODIFIED
F-EXPANSION METHOD

In this section, we apply the F-expansion method to construct the traveling wave solutions for the mentioned (3+1)-dimensional breaking soliton equation, that is,

$$u_{xt} - 4u_x(u_{xy} + u_{xz}) - 2u_{xx}(u_y + u_z) - (u_{xxx} + u_{xxxz}) = 0, \quad (25)$$

we suppose that

$$u(x, y, z, t) = u(\xi), \quad \xi = k(x + my + nz + wt) \quad (26)$$

k, m, n, w are constants that to be determined later. By substituting eq. (26) into eq. (25), we obtain

$$w u'' - 6k(m+n)u' u'' - k^2(m+n)u^{(4)} = 0. \quad (27)$$

By balancing the order of $u' u''$ and $u^{(4)}$ in eq. (27), we have $2N + 3 = N + 4$ then $N = 1$. So we can write

$$u(\xi) = a_0 + a_{-1}F^{-1}(\xi) + a_1F(\xi) \quad (28)$$

a_0, a_1, a_{-1} are constants to be determined later. Substituting (28) into (27), and using (24), the left-hand side of (27) can be converted into a finite series in $F^p(\xi)$, for $p = -5, \dots, -1, 0, 1, \dots, 5$. Equating each coefficient of $F^p(\xi)$ to zero yields a system of algebraic equations for $a_{-1}, a_0, a_1, m, n, w$. In fact, we have

$$F^5 : -12kma_1^2C^3 + 24k^2na_1C^4 - 12kna_1^2C^3 + 24k^2ma_1C^4$$

$$F^4 : 60k^2na_1C^3B - 30kma_1^2BC^2 + 60k^2ma_1C^3B - 30kna_1^2BC^2$$

$$F^3 : -24kma_1^2B^2C - 24kna_1^2B^2C + 50k^2na_1C^2B^2 + 2wa_1C^2 - 24kna_1^2AC^2 + 50k^2ma_1C^2B^2 - 24kma_1^2AC^2 + 40k^2na_1C^3A + 12kna_1a_{-1}C^3 + 40k^2ma_1C^3A + 12kma_1a_{-1}C^3$$

$$F^2 : 60k^2ma_1C^2AB - 36kna_1^2ABC + 3wa_1BC + 24kna_1a_{-1}BC^2 - 36kma_1^2ABC + 24kma_1a_{-1}BC^2 + 60k^2na_1C^2AB + 15k^2ma_1CB^3 + 15k^2na_1CB^3 - 6kma_1^2B^3 - 6kna_1^2B^3$$

$$F^1 : k^2na_1B^4 + 12kma_1a_{-1}B^2C + k^2ma_1B^4 + 22k^2na_1CB^2A + 2wa_1CA + 12kna_1a_{-1}B^2C + 16k^2na_1C^2A^2 - 12kma_1^2AB^2 + 16k^2ma_1C^2A^2 + 12kna_1a_{-1}AC^2 - 12kna_1^2A^2C + wa_1B^2 + 22k^2ma_1CB^2A - 12kma_1^2A^2C + 12kma_1a_{-1}AC^2 - 12kna_1^2AB^2$$

$$F^0 : 6kma_{-1}^2BC^2 + wa_{-1}BC + k^2ma_1B^3A + 8k^2ma_1BCA^2 + k^2ma_{-1}B^3C + 8k^2ma_{-1}ABC^2 + 8k^2na_1BCA^2 + 6kna_{-1}^2BC^2 - 6kma_1^2A^2B + wa_1BA - 6kna_1^2A^2B + 8k^2na_{-1}ABC^2 + k^2na_1B^3A + k^2na_{-1}B^3C$$

$$F^{-1} : -12kna_1a_{-1}A^2C + 12kna_{-1}^2B^2C - 12kma_1a_{-1}AB^2 + k^2ma_{-1}B^4 - 12kma_1a_{-1}A^2C + 12kma_{-1}^2AC^2 - 12kna_1a_{-1}AB^2 + wa_{-1}B^2 + 22k^2ma_{-1}AB^2C + 12kna_{-1}^2AC^2 + 12kma_{-1}^2B^2C + 16k^2ma_{-1}A^2C^2 + k^2na_{-1}B^4 + 22k^2na_{-1}AB^2C + 16k^2na_{-1}A^2C^2 + 2wa_{-1}AC$$

$$F^{-2} : -24kna_1a_{-1}A^2B - 24kma_1a_{-1}A^2B + 15k^2na_{-1}AB^3 + 3wa_{-1}AB + 15k^2ma_{-1}AB^3 + 60k^2na_{-1}A^2BC + 60k^2ma_{-1}A^2BC + 36kma_{-1}^2ABC + 6kma_{-1}^2B^3 + 6kna_{-1}^2B^3 + 36kna_{-1}^2ABC$$

$$F^{-3} : 50k^2ma_{-1}A^2B^2 + 24kma_{-1}^2AB^2 + 24kna_{-1}^2AB^2 + 40k^2ma_{-1}A^3C - 12kna_1a_{-1}A^3 + 24kna_{-1}^2A^2C - 12kma_1a_{-1}A^3 + 2wa_{-1}A^2 + 50k^2na_{-1}A^2B^2 + 24kma_{-1}^2A^2C + 40k^2na_{-1}A^3C$$

$$F^{-4} : 60k^2ma_{-1}A^3B + 60k^2na_{-1}A^3B + 30kma_{-1}^2A^2B + 30kna_{-1}^2A^2B$$

$$F^{-5} : 12kna_{-1}^2A^3 + 24k^2ma_{-1}A^4 + 12kma_{-1}^2A^3 + 24k^2na_{-1}A^4.$$

Solving the obtained algebraic equations by using Maple, we have the following solutions:

Case 1: when $A = 0$, we have

$$a_0 = a_0, a_{-1} = 0, a_1 = 2Ck, m = m, n = n, w = -k^2B^2(n+m) \quad (29)$$

Case 2: when $B = 0$, we have

$$a_0 = a_0, a_{-1} = -2kA, a_1 = 2kC, m = m, n = n, w = 16k^2AC(m+n) \quad (30)$$

$$a_0 = a_0, a_{-1} = -2kA, a_1 = 0, m = m, n = n, w = 4k^2AC(m+n)$$

Case 3: when $A = B = 0$, we have

$$a_0 = a_0, a_{-1} = -\frac{w}{6Ck(m+n)}, a_1 = 2Ck, m = m, n = n, w = w \quad (31)$$

A. The soliton-like solutions:

1) When $A = 0, B = 1, C = -1$, from Table I and Case 1, we have

$$u_1 = a_0 - k - k \tanh\left[\frac{1}{2}k(x + my + nz - k^2(n+m)t)\right]$$

2) When $A = 0, B = -1, C = 1$, from Table I and Case 1, we have

$$u_2 = a_0 + k - k \coth\left[\frac{1}{2}k(x + my + nz - k^2(n + m)t)\right]$$

3) When $A = \frac{1}{2}, B = 0, C = -\frac{1}{2}$, from Table I and Case 2, we have

$$u_3 = a_0 - k \coth[k(x + my + nz - k^2(n + m)t)]$$

$$\mp kcsch[k(x + my + nz - k^2(n + m)t)]$$

$$u_4 = a_0 - k \tanh[k(x + my + nz - k^2(n + m)t)]$$

$$\mp k i \operatorname{sech}[k(x + my + nz - k^2(n + m)t)]$$

4) When $A = 1, B = 0, C = 1$, from Table I and Case 2, we have

$$u_5 = a_0 + 2k \tanh[k(x + my + nz + 16k^2(n + m)t)]$$

$$-2k \coth[k(x + my + nz + 16k^2(n + m)t)]$$

B. The trigonometric function solutions:

1) When $A = C = \frac{1}{2}, B = 0$, from Table I and Case 2, we have

$$u_6 = a_0 + k \sec[k(x + my + nz + k^2(n + m)t)]$$

$$+k \tan[k(x + my + nz + k^2(n + m)t)]$$

$$u_7 = a_0 + k \csc[k(x + my + nz + k^2(n + m)t)]$$

$$-k i \cot[k(x + my + nz + k^2(n + m)t)]$$

2) When $A = C = -\frac{1}{2}, B = 0$, from Table I and Case 2, we have

$$u_8 = a_0 - k \sec[k(x + my + nz + k^2(n + m)t)]$$

$$+k \tan[k(x + my + nz + k^2(n + m)t)]$$

$$u_9 = a_0 - k \csc[k(x + my + nz + k^2(n + m)t)]$$

$$-k i \cot[k(x + my + nz + k^2(n + m)t)]$$

C. The rational solution:

When $A = B = 0, C \neq 0$, from Table I and Case 3, we have

$$u_{10} = a_0 + \frac{w}{6Ck(m+n)}[Ck(x + my + nz + wt) + m] -$$

$$\frac{2Ck}{Ck(x + my + nz + wt) + m}$$

VI. CONCLUSIONS

In this paper, by means of the $\left(\frac{G'}{G}\right)$ -expansion method and modified F-expansion method, some exact traveling wave solutions for the (3+1)-dimensional breaking soliton equation are obtained. These methods are very simple and with the aid of a symbolic computation like Maple or Mathematica are easy and straightforward methods which can be applied to other nonlinear partial differential equations. It must be noted that, all obtained solutions have checked in the (3+1)-dimensional breaking soliton equation. All solutions satisfy in the equation.

REFERENCES

- [1] L. Debnath, *Nonlinear partial differential equations for scientists and engineers*, Birkhauser, Boston, 1997.
- [2] A.M. Wazwaz, *Partial differential equations methods and applications*, Rotterdam, Balkema, 2002.
- [3] J.H. He, *New interpretation of homotopy perturbation method*, Int. J. Mod. Phys. B 20(18) (2006) 2561–2568.
- [4] J.H. He, *Application of homotopy perturbation method to nonlinear wave equations*, Chaos, Solitons and Fractals 26(3) (2005) 695–700.
- [5] J.H. He, *Homotopy perturbation method for bifurcation of nonlinear problems*, Int. J. Nonlinear Sci. Numer. Simul. 6(2) (2005) 207–208.
- [6] M.T. Darvishi, F. Khani, *Application of He's homotopy perturbation method to stiff systems of ordinary differential equations*, Zeitschrift fur Naturforschung A, 63a (1-2) (2008) 19–23.
- [7] M.T. Darvishi, F. Khani, S. Hamed-Nezhad, S.-W. Ryu, *New modification of the HPM for numerical solutions of the sine-Gordon and coupled sine-Gordon equations*, Int. J. Comput. Math. 87(4) (2010) 908–919.
- [8] J.H. He, *Variational iteration method-a kind of non-linear analytical technique: some examples*, Int. J. Non-linear Mech., 34(4) (1999) 699–708.
- [9] M.T. Darvishi, F. Khani, A.A. Soliman, *The numerical simulation for stiff systems of ordinary differential equations*, Comput. Math. Appl., 54(7-8) (2007) 1055–1063.
- [10] M.T. Darvishi, F. Khani, *Numerical and explicit solutions of the fifth-order Korteweg-de Vries equations*, Chaos, Solitons and Fractals, 39 (2009) 2484–2490.
- [11] J.H. He, *Bookkeeping parameter in perturbation methods*, Int. J. Nonlin. Sci. Numer. Simul., 2 (2001) 257–264.
- [12] M.T. Darvishi, A. Karami, B.-C. Shin, *Application of He's parameter-expansion method for oscillators with smooth odd nonlinearities*, Phys. Lett. A, 372(33) (2008) 5381–5384.
- [13] B.-C. Shin, M.T. Darvishi, A. Karami, *Application of He's parameter-expansion method to a nonlinear self-excited oscillator system*, Int. J. Nonlin. Sci. Num. Simul., 10(1) (2009) 137–143.
- [14] M.T. Darvishi, S. Kheybari, A. Yildirim, *Application of He's parameter-expansion method to a system of two van der Pol oscillators coupled via a bath*, Nonlin. Science Lett. A., 1(4) (2010) 399–405.
- [15] M.T. Darvishi, *Preconditioning and domain decomposition schemes to solve PDEs*, Int'l J. of Pure and Applied Math., 1(4) (2004) 419–439.
- [16] M.T. Darvishi, S. Kheybari and F. Khani, *A numerical solution of the Korteweg-de Vries equation by pseudospectral method using Darvishi's preconditionings*, Appl. Math. Comput., 182(1) (2006) 98–105.
- [17] M.T. Darvishi, M. Javidi, *A numerical solution of Burgers' equation by pseudospectral method and Darvishi's preconditioning*, Appl. Math. Comput., 173(1) (2006) 421–429.
- [18] M.T. Darvishi, F. Khani and S. Kheybari, *Spectral collocation solution of a generalized Hirota-Satsuma KdV equation*, Int. J. Comput. Math., 84(4) (2007) 541–551.
- [19] M.T. Darvishi, F. Khani, S. Kheybari, *Spectral collocation method and Darvishi's preconditionings to solve the generalized Burgers-Huxley equation*, Commun., Nonlinear Sci. Numer. Simul., 13(10) (2008) 2091–2103.
- [20] S.J. Liao, *An explicit, totally analytic approximate solution for Blasius viscous flow problems*, Int. J. Non-Linear Mech., 34 (1999) 759–778.
- [21] S.J. Liao, *Beyond Perturbation: Introduction to the Homotopy Analysis Method*, Chapman & Hall/CRC Press, Boca Raton, 2003.
- [22] S.J. Liao, *On the homotopy analysis method for nonlinear problems*, Appl. Math. Comput., 147 (2004) 499–513.
- [23] S.J. Liao, *A new branch of solutions of boundary-layer flows over an impermeable stretched plate*, Int. J. Heat Mass Transfer, 48 (2005) 2529–2539.

- [24] S.J. Liao, *A general approach to get series solution of non-similarity boundary-layer flows*, Commun. Nonlinear Sci. Numer. Simul., 14(5) (2009) 2144–2159.
- [25] M.T. Darvishi, F. Khani, *A series solution of the foam drainage equation*, Comput. Math. Appl., 58 (2009) 360–368.
- [26] J.H. He, X.H. Wu, *Construction of solitary solution and compacton-like solution by variational iteration method*, Chaos, Solitons and Fractals, 29 (2006) 108–113.
- [27] F. Khani, S. Hamed-Nezhad, M.T. Darvishi, S.-W. Ryu, *New solitary wave and periodic solutions of the foam drainage equation using the Exp-function method*, Nonlin. Anal.: Real World Appl., 10 (2009) 1904–1911.
- [28] B.-C. Shin, M.T. Darvishi, A. Barati, *Some exact and new solutions of the Nizhnik-Novikov-Vesselov equation using the Exp-function method*, Comput. Math. Appl. 58(11/12) (2009) 2147–2151.
- [29] X.H. Wu, J.H. He, *Exp-function method and its application to nonlinear equations*, Chaos, Solitons and Fractals, 38(3) (2008) 903–910.
- [30] J.H. He, X.H. Wu, *Exp-function method for nonlinear wave equations*, Chaos Solitons Fractals, 30(3) (2006) 700–708.
- [31] J.H. He, M.A. Abdou, *New periodic solutions for nonlinear evolution equations using Exp-function method*, Chaos Solitons Fractals, 34 (2007) 1421–1429.
- [32] Ma S-H, Peng J and Zhang C, *New exact solutions of the (2+1)-dimensional breaking soliton system via an extended mapping method*, Chaos Solitons Fractals, 46 (2009) 210–214.
- [33] A.M. Wazwaz, *Integrable (2+1)-dimensional and (3+1)-dimensional breaking soliton equations*, Phys. Scr., 81 (2010) 1–5.
- [34] M.-L. Wang., X.-Z. Li, J.-L. Zhang, *The $(\frac{G'}{G})$ -expansion method and travelling wave solutions of nonlinear evolution equations in mathematical physics*, Physics Letters A, 372 (2008) 417–423.
- [35] M.T. Darvishi, M. Najafi, M. Najafi, *New exact solutions for the (3+1)-dimensional breaking soliton equation*, International Journal of Engineering and Mathematical Sciences, 6(2) (2010) 137–140.
- [36] J. Wang, *Construction of new exact traveling wave solutions to (2+1)-dimensional mVN equation*, International Journal of Nonlinear Science, 9(3) (2010) 325–329.