

# The Bipartite Ramsey Numbers $b(C_{2m}; C_{2n})$

Rui Zhang, Yongqi Sun, and Yali Wu,

**Abstract**—Given bipartite graphs  $H_1$  and  $H_2$ , the bipartite Ramsey number  $b(H_1; H_2)$  is the smallest integer  $b$  such that any subgraph  $G$  of the complete bipartite graph  $K_{b,b}$ , either  $G$  contains a copy of  $H_1$  or its complement relative to  $K_{b,b}$  contains a copy of  $H_2$ . It is known that  $b(K_{2,2}; K_{2,2}) = 5$ ,  $b(K_{2,3}; K_{2,3}) = 9$ ,  $b(K_{2,4}; K_{2,4}) = 14$  and  $b(K_{3,3}; K_{3,3}) = 17$ . In this paper we study the case that both  $H_1$  and  $H_2$  are even cycles, prove that  $b(C_{2m}; C_{2n}) \geq m + n - 1$  for  $m \neq n$ , and  $b(C_{2m}; C_6) = m + 2$  for  $m \geq 4$ .

**Keywords**—bipartite graph; Ramsey number; even cycle

## I. INTRODUCTION

WE consider only finite undirected graphs without loops or multiple edges. For a graph  $G$  with vertex-set  $V(G)$  and edge-set  $E(G)$ , we denote the order and the size of  $G$  by  $p(G) = |V(G)|$  and  $q(G) = |E(G)|$ .  $\delta(G)$  and  $\Delta(G)$  are the minimum degree and the maximum degree of  $G$  respectively.

Let  $K_{m,n}$  be a complete  $m$  by  $n$  bipartite graph, that is,  $K_{m,n}$  consists of  $m + n$  vertices, partitioned into sets of size  $m$  and  $n$ , and the  $mn$  edges between them.  $P_k$  is a path on  $k$  vertices, and  $C_k$  is a cycle of length  $k$ . Let  $H_1$  and  $H_2$  be bipartite graphs, the bipartite Ramsey number  $b(H_1; H_2)$  is the smallest integer  $b$  such that given any subgraph  $G$  of the complete bipartite graph  $K_{b,b}$ , either  $G$  contains a copy of  $H_1$  or there exists a copy of  $H_2$  in the complement of  $G$  relative to  $K_{b,b}$ . Obviously, we have  $b(H_1; H_2) = b(H_2; H_1)$ .

Beineke and Schwenk<sup>[1]</sup> showed that  $b(K_{2,2}; K_{2,2}) = 5$ ,  $b(K_{2,4}; K_{2,4}) = 13$ ,  $b(K_{3,3}; K_{3,3}) = 17$ . In particular, they proved that  $b(K_{2,n}; K_{2,n}) = 4n - 3$  for  $n$  odd and less than 100 except  $n = 59$  or  $n = 95$ . Carnielli and Carmelo<sup>[2]</sup> proved that  $b(K_{2,n}; K_{2,n}) = 4n - 3$  if  $4n - 3$  is a prime power. They also showed that  $b(K_{2,2}; K_{1,n}) = n + q$  for  $q^2 - q + 1 \leq n \leq q^2$ , where  $q$  is a prime power. Irving<sup>[6]</sup> showed that  $b(K_{4,4}; K_{4,4}) \leq 48$ . Hattingh and Henning<sup>[4]</sup> proved that  $b(K_{2,2}; K_{3,3}) = 9$ ,  $b(K_{2,2}; K_{4,4}) = 14$ . They also determined the values of  $b(P_m; K_{1,n})$ <sup>[5]</sup>. Faudree and Schelp proved the values of  $b(H_1; H_2)$  when both  $H_1$  and  $H_2$  are two paths<sup>[3]</sup>. It was shown that  $b(C_6; K_{2,2}) = 5$  and  $b(C_{2m}; K_{2,2}) = m + 1$  for  $m \geq 4$  in [7].

Let  $G_i$  be the subgraph of  $G$  whose edges are in the  $i$ -th color in an  $r$ -coloring of the edges of  $G$ . If there exists an  $r$ -coloring of the edges of  $G$  such that  $H_i \not\subseteq G_i$  for all  $1 \leq i \leq r$ , then  $G$  is said to be  $r$ -colorable to  $(H_1, H_2, \dots, H_r)$ . The neighborhood of a vertex  $v \in V(G)$  are denoted by  $N(v) = \{u \in V(G) | uv \in E(G)\}$ , and let  $d(v) = |N(v)|$ .  $G^c$

Yongqi Sun is with the School of Computer and Information Technology, Beijing Jiaotong University, Beijing, 100044, P. R. China, e-mail: yqsun@bjtu.edu.cn.

Rui Zhang and Yali Wu are doctoral students of the School of Computer and Information Technology, Beijing Jiaotong University, Beijing, 100044, P. R. China.

denotes the complement of  $G$  relative to  $K_{b,b}$ .  $G\langle W \rangle$  denotes the subgraph of  $G$  induced by  $W \subseteq V(G)$ . Let  $G \cup H$  denote a disjoint sum of  $G$  and  $H$ , and  $nG$  is a disjoint sum of  $n$  copies of  $G$ .

Obviously, if  $H_1$  and  $H_2$  are cycles, then they are both even cycles. In this paper we study the case that both  $H_1$  and  $H_2$  are even cycles. Firstly, we prove that  $b(C_{2m}; C_{2n}) \geq m + n - 1$  for  $m \neq n$  and  $b(C_{2m}; C_{2m}) \geq 2m$ . Then setting  $n = 3$ , we prove that  $b(C_6; C_6) = 6$  and  $b(C_{2m}; C_6) = m + 2$  for  $m \geq 4$ . For the sake of convenience, let  $V(K_{m,n}) = X \cup Y$ , where  $X = \{x_i | 1 \leq i \leq m\}$ ,  $Y = \{y_j | 1 \leq j \leq n\}$ , and  $E(K_{m,n}) = \{x_i y_j | 1 \leq i \leq m, 1 \leq j \leq n\}$ .

## II. THE LOWER BOUNDS OF $b(C_{2m}; C_{2n})$

**Theorem 1:**  $b(C_{2m}; C_{2n}) \geq \begin{cases} m + n - 1, & m \neq n, \\ 2m, & m = n. \end{cases}$

**Proof:** If  $m \neq n$ , let  $G_1$  and  $G_2$  be subgraphs of  $K_{m+n-2, m+n-2}$ , where  $G_1$  is a complete  $m-1$  by  $m+n-2$  bipartite graph, and  $G_2$  is a complete  $n-1$  by  $m+n-2$  bipartite graph. And let  $V(G_1) = X_1 \cup Y$ , where  $X_1 = \{x_i | 1 \leq i \leq m-1\}$  and  $Y = \{y_j | 1 \leq j \leq m+n-2\}$ ;  $V(G_2) = X_2 \cup Y$ , where  $X_2 = \{x_i | m \leq i \leq m+n-2\}$ ,  $Y = \{y_j | 1 \leq j \leq m+n-2\}$ . Then we have  $E(G_1) \cap E(G_2) = \emptyset$  and  $E(G_1) \cup E(G_2) = E(K_{m+n-2, m+n-2})$ . Note that  $C_{2m} \not\subseteq G_1$  and  $C_{2n} \not\subseteq G_2$ . So  $K_{m+n-2, m+n-2}$  is 2-colorable to  $(C_{2m}, C_{2n})$ , that is,  $b(C_{2m}; C_{2n}) \geq m + n - 1$ .

If  $m = n$ , let  $G_1$  and  $G_2$  be the spanning subgraphs of  $K_{2m-1, 2m-1}$ . And let  $E(G_1) = \{x_i y_j | 1 \leq i, j \leq m-1\} \cup \{x_i y_j | m \leq i, j \leq 2m-2\} \cup \{x_{2m-1} y_j | 1 \leq j \leq 2m-1\}$ ;  $E(G_2) = \{x_i y_j | 1 \leq i \leq m-1, m \leq j \leq 2m-2\} \cup \{x_i y_j | m \leq i \leq 2m-2, 1 \leq j \leq m-1\} \cup \{x_i y_{2m-1} | 1 \leq i \leq 2m-2\}$ . Then we have  $E(G_1) \cap E(G_2) = \emptyset$  and  $E(G_1) \cup E(G_2) = E(K_{2m-1, 2m-1})$ . Note that  $C_{2m} \not\subseteq G_1$  and  $C_{2m} \not\subseteq G_2$ . So  $K_{2m-1, 2m-1}$  is 2-colorable to  $(C_{2m}, C_{2m})$ , that is,  $b(C_{2m}; C_{2m}) \geq 2m$ . ■

Setting  $n = 3$  in Theorem 1, we have

**Corollary 1:**  $b(C_{2m}; C_6) \geq \begin{cases} m + 2, & m \neq 3, \\ 6, & m = 3. \end{cases}$


## III. THE UPPER BOUNDS OF $b(C_{2m}; C_6)$ ( $m \geq 3$ )

**Lemma 1:** Let  $G$  be a spanning subgraph of  $K_{3,3}$ , if  $C_6 \not\subseteq G^c$ , then  $P_3 \subseteq G$ .

**Proof:** If  $P_3 \not\subseteq G$ , then  $G$  is isomorphic to one graph of  $\{6P_1, 4P_1 \cup P_2, 2P_1 \cup 2P_2, 3P_2\}$ . In any case, we have  $C_6 \subseteq G^c$ . ■


**Lemma 2:**  $b(C_6; C_6) \leq 6$ .

**Proof:** By contradiction, we assume that  $b(C_6; C_6) > 6$ , that is,  $K_{6,6}$  is 2-colorable to  $C_6$ . Let  $V(K_{5,5}) = V(K_{6,6}) - \{x_6, y_6\}$ . By Theorem 1,  $K_{5,5}$  is 2-colorable to  $C_6$ , and  $E(G_1 \langle V(K_{5,5}) \rangle) = \{x_i y_j | 1 \leq i, j \leq 2\} \cup \{x_i y_j | 3 \leq i, j \leq$

Fig. 1. The graphs  $G'_1(V(K_{5,5}))$  and  $G'_2(V(K_{5,5}))$ 

$4\} \cup \{x_5y_j | 1 \leq j \leq 5\}; E(G_2(V(K_{5,5}))) = \{x_iy_j | 1 \leq i \leq 2, 3 \leq j \leq 4\} \cup \{x_iy_j | 3 \leq i \leq 4, 1 \leq j \leq 2\} \cup \{x_iy_5 | 1 \leq i \leq 4\}$ . Besides this, there is one coloring way without resulting monosubgraph  $C_6$  in the 2-coloring edges of  $K_{5,5}$ , namely  $G'_1(V(K_{5,5})) \cong G_1(V(K_{5,5})) - x_5y_4$  and  $G'_2(V(K_{5,5})) \cong G_2(V(K_{5,5})) + x_5y_4$  (see Figure 1). Now we consider the vertices  $x_6$  and  $y_6$ . Since  $C_6 \not\subseteq G_2$  (or  $G'_2$ ),  $x_6$  is adjacent to at most one vertex of  $\{y_1, y_2, y_3, y_4\}$ . Hence  $x_6$  has to be adjacent to at least three vertices of  $\{y_1, y_2, y_3, y_4\}$  in  $G_1$  (or  $G'_1$ ), we have  $C_6 \subseteq G_1$  (or  $G'_1$ ), a contradiction. So,  $K_{6,6}$  is not 2-colorable to  $C_6$ , that is,  $b(C_6; C_6) \leq 6$ . ■

In order to prove Lemma 3, we need the following claims. Let  $H_{2k+3}$  and  $H_{2k+4}$  denote the two graphs as shown in Figure 2, and  $G$  be a spanning subgraph of  $K_{k+3, k+3}$  for  $k \geq 3$  such that  $C_{2(k+1)} \not\subseteq G$  and  $C_6 \not\subseteq G^c$ , then we have

Fig. 2. The graphs  $H_{2k+3}$  and  $H_{2k+4}$ 

**Claim 1:**  $H_{2k+3} \not\subseteq G$ .

**Proof:** By contradiction, we assume that  $H_{2k+3} \subseteq G$ , and label the vertices of  $H_{2k+3}$  as shown in Fig. 2(a).

Let  $x_{k+2}$ ,  $x_{k+3}$  and  $y_{k+3}$  be the remaining vertices of  $V(G)$ . Since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G(\{x_{k+1}, x_{k+2}, x_{k+3}, y_{k-1}, y_k, y_{k+3}\})$ . Since  $C_{2(k+1)} \not\subseteq G$ ,  $x_{k+1}$  is nonadjacent to  $y_{k-1}$  or  $y_k$ . By symmetry, it is sufficient to consider the five cases. We may assume  $x_{k+2}y_{k-1}, x_{k+2}y_{k+3} \in E(G)$ ,  $y_{k-1}x_{k+2}, y_{k-1}x_{k+3} \in E(G)$ ,  $x_{k+2}y_{k-1}, x_{k+2}y_k \in E(G)$ ,  $y_{k+3}x_{k+1}, y_{k+3}x_{k+2} \in E(G)$  or  $y_{k+3}x_{k+2}, y_{k+3}x_{k+3} \in E(G)$ .


**Case 1.** Suppose  $x_{k+2}y_{k-1}, x_{k+2}y_{k+3} \in E(G)$ . Since  $C_{2(k+1)} \not\subseteq G$ , each vertex of  $\{y_{k+1}, y_{k+2}\}$  is nonadjacent to any vertex of  $\{x_1, x_{k-1}, x_{k+2}\}$ , and  $y_{k+3}$  is nonadjacent to  $x_{k-1}$ . And since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G(\{x_1, x_{k-1}, x_{k+2}, y_{k+1}, y_{k+2}, y_{k+3}\})$ . Hence  $y_{k+3}$  has to be adjacent to  $x_1$ . Since  $C_{2(k+1)} \not\subseteq G$ ,  $y_k$  is nonadjacent to any vertex of  $\{x_{k-1}, x_{k+2}\}$ . Hence we have we have  $P_3 \not\subseteq G(\{x_1, x_{k-1}, x_{k+2}, y_k, y_{k+1}, y_{k+2}\})$ . By Lemma 1, we have  $C_6 \subseteq G^c$ , a contradiction.

**Case 2.** Suppose  $y_{k-1}x_{k+2}, y_{k-1}x_{k+3} \in E(G)$ . Since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G(\{x_{k-1}, x_{k+2}, x_{k+3}, y_{k+1}, y_{k+2}, y_{k+3}\})$ . Since  $C_{2(k+1)} \not\subseteq G$ , each vertex of  $\{y_{k+1}, y_{k+2}\}$  is nonadjacent to any vertex of  $\{x_{k-1}, x_{k+2}, x_{k+3}\}$ . Hence  $y_{k+3}$  has to be adjacent to at least one vertex of  $\{x_{k+2}, x_{k+3}\}$ . The proof is same as Case 1.

**Case 3.** Suppose  $x_{k+2}y_{k-1}, x_{k+2}y_k \in E(G)$ . Since  $C_{2(k+1)} \not\subseteq G$ , each vertex of  $\{y_{k+1}, y_{k+2}\}$  is nonadjacent to any vertex of  $\{x_1, x_{k-1}, x_{k+2}\}$ . And since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G(\{x_1, x_{k-1}, x_{k+2}, y_{k+1}, y_{k+2}, y_{k+3}\})$ . Hence  $y_{k+3}$  is adjacent to at least two vertices of  $\{x_1, x_{k-1}, x_{k+2}\}$ . Therefore since  $C_{2(k+1)} \not\subseteq G$ , we have  $y_{k+3}$  has to be adjacent to  $x_1$  and  $x_{k-1}$ . Similarly, since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G(\{x_k, x_{k+1}, x_{k+3}, y_1, y_k, y_{k+3}\})$ . Since  $C_{2(k+1)} \not\subseteq G$ ,  $x_k$  is nonadjacent to  $y_1$  or  $y_{k+3}$ , and  $x_{k+1}$  is nonadjacent to any vertex of  $\{y_1, y_k, y_{k+3}\}$ . If  $x_{k+3}$  is adjacent to  $y_k$ , the proof is same as Case 2. If  $x_{k+3}$  is adjacent to both  $y_1$  and  $y_{k+3}$ , we have  $C_{2(k+1)} \subseteq G$ , a contradiction.

**Case 4.** Suppose  $y_{k+3}x_{k+1}, y_{k+3}x_{k+2} \in E(G)$ . Since  $C_{2(k+1)} \not\subseteq G$ , each vertex of  $\{x_1, x_{k-1}\}$  is nonadjacent to any vertex of  $\{y_{k+1}, y_{k+2}, y_{k+3}\}$ . And since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G(\{x_1, x_{k-1}, x_{k+2}, y_{k+1}, y_{k+2}, y_{k+3}\})$ . Hence  $x_{k+2}$  is adjacent to at least one vertex of  $\{y_{k+1}, y_{k+2}\}$ , say  $x_{k+2}y_{k+1} \in E(G)$  as shown in Fig. 3. And since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G(\{x_1, x_{k-1}, x_{k+3}, y_{k+1}, y_{k+2}, y_{k+3}\})$ . Hence we have  $x_{k+3}$  is adjacent to at least two vertices of  $\{y_{k+1}, y_{k+2}, y_{k+3}\}$ . In any case, since  $C_{2(k+1)} \not\subseteq G$ ,  $x_{k+3}$  is nonadjacent to any vertex of  $\{y_{k-2}, y_{k-1}, y_k\}$ . And each vertex of  $\{x_{k+1}, x_{k+2}\}$  is nonadjacent to any vertex of  $\{y_{k-2}, y_{k-1}, y_k\}$ . Hence we have  $P_3 \not\subseteq G(\{x_{k+1}, x_{k+2}, x_{k+3}, y_{k-2}, y_{k-1}, y_k\})$ . By Lemma 1, we have  $C_6 \subseteq G^c$ , a contradiction.

**Case 5.** Suppose  $y_{k+3}x_{k+2}, y_{k+3}x_{k+3} \in E(G)$ . Since  $C_{2(k+1)} \not\subseteq G$ ,  $x_{k+1}$  is nonadjacent to  $y_{k-1}$  or  $y_k$ . Since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G(\{x_{k+1}, x_{k+2}, x_{k+3}, y_{k-1}, y_k, y_{k+1}\})$ . If there is one edge between  $\{x_{k+2}, x_{k+3}\}$  and  $\{y_{k-1}, y_k\}$ , the proof is same as Case 2. Hence  $y_{k+1}$  has to be adjacent to at least one vertex of  $\{x_{k+2}, x_{k+3}\}$ , say  $y_{k+1}x_{k+2} \in E(G)$ . And since  $C_{2(k+1)} \not\subseteq G$ ,  $y_1$  is nonadjacent to  $x_{k+2}$  or  $x_{k+3}$ . Therefore we have  $P_3 \not\subseteq G(\{x_{k+1}, x_{k+2}, x_{k+3}, y_1, y_{k-1}, y_k\})$ . By Lemma 1, we have

Fig. 3.  $x_{k+2}$  being adjacent to  $y_{k+1}$ 

$C_6 \subseteq G^c$ , a contradiction.

By Cases 1-5, we have  $H_{2k+3} \not\subseteq G$ . ■

**Claim 2:**  $H_{2k+4} \not\subseteq G$ .

*Proof:* By contradiction, we assume that  $H_{2k+4} \subseteq G$ , and label the vertices of  $H_{2k+4}$  as shown in Fig. 2(b). Let  $x_{k+2}$  and  $x_{k+3}$  be the remaining vertices of  $V(G)$ . Since  $C_{2(k+1)} \not\subseteq G$ ,  $x_{k+1}$  is nonadjacent to any vertex of  $\{y_{k-1}, y_k\}$ . If  $x_{k+1}$  is adjacent to  $y_{k+3}$ , then we have  $H_{2k+3} \subseteq G$ , a contradiction to Claim 1. And since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G\langle\{x_{k+1}, x_{k+2}, x_{k+3}, y_{k-1}, y_k, y_{k+3}\}\rangle$ . If  $x_{k+2}$  (or  $x_{k+3}$ ) is adjacent to both  $y_{k-1}$  and  $y_{k+3}$ , we have  $C_{2(k+1)} \subseteq G$ , a contradiction. By symmetry, we may assume  $x_{k+2}y_{k-1}, x_{k+2}y_k \in E(G)$ ,  $y_{k-1}x_{k+2}, y_{k-1}x_{k+3} \in E(G)$  or  $y_{k+3}x_{k+2}, y_{k+3}x_{k+3} \in E(G)$ .

**Case 1.** Suppose  $x_{k+2}y_{k-1}, x_{k+2}y_k \in E(G)$ . Since  $C_{2(k+1)} \not\subseteq G$ , each vertex of  $\{y_{k+1}, y_{k+3}\}$  is nonadjacent to any vertex of  $\{x_1, x_{k-1}, x_{k+2}\}$ , and  $y_{k+2}$  is nonadjacent to any vertex of  $\{x_1, x_{k-1}\}$ . Then  $P_3 \not\subseteq G\langle\{x_1, x_{k-1}, x_{k+2}, y_{k+1}, y_{k+2}, y_{k+3}\}\rangle$ . By Lemma 1, we have  $C_6 \subseteq G^c$ , a contradiction.

**Case 2.** Suppose  $y_{k-1}x_{k+2}, y_{k-1}x_{k+3} \in E(G)$ . Since  $C_{2(k+1)} \not\subseteq G$ , each vertex of  $\{y_{k+1}, y_{k+3}\}$  is nonadjacent to any vertex of  $\{x_{k+2}, x_{k+3}\}$ , and  $y_k$  is nonadjacent to  $x_{k+1}$ . If  $y_k$  is adjacent to one vertex of  $\{x_{k+2}, x_{k+3}\}$ , the proof is same as Case 1. If  $x_{k+1}y_{k+3} \in E(G)$ , then  $H_{2k+3} \subseteq G$ , a contradiction to Claim 1. Hence  $P_3 \not\subseteq G\langle\{x_{k+1}, x_{k+2}, x_{k+3}, y_k, y_{k+1}, y_{k+3}\}\rangle$ . By Lemma 1, we have  $C_6 \subseteq G^c$ , a contradiction.

**Case 3.** Suppose  $y_{k+3}x_{k+2}, y_{k+3}x_{k+3} \in E(G)$ . Since  $C_{2(k+1)} \not\subseteq G$ , each vertex of  $\{y_{k-1}, y_k\}$  is nonadjacent to any vertex of  $\{x_{k+1}, x_{k+2}, x_{k+3}\}$ . And since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G\langle\{x_{k+1}, x_{k+2}, x_{k+3}, y_{k-1}, y_k, y_{k+1}\}\rangle$ . Hence  $y_{k+1}$  is adjacent to at least one vertex of  $\{x_{k+2}, x_{k+3}\}$ . In any case, we have  $H_{2k+3} \subseteq G$ , a contradiction to Claim 1.

By Cases 1-3, we have  $H_{2k+4} \not\subseteq G$ . ■

By an argument similar to the above proofs, we can prove Claim 3 and 4. However, their proofs are more complicated than Claim 2.

**Claim 3:**  $(C_{2k} \cup C_4) \not\subseteq G$ .


**Claim 4:**  $(C_{2k} \cup P_5) \not\subseteq G$ .

**Lemma 3:** Let  $G$  be a spanning subgraph of  $K_{k+3, k+3}$  for  $k \geq 3$ . If  $C_{2k} \subseteq G$  and  $C_6 \not\subseteq G^c$ , then  $C_{2(k+1)} \subseteq G$ .

*Proof:* We may assume that  $C_{2(k+1)} \not\subseteq G$ . Without loss of generality, let  $E(C_{2k}) = \{x_1y_1, y_1x_2, x_2y_2, \dots, x_ky_k, y_kx_1\}$ . Since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G\langle\{x_{k+1}, x_{k+2}, x_{k+3}, y_{k+1}, y_{k+2}, y_{k+3}\}\rangle$ , say  $x_{k+1}y_{k+1}, x_{k+1}y_{k+2} \in E(G)$ .

Similarly, since  $C_6 \not\subseteq G^c$ , we have  $P_3 \subseteq G\langle\{x_k, x_{k+2}, x_{k+3}, y_{k+1}, y_{k+2}, y_{k+3}\}\rangle$ . If  $x_k$  is adjacent to both  $y_{k+1}$  and  $y_{k+2}$ , then  $H_{2k+3} \subseteq G$ , a contradiction to Claim 1. If  $x_k$  is adjacent to both  $y_{k+1}$  and  $y_{k+3}$  (or both  $y_{k+2}$  and  $y_{k+3}$ ), then  $H_{2k+4} \subseteq G$ , a contradiction to Claim 2. If there exists one vertex of  $\{x_{k+2}, x_{k+3}\}$  being adjacent to both  $y_{k+1}$  and  $y_{k+2}$ , then  $(C_{2k} \cup C_4) \subseteq G$ , a contradiction to Claim 3. If there exists one vertex of  $\{x_{k+2}, x_{k+3}\}$  being adjacent to both  $y_{k+1}$  and  $y_{k+3}$  (or both  $y_{k+2}$  and  $y_{k+3}$ ), then  $(C_{2k} \cup P_5) \subseteq G$ , a contradiction to Claim 4. So, by symmetry, it is sufficient to consider the four cases as follows.

**Case 1.** Suppose  $y_{k+1}x_k, y_{k+1}x_{k+2} \in E(G)$ . Since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G\langle\{x_{k+1}, x_{k+2}, x_{k+3}, y_{k-1}, y_k, y_{k+2}\}\rangle$ . Since  $C_{2(k+1)} \not\subseteq G$ , each vertex of  $\{x_{k+1}, x_{k+2}\}$  is nonadjacent to any vertex of  $\{y_{k-1}, y_k\}$ . If  $x_{k+2}$  is adjacent to  $y_{k+2}$ , then we have  $(C_{2k} \cup C_4) \subseteq G$ , a contradiction to Claim 3. If  $x_{k+3}$  is adjacent to  $y_{k+2}$ , then we have  $(C_{2k} \cup P_5) \subseteq G$ , a contradiction to Claim 4. Hence  $x_{k+3}$  has to be adjacent to both  $y_{k-1}$  and  $y_k$ . Similarly since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G\langle\{x_1, x_{k-1}, x_{k+3}, y_{k+1}, y_{k+2}, y_{k+3}\}\rangle$ . Since  $C_{2(k+1)} \not\subseteq G$ ,  $y_{k+1}$  is nonadjacent to any vertex of  $\{x_1, x_{k-1}, x_{k+3}\}$ ,  $y_{k+2}$  is nonadjacent to any vertex of  $\{x_1, x_{k-1}\}$ . If  $y_{k+2}x_{k+3} \in E(G)$ , we have  $(C_{2k} \cup P_5) \subseteq G$ , a contradiction to Claim 4. If  $y_{k+3}$  is adjacent to both  $x_1$  and  $x_{k+3}$  (or both  $x_{k-1}$  and  $x_{k+3}$ ), we have  $C_{2(k+1)} \subseteq G$ , a contradiction too. Hence we have  $y_{k+3}x_1, y_{k+3}x_{k-1} \in E(G)$  as shown in Fig. 4. However, since  $C_{2(k+1)} \not\subseteq G$ , each vertex of  $\{x_{k+1}, x_{k+2}\}$  is nonadjacent to any vertex of  $\{y_1, y_{k-1}, y_{k+3}\}$  and  $x_{k+3}$  is nonadjacent to any vertex of  $\{y_1, y_{k+3}\}$ . So, we have  $P_3 \not\subseteq G\langle\{x_{k+1}, x_{k+2}, x_{k+3}, y_1, y_{k-1}, y_{k+3}\}\rangle$ . By Lemma 1, we have  $C_6 \subseteq G^c$ , a contradiction.

Fig. 4.  $y_{k+3}$  being adjacent to both  $x_1$  and  $x_{k-1}$ 

**Case 2.** Suppose  $y_{k+1}x_{k+2}, y_{k+1}x_{k+3} \in E(G)$ . Since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G\langle\{x_{k+1}, x_{k+2}, x_{k+3}, y_k, y_{k+2}, y_{k+3}\}\rangle$ . If  $x_{k+1}$  is adjacent to  $y_k$ , the proof is same as Case 1. If there exists one vertex of  $\{x_{k+2}, x_{k+3}\}$  being adjacent to  $y_{k+2}$ , then we have  $(C_{2k} \cup C_4) \subseteq G$ , a contradiction to Claim 3. If there exists one vertex of  $\{x_{k+2}, x_{k+3}\}$  being adjacent to  $y_{k+3}$ , then we have  $(C_{2k} \cup P_5) \subseteq G$ , a contradiction to Claim 4. If  $y_k$  is adjacent to both  $x_{k+2}$  and  $x_{k+3}$ , we have  $H_{2k+3} \subseteq G$ , a contradiction to Claim 1. Hence  $y_{k+3}$  has to be adjacent to  $x_{k+1}$ . Similarly, since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G\langle\{x_1, x_k, x_{k+2}, y_{k+1}, y_{k+2}, y_{k+3}\}\rangle$ . If there exists one vertex of  $\{x_1, x_k\}$  being adjacent to  $y_{k+1}$ , the proof is same as Case 1. If there exists one vertex of  $\{x_1, x_k\}$  being adjacent to both  $y_{k+2}$  and  $y_{k+3}$ , then we have  $H_{2k+3} \subseteq G$ , a contradiction to Claim 1. If  $x_{k+2}$  is adjacent to  $y_{k+2}$  or  $y_{k+3}$ , then we have  $(C_{2k} \cup C_4) \subseteq G$ , a contradiction to Claim 3. If

there exists one vertex of  $\{y_{k+2}, y_{k+3}\}$  being adjacent to both  $x_1$  and  $x_k$ , the proof is same as Case 1.

Case 3. Suppose  $y_{k+3}x_k, y_{k+3}x_{k+2} \in E(G)$ . And since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G[\{x_k, x_{k+2}, x_{k+3}, y_{k-1}, y_{k+1}, y_{k+2}\}]$ . If  $x_k$  is adjacent to  $y_{k+1}$  or  $y_{k+2}$ , then we have  $H_{2k+4} \subseteq G$ , a contradiction to Claim 2. If  $x_{k+2}$  is adjacent to  $y_{k+1}$  or  $y_{k+2}$ , then we have  $(C_{2k} \cup P_5) \subseteq G$ , a contradiction to Claim 4. If  $x_{k+3}$  is adjacent to both  $y_{k+1}$  and  $y_{k+2}$ , then we have  $(C_{2k} \cup C_4) \subseteq G$ , a contradiction to Claim 3. Since  $C_{2(k+1)} \not\subseteq G$ ,  $y_{k-1}$  is non-adjacent to  $x_{k+2}$ . Hence  $x_{k+3}$  has to be adjacent to  $y_{k-1}$ . Similarly, we have  $y_kx_{k+3} \in E(G)$ , since otherwise  $P_3 \not\subseteq G[\{x_k, x_{k+2}, x_{k+3}, y_k, y_{k+1}, y_{k+2}\}]$ .

Since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G[\{x_1, x_{k+2}, x_{k+3}, y_{k+1}, y_{k+2}, y_{k+3}\}]$ . If there exists one vertex of  $\{x_1, x_{k+3}\}$  being adjacent to both  $y_{k+1}$  and  $y_{k+2}$ , then we have  $H_{2k+3} \subseteq G$ , a contradiction to Claim 1. If  $x_{k+2}$  is adjacent to  $y_{k+1}$  or  $y_{k+2}$ , then we have  $(C_{2k} \cup P_5) \subseteq G$ , a contradiction to Claim 4. Since  $C_{2(k+1)} \not\subseteq G$ ,  $y_{k+3}$  is nonadjacent to  $x_1$  or  $x_{k+3}$ . If there exists one vertex of  $\{y_{k+1}, y_{k+2}\}$  being adjacent to both  $x_1$  and  $x_{k+3}$ , we have  $C_{2(k+1)} \subseteq G$ , a contradiction.

Case 4. Suppose  $y_{k+3}x_{k+2}, y_{k+3}x_{k+3} \in E(G)$ . Since  $C_6 \not\subseteq G^c$ , by Lemma 1, we have  $P_3 \subseteq G[\{x_k, x_{k+2}, x_{k+3}, y_k, y_{k+1}, y_{k+2}\}]$ . If there exists one edge between  $\{x_{k+2}, x_{k+3}\}$  and  $\{y_{k+1}, y_{k+2}\}$ , we have  $(C_{2k} \cup P_5) \subseteq G$ , a contradiction to Claim 4. If  $x_k$  is adjacent to  $y_{k+1}$  or  $y_{k+2}$ , the proof is same as Case 3. If  $y_k$  is adjacent to  $x_{k+2}$  or  $x_{k+3}$ , the proof is also same as Case 3.

By Cases 1-4, we have  $C_{2k+1} \subseteq G$ . ■

Let  $G$  be a spanning subgraph of  $K_{6,6}$ . If  $C_6 \not\subseteq G^c$ , by Lemma 2, we have  $C_6 \subseteq G$ . Hence we have the following corollary by Lemma 3.

Corollary 2:  $b(C_8; C_6) \leq 6$ .

Lemma 4: If  $m \geq 4$ , we have  $b(C_{2m}; C_6) \leq m + 2$ .

Proof: We will prove it by induction.

(1) For  $m = 4$ , the lemma holds by Corollary 2.

(2) Suppose that  $b(C_{2k}; C_6) \leq k + 2$  for  $k \geq 5$ . We assume that  $b(C_{2(k+1)}; C_6) > k + 3$  for  $k \geq 5$ . Since  $C_6 \not\subseteq G^c$ , we have  $C_{2k} \subseteq G$ . By Lemma 3, we have  $C_{2(k+1)} \subseteq G$ , a contradiction. So the assumption does not hold, that is,  $b(C_{2(k+1)}; C_6) \leq k + 3$ . This completes the induction step, and the proof is finished. ■

#### IV. CONCLUSION

Setting  $m = 3$  in Corollary 1, we have  $b(C_6; C_6) \geq 6$ . By Theorem 1, Lemma 2 and Lemma 4, we obtain the values of  $b(C_{2m}; C_6)$  as follows.

Theorem 2:  $b(C_{2m}; C_6) = \begin{cases} 6, & m = 3, \\ m + 2, & m \geq 4. \end{cases}$

Furthermore, we have the following conjecture,

Conjecture 1:  $b(C_{2m}; C_{2n}) = m + n - 1$  for  $m > n$ .

By the results in [7] and Theorem 2, it is true for  $n = 2$  and 3.

#### ACKNOWLEDGMENT

This research was supported by NSFC(60973011), SRFDF(20090009120007) and the Fundamental Research Funds for the Central Universities.

#### REFERENCES

- [1] L. W. Beineke and A. J. Schwenk. *On a bipartite form of the Ramsey problem*. Proc. 5th British Combin. Conf. 1975, Congressus Number, XV: 17–22, 1975.
- [2] W. A. Camielli and E. L. Monte Carmelo.  $K_{2,2} - K_{1,n}$  and  $K_{2,n} - K_{2,n}$  bipartite Ramsey numbers. Discrete Math., 223: 83–92, 2000.
- [3] R. J. Faudree and R. H. Schelp. *Path-path Ramsey-type numbers for the complete bipartite graphs*. J. Combin. Theory Ser. B, 19: 161–173, 1975.
- [4] J. H. Hattingh and M. A. Henning. *Bipartite Ramsey numbers*. Utilitas Math., 53: 217–230, 1998.
- [5] J. H. Hattingh and M. A. Henning. *Star-path bipartite Ramsey numbers*. Discrete Math., 185: 255–258, 1998.
- [6] R. W. Irving. *A bipartite Ramsey problem and the Zarankiewicz numbers*. Discrete Math., 19: 13–26, 1978.
- [7] R. Zhang and Y. Q. Sun. *The bipartite Ramsey numbers  $b(C_{2m}; K_{2,2})$* . The Elec. J. of Combin., 18: P51, 2011.