

# Fundamental Groups in Chaotic Flat Space and Its Retractions

A. E. El-Ahmady, M. Abu-Saleem

**Abstract**—The purpose of this paper is to give a combinatorial characterization and construct representations of the chaotic fundamental groups of the chaotic submanifolds of chaotic flat space by using some geometrical transformations. The chaotic homotopy groups of the limit folding for chaotic flat space are presented. The chaotic fundamental groups of some types of chaotic geodesics in chaotic flat space are deduced.

**Keywords**—Chaotic flat space, Chaotic folding, Chaotic retractions, Chaotic fundamental groups.

## I. INTRODUCTION AND DEFINITIONS

FLAT space represents one of the most intriguing and emblematic discoveries in the history of geometry. If it was introduced for a purely geometric purpose, it came into prominence in many branches of mathematics and physics. This association with applied science and geometry generated synergistic effect: Applied science gave relevance to flat space and flat space allowed formalizing practical problems [6], [8].

Vector spaces, linear maps, topological spaces, and continuous maps, groups and homomorphisms together with the distinguished family of maps are referred to a category.

An operator which assigns to every object in one category a corresponding object in another category and to every map in the first map in the second in such a way that compositions are preserved and the identity map is taken to the identity map is called a functor. Thus, we may summarize our activities thus far by saying that we have constructed a functor (the fundamental group functor) from the category of pointed spaces and maps to the category of groups and homomorphisms. Such functors are the vehicles by which one translates topological problems into algebraic problem [21]-[24], [26], [28].

Most folding problems are attractive from a pure mathematical standpoint, for the beauty of the problems themselves. The folding problems have close connections to important industrial applications. Linkage folding has applications in robotics and hydraulic tube bending. Paper folding has application in sheet-metal bending, packaging, and air-bag folding [10]-[12], [18]. Isometric folding between two Riemannian manifolds may be characterized as maps that send piecewise geodesic segments to a piecewise geodesic segments of the same length [4]. For a topological folding the

maps do not preserve lengths [5], [6], i.e. A map  $\mathfrak{S}: M \rightarrow N$ , where  $M$  and  $N$  are  $C^\infty$ -Riemannian manifolds of dimension  $m$  and  $n$  respectively is said to be an isometric folding of  $M$  into  $N$ , iff for any piecewise geodesic path  $\gamma: J \rightarrow M$ , the induced path  $\mathfrak{S} \circ \gamma: J \rightarrow N$  is a piecewise geodesic and of the same length as  $\gamma$ . If  $\mathfrak{S}$  does not preserve length, then  $\mathfrak{S}$  is a topological folding [1]-[5], [7].

An  $n$ -dimensional topological manifold  $M$  is a Hausdorff topological space with a countable basis for the topology which is locally homeomorphic to  $\mathbb{R}^n$ . If  $h: U \rightarrow U'$  is a homeomorphism of  $U \subseteq M$  onto  $U' \subseteq \mathbb{R}^n$ , then  $h$  is called a chart of  $M$  and  $U$  is the associated chart domain. A collection  $(h_\alpha, U_\alpha)$  is said to be an atlas for  $M$  if  $\bigcup_{\alpha \in A} U_\alpha = M$ . Given two charts  $h_\alpha, h_\beta$  such that  $U_{\alpha\beta} = U_\alpha \cap U_\beta \neq \emptyset$ , the transformation chart  $h_\beta \circ h_\alpha^{-1}$  between open sets of  $\mathbb{R}^n$  is defined, and if all of these charts transformation are  $C^\infty$ -mappings, then the manifolds under consideration is a  $C^\infty$ -manifolds. A differentiable structure on  $M$  is a differentiable atlas and a differentiable manifold is a topological manifold with a differentiable structure [29]-[31].  $M$  may have other structures as colors, density or any physical structures. The number of structures may be infinite. In this case the manifold is said to be a chaotic manifold and may become relevant to vacuum fluctuation and chaotic quantum field theories. The magnetic field of a magnet bar is a kind of chaotic 1-dimensional manifold represented by the magnetic flux lines. The geometric manifold is the magnetic bar itself [1]-[5].

Fuzzy manifolds are special type of the category of chaotic manifolds. Usually we denote by  $M = M_{0123\dots h}$  to a chaotic manifolds [6], [7], where  $M_{0h}$  is the geometric (essential) manifold and the associated pure chaotic manifolds, the manifolds with physical characters, are denoted by  $M_{1h}, \dots, M_{\infty h}$  [11], [13], [18], [19].

The aim of this paper is to describe the connection between the chaotic fundamental groups and the chaotic homotopy group geometrically, specifically concerned with the study of the new type of chaotic retractions, chaotic deformation retracts, chaotic foldings and the chaotic fundamental groups of chaotic flat space  $F_{0123\dots h}$  as presented by El-Ahmady [1]-[26] and M. Abu-saleem [27]-[40]. The set of chaotic homotopy classes of chaotic loops based at the point  $x_0(\mu)$  with the product operation  $[f(\mu)][g(\mu)] = [f(\mu).g(\mu)]$  is called the chaotic fundamental groups and denoted by  $\pi_1(X(\mu), x_0(\mu))$  [7], [8], [18]-[44].

A subset  $A$  of a topological space  $X$  is called a retract of  $X$  if there exists a continuous map  $r: X \rightarrow A$  such that  $r(a) = a, \forall a \in A$  where  $A$  is closed and  $X$  is open [3], [7]. Also, let  $X$  be

A. E. El-Ahmady is with the Mathematics Department, Faculty of Science, Taibah University, Madinah, Saudi Arabia, (e-mail: a\_elahmady@hotmail.com)

M. Abu-Saleem is with the Department of Mathematics, Allaiith University College (Girls Branch), Umm Al-Qura University, Allaiith, Saudi Arabia, (corresponding author to e-mail: mohammedabusaleem2005@yahoo.com).

a space and  $A$  a subspace. A map  $r: X \rightarrow A$  such that  $r(a) = a$ , for all  $a \in A$ , is called a retraction of  $X$  onto  $A$  and  $A$  is called a retract of  $X$ . This can be restated as follows. If  $i: A \rightarrow X$  is the inclusion map, then  $r: X \rightarrow A$  is a map such that  $ri = id_A$ . If, in addition,  $ri \approx id_X$ , we call  $r$  a deformation retract and  $A$  a deformation retract of  $X$ . Imagine a point light source on a two-dimensional surface. If we switch this light source on, then light will propagate in an ever increasing circle, around the source with a velocity  $C$  which is that of light. Imagine now we blot the elapsing time needed for the propagation of light on a vertical axis perpendicular to our surface with its zero point exactly at the light source point. That way we obtain the light cone which plays a fundamental role in relatively theory [25], [26]. Here we will use an E-Infinity fuzzy light cone. The light cone gives us a way of differentiating between three types of geodesics. They correspond to velocities less than  $C$ , exactly equal to  $C$  and larger than  $C$ . We call them time-like, null and space-like geodesics. Special relativity states that all matter must move at speed of light. This means that all bodies must move on time like or null geodesics i.e. unless acted upon by forces other than gravity [25], [26].

Next let us recall the concept of a metric in four-dimensions. Considering only flat space, we have

$$ds^2 = dx^2 + dy^2 + dz^2 - c^2 dt^2$$

Now we see that  $ds^2 > 0$ ,  $ds^2 = 0$  and  $ds^2 < 0$  correspond to space-like, null and time-like geodesics, we note that massless particles, such as the photon, move on null geodesics. That can be interpreted as saying that in 4-dimensional space, the photon does not move and the time for that photon does not pass. Particularly intriguing is the mathematical possibility of a negative metric. Now it is extremely interesting that there is a geometry in which two separated points may still have a zero distance analogous to  $ds^2 = 0$  the corresponding to a null geodesic [25], [26]. In what follows we would like to introduce a new type of geodesics of chaotic flat space namely "the retraction of chaotic flat space". The 0-dimensional flat space will be discussed.

## II. MAIN RESULTS

**Theorem 1.** The chaotic fundamental groups of types of the chaotic retractions of chaotic flat space  $F_{0123...h}$  are isomorphic to  $Z_{0123...h}$  or chaotic identity group  $0 = 0_{0123...h}$ .

**Proof.** The chaotic flat space is defined as

$$ds^2 = -(c_1(t)x_2 + c_2(t))^2 dx_1^2 - \frac{c_1(t)(c_1(t)x_2 + c_2(t))}{x_2 + c_3(t)} dx_2^2 - \frac{x_2 + c_3(t)}{c_1(t)(c_1(t)x_2 + c_3(t))} dx_3^2 + (c_1(t)x_2 + c_3(t))^2 \cos^2 x_1 dx_4^2 \dots \quad (1)$$

where  $c_1(t)$ ,  $c_2(t)$  and  $c_3(t)$  are functions of time. Then, the coordinates of the chaotic flat space  $F_{0123...h}$  are

$$x_1 = \frac{iA_1}{1 - i(c_1(t)x_2 + c_2(t))}, x_2 = \frac{ic_1(t)}{2} (\log B_1 - N \log B_2),$$

$$B_1 = \sqrt{x_2^2 + Lx_2 + M}, B_2 = \frac{x_2 + \frac{L}{2} + \sqrt{x_2^2 + Lx_2 + M}}{\sqrt{M - L^2/4}} \quad (2)$$

$$L = (c_3(t) + \frac{c_3(t)}{c_1(t)}), M = \frac{c_2(t)c_3(t)}{c_1(t)}, N = (c_3(t) - \frac{c_2(t)}{c_1(t)})$$

$$x_3 = \frac{iA_2}{(1 - i\sqrt{\frac{x_2 + c_3(t)}{c_1(t)(c_1(t)x_2 + c_3(t))}})}, x_4 = \frac{A_3}{(1 - (c_1(t)x_2 + c_2(t))\cos x_1)}$$

where  $A_1, A_2$  and  $A_3$  are the constant of integration.

In a position, using Lagrangian equations:

$$\frac{d}{ds} \left( \frac{\partial T}{\partial x'_i} \right) - \frac{\partial T}{\partial x_i} = 0, \quad i = 1, 2, 3, 4, 5.$$

To deduce a chaotic geodesic which is a subset of chaotic flat space  $F_{0123...h}$ . Since  $T = \frac{1}{2} \dot{ds}^2$ , this yields

$$T = \frac{1}{2} \left\{ - (c_1(t)x_2 + c_2(t))^2 - \frac{c_1^2(t)x_2 + c_1(t)c_2(t)}{x_2 + c_3(t)} x_2'^2 - \frac{x_2 + c_3(t)}{(c_1^2(t)x_2 + c_1(t)c_2(t))} x_3'^2 + (c_1(t)x_2 + c_3(t))^2 \cos^2 x_1 x_4'^2 \right\} \quad (3)$$

Then, the Lagrangian equations are

$$\frac{d}{ds} \left\{ - (c_1(t)x_2 + c_2(t))^2 x_1' + (c_1(t)x_2 + c_2(t))^2 \cos x_1 \sin x_1 x_4'^2 \right\} = 0 \quad (4)$$

$$\frac{d}{ds} \left\{ - \frac{(c_1^2(t)x_2 + c_1(t)c_2(t))}{x_2 + c_3(t)} x_2' \right\} + (c_1^2(t)x_2 + c_1(t)c_2(t)) x_1'^2 + \frac{c_1^2(t)c_3(t) - c_1(t)c_2(t)}{2(x_2 + c_3(t))^2} x_2'^2 + \frac{c_1(t)c_3(t) - c_1^2(t)c_3(t)}{2(c_1^2(t)x_2 + c_1(t)c_3(t))} x_3'^2 - (c_1^2(t)x_2 + c_1(t)c_2(t)) \cos^2 x_1 x_4'^2 = 0 \quad (5)$$

$$\frac{d}{ds} \left\{ \frac{x_2 + c_3(t)}{c_1^2(t)x_2 + c_1(t)c_3(t)} \right\} x_3' = 0 \quad (6)$$

$$\frac{d}{ds} \left\{ (c_1(t)x_2 + c_2(t))^2 \cos^2 x_1 x_4' \right\} = 0 \quad (7)$$

$$- 2(c_1(t)x_2 + c_2(t))(c_1'(t)x_2 + c_2'(t))x_4'^2 +$$

$$\frac{(x_2 + c_3(t))(2c_1(t)c_1'(t)x_2 + c_1'(t)c_2(t) + c_1(t)c_2'(t)) - (c_1^2(t)x_2 + c_1(t)c_2(t))c_3'(t)}{(x_2 + c_3(t))^2} x_2'^2 - \frac{(c_1^2(t)x_2 + c_1(t)c_2(t))c_3'(t) - (x_2 + c_3(t))(2c_1(t)c_1'(t)x_2 + c_1'(t)c_2(t) + c_1(t)c_2'(t))}{(c_1^2(t)x_2 + c_1(t)c_2(t))^2} x_3'^2 - 2(c_1(t)x_2 + c_2(t))(c_1'(t)x_2 + c_2'(t))\cos^2 x_1 x_4'^2 = 0 \quad (8)$$

From (7), we have  $(c_1(t)x_2 + c_2(t))^2 \cos^2 x_1 x_4' = \text{constant}$  (say  $= \gamma$ ),

if  $\gamma = 0$ , we get  $x_4' = 0$ , which implies to  $x_4 = 0$ . Then, we obtain the following coordinates

$$x_1 = \frac{iA_1}{1 - i(c_1(t)x_2 + c_2(t))}, x_2 = \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), x_3 = \frac{iA_2}{1 - i\sqrt{\frac{x_2 + c_3(t)}{c_1(t)(c_1(t)x_2 + c_3(t))}}}, x_4 = 0 \quad (9)$$

which is the chaotic hypersurface time-like geodesic  $F_{i1}$  in chaotic flat space  $F_{0123...h}$ , which is the chaotic retraction. Therefore,  $\pi_1(F_{i1} \subset F_{0123...h})$  is isomorphic to identity group.

In a special case if  $x_2 = \frac{-c_2(t)}{c_1(t)}$ , we obtain the chaotic spheres  $S_{i1}^2$  in chaotic flat space  $F_i$ , which represented by

$$x_1 = iA_1, x_2 = iB_1', x_3 = iB_2', x_4 = 0 \quad (10)$$

Then,  $x_1^2 + x_2^2 + x_3^2 - x_4^2 = (A_1^2 + B_1'^2 + B_2'^2) = -k^2$ , which is the geodesic hypersphere in chaotic time-like flat space which is a retraction, where  $A_1, B_1'$  and  $B_2'$  are constants. Therefore  $\pi_1(S_{i1}^2 \subset F_{0123...h})$  is isomorphic to identity group.

Also, if  $c_2(t) = 0$ , then we get the following coordinates.

$$x_1 = iA_1, x_2 = 0, x_3 = \frac{iA_2}{1 - i\sqrt{1/c_1(t)}}, x_4 = 0 \quad (11)$$

Hence,

$$x_1^2 + x_2^2 + x_3^2 - x_4^2 = (iA_1)^2 + \left(\frac{iA_2}{1 - i\sqrt{1/c_1(t)}}\right)^2,$$

which is the chaotic great circle  $S_{i1}^1$  in chaotic flat space-time geodesic. These geodesic is a retraction in chaotic flat space. Therefore,  $\pi_1(S_{i1}^1 \subset F_{0123...h})$  is isomorphic to  $Z_{0123...h}$ . But, if we put  $A_1^* = iA_1, A_2^* = iA_2$  and  $c_1(t) = -c_1^*(t)$ , we have

$$x_1 = A_1^*, x_2 = 0, x_3 = \frac{iA_2^*}{1 - i\sqrt{1/c_1^*(t)}}, x_4 = 0 \quad (12)$$

In this case we obtain a chaotic circle  $S_{i2}^1$ , which is a space-time geodesic. Therefore,  $\pi_1(S_{i2}^1 \subset F_{0123...h})$  is isomorphic to  $Z_{0123...h}$ .

If  $\cos^2 x_1 = 0$ , then  $x_1 = n\frac{\pi}{2}$ ,  $n$  is odd and the chaotic hyper space  $F_{i2}$  is represented by the following coordinate

$$x_1 = n\frac{\pi}{2}, x_2 = \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), x_3 = \frac{iA_2}{1 - i\sqrt{\frac{x_2 + c_3(t)}{c_1(t)(c_1(t)x_2 + c_3(t))}}}, x_4 = A_3 \quad (13)$$

which is a geodesic in time-like chaotic flat space. Also, these geodesics is a retraction on  $F_{0123...h}$ . Therefore,  $\pi_1(F_{i2} \subset F_{0123...h})$  is isomorphic to identity group.

From (6), we obtain

$$\frac{x_2 + c_3(t)}{c_1^2(t)x_2 + c_1(t)c_3(t)} x_3' = \text{constant (say } \alpha).$$

If  $\alpha = 0$ , we get  $x_3' = 0 \Rightarrow x_3 = 0$ , we obtain the following coordinates

$$x_1 = \frac{iA_1}{1 - i(c_1(t)x_2 + c_2(t))}, x_2 = \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), x_3 = 0, x_4 = \frac{A_3}{1 - (c_1(t)x_2 + c_2(t))\cos x_1} \quad (14)$$

which is the chaotic hyperspace  $F_{i3}$ , which is the chaotic retraction and chaotic geodesic in time-like. Therefore  $\pi_1(F_{i3} \subset F_{0123...h})$  is isomorphic to identity group.

Andif

$$\frac{x_2 + c_3(t)}{c_1^2(t)x_2 + c_1(t)c_3(t)} = 0, \text{ then } x_2 = -c_3(t)$$

which deduce also a chaotic time-like geodesic hyperspace  $F_{i4}$  having the following coordinates

$$x_1 = \frac{iA_1}{1 - i(c_1(t)x_2 + c_2(t))}, x_2 = -c_3(t), x_3 = \frac{iA_2}{1 - i\sqrt{\frac{x_2 + c_3(t)}{c_1(t)(c_1(t)x_2 + c_3(t))}}}, x_4 = \frac{A_4}{1 - (c_1(t)x_2 + c_2(t))\cos x_1} \quad (15)$$

Therefore,  $\pi_1(F_{i4} \subset F_{0123\dots h})$  is isomorphic to identity group.

**Corollary 1.** The chaotic fundamental groups of chaotic flat space are time-like geodesics.

**Theorem 2.** The chaotic fundamental group of the limit folding of the chaotic flat space into itself, under (16), is the same as the chaotic fundamental group of the retraction of the chaotic flat space into the geodesic  $F_{i1}$ .

**Proof.** Now, we are going to discuss the folding  $\eta_n$  of the chaotic flat space  $F_{0123\dots h}$ . Let  $\eta_n : F_{0123\dots h} \longrightarrow F_{0123\dots h}$ , be given by  $\eta_n(x_1, x_2, x_3, x_4) = (x_1, x_2, x_3, \frac{|x_4|}{n})$  where

$$n = 1, 2, \dots \tag{16}$$

An isometric chain folding of  $F_{0123\dots h}$  into itself may be defined by

$$\eta_1 : \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\} \rightarrow \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\},$$

$$\eta_2 : \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\} \rightarrow \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\}, \dots$$

$$\eta_n : \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\} \rightarrow \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\}$$

Then, we have

$$\lim_{n \rightarrow \infty} \eta_n = \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \frac{ic_1(t)}{2}(\log B_1 - N \log B_2), \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, 0 \right\},$$

which is a chaotic hypersurface  $F_{i1}$  in chaotic flat space  $F_{0123\dots h}$ , which is the chaotic retraction. Therefore,  $\pi_1(F_{i1} \subset F_{0123\dots h})$  is isomorphic to identity group.

**Theorem 3.** The chaotic fundamental group of the limit folding of the chaotic flat space into itself, under condition (17),  $\pi_1(F_{i5} \subset F_{0123\dots h})$ , is isomorphic to identity group.

**Proof.** If we let  $\Pi_m : F_{0123\dots h} \longrightarrow F_{0123\dots h}$  be given by

$$\Pi_m(x_1, x_2, x_3, x_4) = (x_1, \frac{|x_2|}{m}, x_3, \frac{|x_4|}{m}) \tag{17}$$

Then, the isometric chain folding of  $F_{0123\dots h}$  into itself may be defined by

$$\Pi_1 : \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \left| \frac{ic_1(t)}{2}(\log B_1 - N \log B_2) \right|, \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\} \rightarrow \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \left| \frac{ic_1(t)}{2}(\log B_1 - N \log B_2) \right|, \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\},$$

$$\Pi_2 : \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \left| \frac{ic_1(t)}{2}(\log B_1 - N \log B_2) \right|, \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\} \rightarrow \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \left| \frac{ic_1(t)}{2}(\log B_1 - N \log B_2) \right|, \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\}, \dots$$

$$\Pi_m: \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \left| \frac{ic_1(t)(\log B_1 - N \log B_2)}{2} \right| \right. \\ \left. \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\} \rightarrow \\ \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \left| \frac{ic_1(t)(\log B_1 - N \log B_2)}{2} \right| \right. \\ \left. \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\}$$

Then, we get

$$\lim_{m \rightarrow \infty} \Pi_m = \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, 0, \right. \\ \left. \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, 0 \right\}$$

which is a chaotic hypersurface  $F_{i5}$  in chaotic flat space  $F_{0123...h}$ . Therefore,  $\pi_1(F_{i5} \subset F_{0123...h})$  is isomorphic to identity group.

**Theorem 4.** The chaotic fundamental group of the limit folding of chaotic flat space into itself, under (18), is equivalent to the zero-dimensional sphere  $S_i^0$  in chaotic flat space which isomorphic to identity group.

**Proof.** If the folding is defined by  $\gamma_n : F_{0123...h} \longrightarrow F_{0123...h}$  such that

$$\gamma_n(x_1, x_2, x_3, x_4) = \left( \frac{|x_1|}{n}, \frac{|x_2|}{n}, \frac{|x_3|}{n}, \frac{|x_4|}{n} \right) \quad (18)$$

Then, the isometric chain folding of  $F_{0123...h}$  into itself may be defined by

$$\gamma_1: \left\{ \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))}, \left| \frac{ic_1(t)(\log B_1 - M \log B_2)}{2} \right| \right. \\ \left. \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\}$$

$$\rightarrow \left\{ \left| \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))} \right|, \left| \frac{ic_1(t)(\log B_1 - M \log B_2)}{2} \right| \right. \\ \left. \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\}, \\ \gamma_2: \left\{ \left| \frac{iA_1}{1-i(c_1(t)x_2+c_2(t))} \right|, \left| \frac{ic_1(t)(\log B_1 - M \log B_2)}{2} \right| \right. \\ \left. \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\} \\ \rightarrow \left\{ \frac{iA_1}{2}, \left| \frac{ic_1(t)(\log B_1 - N \log B_2)}{2} \right| \right. \\ \left. \frac{iA_2}{2}, \left| \frac{A_3}{2} \right| \right\},$$

$$\left\{ \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\}, \dots,$$

$$\gamma_n: \left\{ \frac{iA_1}{n-1}, \left| \frac{ic_1(t)(\log B_1 - N \log B_2)}{2} \right| \right. \\ \left. \frac{iA_2}{n-1}, \left| \frac{A_3}{n-1} \right| \right\}$$

$$\left\{ \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\}$$

$$\rightarrow \left\{ \frac{iA_1}{n}, \left| \frac{ic_1(t)(\log B_1 - M \log B_2)}{2} \right| \right. \\ \left. \frac{iA_2}{n}, \left| \frac{A_3}{n} \right| \right\},$$

$$\left\{ \frac{iA_2}{1-i\sqrt{\frac{x_2+c_3(t)}{c_1(t)(c_1(t)x_2+c_3(t))}}}, \left| \frac{A_3}{1-(c_1(t)x_2+c_2(t))\cos x_1} \right| \right\}$$

Then, we get  $\lim_{n \rightarrow \infty} \gamma_n = \{0,0,0,0\}$ , which a zero-dimensional

hypersphere  $S_i^0$  in chaotic flat space. Therefore  $\pi_1(S_i^0 \subset F_{0123...h})$  is isomorphic to identity group.

**Theorem 5.** The end of the limits of the foldings of chaotic flat space of dimension  $n$  is a 0-dimensional chaotic flat space and  $\pi_1(F_i^0 \subset F_{0123...h})$  is isomorphic to identity group.

**Proof:** If we let  $\eta_1 : F_i^n \longrightarrow F_i^n$ ,  $\eta_2 : \eta_1(F_i^n) \longrightarrow \eta_1(F_i^n)$ ,

A.  $\eta_3 : \eta_2(\eta_1(F_i^n)) \longrightarrow \eta_1(F_i^n), \dots$

$\eta_n : \eta_{n-1}(\eta_{n-2}(\eta_1(F_i^n))) \longrightarrow \eta_{n-1}(\eta_{n-2} \cdots (\eta_1(F_i^n)), \dots),$   
 then

$\lim_{n \rightarrow \infty} \eta_n : \eta_{n-1}(\eta_{n-2}(\cdots(\eta_1(F_i^n))\cdots)) = F_i^{n-1}$ , which is the chaotic flat space of dimension  $n - 1$ .

Also, if we consider

$\gamma_1 : F_i^{n-1} \longrightarrow F_i^{n-1}$ ,  $\gamma_2 : \gamma_1(F_i^{n-1}) \longrightarrow \gamma_1(F_i^{n-1})$ ,  
 $\gamma_3 : \gamma_2(\gamma_1(F_i^{n-1})) \longrightarrow \gamma_1((F_i^{n-1}))$ ,  
 $\gamma_m : \gamma_{m-1}(\gamma_{m-2}(\gamma_1(F_i^{n-1}))) \longrightarrow \gamma_{m-1}(\gamma_{m-2} \cdots (\gamma_1(F_i^{n-1})), \dots)$ , then  
 $\lim_{m \rightarrow \infty} \gamma_m : \gamma_{m-1}(\gamma_{m-2}(\cdots(\eta_1(F_i^{n-1})))\cdots) = F_i^{n-2}$ , which is the chaotic flat space of dimension  $n-2$ .

Consequently,  $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \gamma_m(\eta_n(\cdots(\eta_1(F_i^n)))) = F_i^0$ ,

which is a zero-dimensional chaotic space. Therefore  $\pi_1(F_i^0 \subset F_{0123\dots h})$  is isomorphic to identity group.

**Theorem 6.** Under (18), the chaotic fundamental group of the limit of foldings of chaotic flat space into itself coincide with minimal retraction.

**Proof.** Let  $\eta_i : F_{0123\dots h}^n \longrightarrow F_{0123\dots h}^n$  be a type of foldings and  $r_i$  are the retractions. Then, we have the following chains

$$\begin{aligned}
 &F_i^n \xrightarrow{\eta_1} F_{i1}^n \xrightarrow{\eta_2} F_{i2}^n \longrightarrow \cdots F_{i(n-1)}^n \xrightarrow{\text{Lim } \eta_i^n} F_i^{n-1} \\
 &F_i^n \xrightarrow{r_1} F_{i1}^n \xrightarrow{r_2} F_{i2}^n \longrightarrow \cdots F_{i(n-1)}^n \xrightarrow{\text{Lim } r_i^n} F_i^{n-1}, \\
 &F_i^{n-1} \xrightarrow{\eta_1^2} F_{i1}^{n-1} \xrightarrow{\eta_2^2} F_{i2}^{n-1} \longrightarrow \cdots F_{i(n-1)}^{n-1} \xrightarrow{\text{Lim } \eta_i^{n-1}} F_i^{n-2}, \\
 &F_i^{n-1} \xrightarrow{r_1^2} F_{i1}^{n-1} \xrightarrow{r_2^2} F_{i2}^{n-1} \longrightarrow \cdots F_{i(n-1)}^{n-1} \xrightarrow{\text{Lim } r_i^{n-1}} F_i^{n-2} \\
 &\vdots \\
 &F_i^1 \xrightarrow{\eta_1^n} F_{i1}^1 \xrightarrow{\eta_2^n} F_{i2}^1 \longrightarrow \cdots F_{i(n-1)}^1 \xrightarrow{\text{Lim } \eta_i^n} F_i^0 \\
 &F_i^1 \xrightarrow{r_1^n} F_{i1}^1 \xrightarrow{r_2^n} F_{i2}^1 \longrightarrow \cdots F_{i(n-1)}^1 \xrightarrow{\text{Lim } r_i^n} F_i^0.
 \end{aligned}$$

Thus from the above chain the end of the limits of folding coincides with the zero-dimensional space which is the limit of retractions and therefore  $\pi_1(F_i^0 \subset F_{0123\dots h})$  is isomorphic to identity group.

### III. CONCLUSION

In this paper we achieved the approval of the important of the chaotic fundamental group of the chaotic geodesic flat space. The relation between the folding of chaotic flat space and chaotic fundamental group are discussed. Theorems which govern these relations are presented.

### REFERENCES

- [1] A.E. El-Ahmady, The variation of the density on chaotic spheres in chaotic space-like Minkowski space time, Chaos, Solitons and Fractals, Vol. 31, (1272-1278), (2007).
- [2] A.E. El-Ahmady, Folding of fuzzy hypertori and their retractions, Proc. Math. Phys. Soc. Egypt, Vol.85, No.1,(1-10), (2007).
- [3] A.E. El-Ahmady, Limits of fuzzy retractions of fuzzy hyperspheres and their foldings, Tamkang Journal of Mathematics, Vol.37, No. 1, (47-55), (2006).
- [4] A.E. El-Ahmady, Fuzzy folding of fuzzy horocycle, Circolo Matematico di Palermo Serie II, Tomo L. III, (443-450), (2004).
- [5] A.E. El-Ahmady, Fuzzy Lobachevskian space and its folding, The Journal of Fuzzy Mathematics, Vol. 12, No. 2, (609-614), (2004).
- [6] A.E. El-Ahmady, The deformation retract and topological folding of Buchdahi space, Periodica Mathematica Hungarica Vol. 28, (19-30), (1994).
- [7] A. E. El-Ahmady, The geodesic deformation retract of Klein bottle and its folding, The International Journal of Nonlinear Science, Vol. 9, No. 3, (1-8), (2011).
- [8] A.E. El-Ahmady, Folding and fundamental groups of Buchdahi space, Indian Journal of Science and Technology, Vol.6, No. 1, (3940-3945), (2013).
- [9] A.E. El-Ahmady, Folding of some types of Einstein spaces, The International Journal of Nonlinear Science, (In press).
- [10] A.E. El-Ahmady, Onelastic Klein bottle and fundamental groups, Applied Mathematics, Vol.4, No.3, (499-504), (2013).
- [11] A.E. El-Ahmady, Retraction of chaotic black hole, The Journal of Fuzzy Mathematics, Vol.19, No.4, (833-838), (2011).
- [12] A.E. El-Ahmady, On the fundamental group and folding of Klein bottle, International Journal of Applied Mathematics and Statistics, Vol.37, No. 6, (56-64), (2013).
- [13] A.E. El-Ahmady, Fuzzy elastic Klein bottle and its retractions, International Journal of Applied Mathematics and Statistics, Vol.42, No. 12, (94-102), (2013).
- [14] A.E. El-Ahmady, Folding and fundamental groups of flat Robertson-Walker Space, Indian Journal of Science and Technology, Vol.6, No. 4, (4235-4242), (2013).
- [15] A.E. El-Ahmady, Fundamental groups and folding of Minkowski space, European Journal of Scientific Research, Vol.104, No. 2, (284-293), (2013).
- [16] A.E. El-Ahmady, Retraction of chaotic Schwarzschild space, European Journal of Scientific Research, Vol.104, No. 3, (333-339), (2013).
- [17] A.E. El-Ahmady, The isonormal folding of fuzzy manifolds and its retractions, European Journal of Scientific Research, Vol.104, No. 3, (340-347), (2013).
- [18] A.E. El-Ahmady and A.Al-Rdade, Deformation retracts of the Reissner-Nordström spacetime and its foldings, American Journal of Applied Sciences, Vol.10, No. 7, (740-745), (2013).
- [19] A.E. El-Ahmady and A.Al-Rdade, On the geometry of fuzzy Reissner-Nordström spacetime and its foldings, European Journal of Scientific Research, Vol.104, No. 2, (294-303), (2013).
- [20] A. E. El-Ahmady, Folding and fundamental groups of Buchdahi space, Indian Journal of Science and Technology, Vol.6, No. 1, (3940-3945), (2013).
- [21] A.E. El-Ahmady and N.Al-Hazmi, On the folding of Buchdahi space, European Journal of Scientific Research, Vol. 100, No. 2, (315-322), (2013).
- [22] A.E. El-Ahmady and N.Al-Hazmi, The exponential deformation retract of the hypertori and its foldings, European Journal of Scientific Research, Vol.104, No. 4, (561-568), (2013).
- [23] A.E. El-Ahmady and K.Al-Onema, The deformation retractions of fuzzy sphere in fuzzy Lobachevsky space and its folding, European Journal of Scientific Research, Vol.104, No. 4, (549-560), (2013).

- [24] A. E. El-Ahmady, Folding and unfolding of chaotic spheres in chaotic space-like Minkowski space-time, *The Scientific Journal of Applied Research*, Vol. 1(2), (34-43), (2012).
- [25] A. E. El-Ahmady and H. Rafat, A calculation of geodesics in chaotic flat space and its folding, *Chaos, Solutions and Fractals*, 30 (836-844) (2006).
- [26] M. El-Ghoul, A. E. El-Ahmady: The deformation retract and topological folding of the flat space, *Jour. Inst. Math. & Comp. Sci. (Math. Ser.)* Vol. 5, No. 3, (1992), 349-356.
- [27] M. Abu-Saleem, Homology group on the dynamical trefoil knot, *Indian Journal of Science and Technology*, Vol 6 (5) (2013) : 4514-4518.
- [28] M. Abu-Saleem, On Chaotic Cartesian Product of Graphs and their Retractions, *International Journal of Nonlinear Science*, UK, Vol.15, No.1(2013): 86-90.
- [29] M. Abu-Saleem, On chaotic homotopy group, *Advanced Studies in Contemporary Mathematics*, Vol.23, No.1 (2013): 69-75.
- [30] M. Abu-Saleem, On the dynamical hyperbolic 3-spaces and their deformation retracts. *Proceedings of the Jangjeon Mathematical society*, 15 No. 2 (2012): 189-193.
- [31] M. Abu-Saleem, Dynamical chaotic homotopy group and its applications, *International Journal of Nonlinear Science*, Vol.11, No.2, (206-212), (2011).
- [32] M. Abu-Saleem, On dynamical chaotic de sitter spaces and their deformation retracts, *Proceedings of the Jangjeon Mathematical society*, Vol.14, (231-238), (2011).
- [33] M. Abu-Saleem, On dynamical chaotic Weyl representations of the vacuum C metric and their retractions, *International journal of Mathematical Combinatorics*, Vol.3, (47-54), (2011).
- [34] M. Abu-Saleem, Folding on the wedge sum of graphs and their fundamental group, *Applied Sciences*, Vol.12, (14-19), (2010).
- [35] M. Abu-Saleem, Homology group on manifolds and their foldings, *Tamkang journal of mathematics*, Vol.41, No.1, (31-38), (2010).
- [36] M. Abu-Saleem, Folding and unfolding of manifolds and their fundamental groups, *Int.J.Contemp.Math.Sciences*, Vol.5, No.1, (1-19), (2010).
- [37] M. Abu-Saleem, Dynamical Knot and their fundamental group, *International journal of Mathematical Combinatorics*, Vol.1, (80-86), (2010).
- [38] M. Abu-Saleem, Dynamical manifolds and their fundamental group, *Advanced Studies in Contemporary Mathematics*, Vol.20, No.1, (125-131), (2010).
- [39] M. Abu-Saleem, Conditional fractional folding of a manifold and their fundamental group, *Advanced Studies in Contemporary Mathematics*, Vol.20, No.2, (271-277), (2010).
- [40] M. Abu-Saleem, Folding on the chaotic Cartesian product of manifolds and their fundamental group, *Tamkang journal of mathematics*, Vol.39, No.4, (353-361), (2008).
- [41] S. Lefschetz, *Algebraic Topology*, American mathematical society, (1942).
- [42] G. L. Naber, *Topology, Geometry and Gauge fields*, Foundations, Springer-Verlag New York, Berlin, (2011).
- [43] M. Arkowitz, *Introduction to homotopy theory*, Springer-Verlag, (2011).
- [44] J. Strom, *Modern classical homotopy theory*, American Mathematical Society, (2011).