

Some Equalities Connected with Fuzzy Soft Matrices

D. R. Jain

Abstract—The aim of this paper is to use matrix representation of Fuzzy soft sets for proving some equalities connected with Fuzzy soft sets based on set-operations.

Keywords—Equality, Fuzzy soft matrix, Fuzzy soft sets, operations.

I. INTRODUCTION

WE cannot always use the classical methods to solve complicated problems in economics, engineering, social sciences, medical sciences etc. because different types of uncertainties are present in these theories. These days so many theories are available to deal with such type of uncertainties, such as, theory of fuzzy sets [1], theory of intuitionistic fuzzy sets [2], [3], theory of vague sets [4], theory of rough sets [5]. Every theory has its own problems because of the inadequacy of the parameterization tool of the theories. In 1999, Molodtsov [6] initiated the concept of fuzzy set to deal with uncertainties which is free from the above difficulties and later in 2001 Maji et al. [7] gave a hybrid model known as fuzzy soft set which is a combination of soft set and fuzzy set. They introduce the concept of union, intersection, complement etc. of Fuzzy Soft Sets and discussed their properties. These results were further revised and improved by Ahmad and Kharal [8]. Manash Jyoti Borah et al. [9] introduced the notions of symmetric, transitive, reflexive, equivalence fuzzy soft relations and discussed some related properties.

The theory of matrix is commonly used in the broad areas of science and engineering. However classical matrix theory sometimes fails to solve the problems involving uncertainties, occurring in an imprecise environment. In [10], Yong Yang and Chenli Ji, using a matrix representation [10], [11] of Fuzzy Soft Set, discussed its properties and applied the same in certain decision making problems. In [12] the notion of Fuzzy Soft Matrices has been extended.

In this paper, by defining some more set-operations on FSM_{mxn} analogous to [13], [14] defined for intuitionistic fuzzy sets, an attempt has been made to derive some new results based on these operations on FSM_{mxn} .

II. PRELIMINARIES

In this section, we give definitions and notions (refer [12]), to be used in our subsequent work.

Definition 1: *Fuzzy Soft Set*- Let X be an initial universal set and E be a set of parameters. Let $\tilde{P}(X)$ denotes the power set of all Fuzzy subsets of X . Let $A \subseteq E$. A pair (F, A) is called

Fuzzy Soft Set over X , where F is a mapping given by $F : A \rightarrow \tilde{P}(X)$.

Definition 2: *Fuzzy Soft Class* - The pair (X, E) denotes the collection of all Fuzzy Soft Sets on X with attributes from E and is called Fuzzy Soft Class.

Definition 3: *Fuzzy Soft Matrices* - Let $X = \{x_1, x_2, \dots, x_m\}$ be the universal set and $E = \{e_1, e_2, \dots, e_n\}$ be the set of parameters. Let $A \subseteq E$ and (F, A) be a Fuzzy Soft Set in the Fuzzy Soft Class (X, E) . Then we represent the Fuzzy Soft Set (F, A) in the matrix form as:

$$A_{mn} = [a_{ij}]_{mxn} \text{ or simply by } A = [a_{ij}]$$

where

$$a_{ij} = \begin{cases} \mu_j(x_i) & \text{if } e_j \in A \\ 0 & \text{if } e_j \notin A \end{cases}$$

Here $\mu_j(x_i)$ represent the membership of x_i in the Fuzzy Set $F(e_j)$. We would identify a Fuzzy Soft Set with its Fuzzy Soft matrix and vice versa. The set of all mxn Fuzzy Soft Matrices will be denoted FSM_{mxn} over X .

Definition 4: *Set operations on FSM_{mxn}* - Let $A = [a_{ij}]_{mxn}$ and $B = [b_{ij}]_{mxn}$ be two Fuzzy Soft matrices over the universal set X .

Some set operations on FSM_{mxn} are defined as follows:

$$A \cup B = C = [c_{ij}]_{mxn} \quad (1)$$

where

$$c_{ij} = \max(a_{ij}, b_{ij}), \text{ for all } i \text{ and } j.$$

$$A \cap B = C = [c_{ij}]_{mxn} \quad (2)$$

where

$$c_{ij} = \min(a_{ij}, b_{ij}), \text{ for all } i \text{ and } j.$$

$$A \diamond B = C = [c_{ij}]_{mxn} \quad (3)$$

where

$$c_{ij} = a_{ij} + b_{ij} - a_{ij} b_{ij}, \text{ for all } i \text{ and } j.$$

$$A \cdot B = C = [c_{ij}]_{mxn} \quad (4)$$

where

$$c_{ij} = a_{ij} \cdot b_{ij} \text{ for all } i \text{ and } j.$$

$$A @ B = C = [c_{ij}]_{m \times n} \quad (5)$$

where

$$c_{ij} = \frac{1}{2} (a_{ij} + b_{ij}), \text{ for all } i \text{ and } j.$$

$$A \$ B = C = [c_{ij}] \quad (6)$$

where

$$c_{ij} = \sqrt{a_{ij} \cdot b_{ij}}, \text{ for all } i \text{ and } j.$$

$$A \# B = C = [c_{ij}] \quad (7)$$

where

$$c_{ij} = \frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}}, \text{ for all } i \text{ and } j,$$

for which we will accept that if $a_{ij} = b_{ij} = 0$, then

$$\frac{a_{ij} b_{ij}}{a_{ij} + b_{ij}} = 0.$$

$$A * B = \left[\frac{a_{ij} + b_{ij}}{2(a_{ij} b_{ij} + 1)} \right] \quad (8)$$

III. EQUALITIES CONNECTED WITH $FSM_{M \times N}$

Theorem 1: If $A = [a_{ij}]$, $B = [b_{ij}] \in FSM_{m \times n}$, then

$$\{(A \cup B) \# (A \cap B)\} \$ \{(A \cup B) @ (A \cap B)\} = A \$ B$$

$$\begin{aligned} \text{Proof: } (A \cup B) \# (A \cap B) &= [\max(a_{ij}, b_{ij})] \# [\min(a_{ij}, b_{ij})] \\ &= \left[\frac{2 \max(a_{ij}, b_{ij}) \cdot \min(a_{ij}, b_{ij})}{\max(a_{ij}, b_{ij}) + \min(a_{ij}, b_{ij})} \right] = \left[\frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right] \end{aligned}$$

and

$$\begin{aligned} (A \cup B) @ (A \cap B) &= [\max(a_{ij}, b_{ij})] @ [\min(a_{ij}, b_{ij})] \\ &= \left[\frac{\max(a_{ij}, b_{ij}) + \min(a_{ij}, b_{ij})}{2} \right] = \left[\frac{a_{ij} + b_{ij}}{2} \right]. \end{aligned}$$

Thus

$$\begin{aligned} \{(A \cup B) \# (A \cap B)\} \$ \{(A \cup B) @ (A \cap B)\} \\ &= \left[\frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right] \$ \left[\frac{a_{ij} + b_{ij}}{2} \right] = \left[\frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \cdot \frac{a_{ij} + b_{ij}}{2} \right] \\ &= [\sqrt{a_{ij} b_{ij}}] = A \$ B \end{aligned}$$

Hence the result is proved.

Theorem 2: If $A = [a_{ij}]$, $B = [b_{ij}] \in FSM_{m \times n}$, then

$$\langle \{(A \cup B) \cdot (A \cap B)\} @ (A \diamond B) \rangle \$ (A \# B) = (A \$ B)$$

$$\begin{aligned} \text{Proof: } (A \cup B) \cdot (A \cap B) &= [\max(a_{ij}, b_{ij})] \cdot [\min(a_{ij}, b_{ij})] \\ &= [\max(a_{ij}, b_{ij}) \cdot \min(a_{ij}, b_{ij})] = [a_{ij} b_{ij}] \end{aligned}$$

and

$$\begin{aligned} \{(A \cup B) \cdot (A \cap B)\} @ (A \diamond B) \\ &= [a_{ij} b_{ij}] @ [a_{ij} + b_{ij} - a_{ij} b_{ij}] = \left[\frac{a_{ij} b_{ij} + a_{ij} + b_{ij} - a_{ij} b_{ij}}{2} \right] \\ &= \left[\frac{a_{ij} + b_{ij}}{2} \right] \end{aligned}$$

Now consider

$$\begin{aligned} \langle \{(A \cup B) \cdot (A \cap B)\} @ (A \diamond B) \rangle \$ (A \# B) \\ &= \left[\frac{a_{ij} + b_{ij}}{2} \right] \$ \left[\frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right] = \left[\sqrt{\frac{a_{ij} + b_{ij}}{2} \cdot \frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}}} \right] \\ &= [\sqrt{a_{ij} b_{ij}}] = A \$ B \end{aligned}$$

Hence the result is proved.

Corollary 1: If $A = [a_{ij}]$, $B = [b_{ij}] \in FSM_{m \times n}$, then

$$\begin{aligned} \{(A \cup B) \# (A \cap B)\} \$ \{(A \cup B) @ (A \cap B)\} \\ &= \langle \{(A \cup B) \cdot (A \cap B)\} @ (A \diamond B) \rangle \$ (A \# B) = (A \$ B). \end{aligned}$$

Proof: It follows from Theorems 1 & 2.

Theorem 3: If $A = [a_{ij}]$, $B = [b_{ij}] \in FSM_{m \times n}$, then

$$(A \diamond B) \cup (A @ B) @ (A \cdot B) \cap (A @ B) = (A @ B)$$

Proof:

$$\begin{aligned} (A \diamond B) \cup (A @ B) &= [a_{ij} + b_{ij} - a_{ij} b_{ij}] \cup \left[\frac{a_{ij} + b_{ij}}{2} \right] \\ &= \left[\max\left(a_{ij} + b_{ij} - a_{ij} b_{ij}, \frac{a_{ij} + b_{ij}}{2}\right) \right] = [a_{ij} + b_{ij} - a_{ij} b_{ij}] \end{aligned}$$

and

$$\begin{aligned} (A \cdot B) \cap (A @ B) &= [a_{ij} b_{ij}] \cap \left[\frac{a_{ij} + b_{ij}}{2} \right] \\ &= \left[\min\left(a_{ij} b_{ij}, \frac{a_{ij} + b_{ij}}{2}\right) \right] = [a_{ij} b_{ij}] \end{aligned}$$

Now Consider

$$\begin{aligned} \{(A \diamond B) \cup (A @ B)\} @ \{(A \cdot B) \cap (A \# B)\} \\ = [a_{ij} + b_{ij} - a_{ij} b_{ij}] @ [a_{ij} b_{ij}] &= \left[\frac{a_{ij} + b_{ij} - a_{ij} b_{ij} + a_{ij} b_{ij}}{2} \right] \\ &= \left[\frac{a_{ij} + b_{ij}}{2} \right] = (A @ B), \end{aligned}$$

which proves the result

Theorem 4: If $A = [a_{ij}]$, $B = [b_{ij}] \in \text{FSM}_{m \times n}$, then

$$\{(A \diamond B) \cap (A @ B)\} \$ \{(A \diamond B) \cap (A \# B)\} = (A \$ B).$$

Proof: $(A \diamond B) \cap (A @ B) = [a_{ij} + b_{ij} - a_{ij} b_{ij}] \cap \left[\frac{a_{ij} + b_{ij}}{2} \right]$

$$= \left[\min \left(a_{ij} + b_{ij} - a_{ij} b_{ij}, \frac{a_{ij} + b_{ij}}{2} \right) \right] = \left[\frac{a_{ij} + b_{ij}}{2} \right]$$

and

$$\begin{aligned} (A \diamond B) \cap (A \# B) &= \left[\min \left(a_{ij} + b_{ij} - a_{ij} b_{ij}, \frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right) \right] \\ &= \left[\frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right] \end{aligned}$$

Thus

$$\begin{aligned} \{(A \diamond B) \cap (A @ B)\} \$ \{(A \diamond B) \cap (A \# B)\} \\ = \left[\frac{a_{ij} + b_{ij}}{2} \right] \$ \left[\frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right] = \sqrt{\frac{a_{ij} + b_{ij}}{2} \cdot \frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}}} = \left[\sqrt{a_{ij} b_{ij}} \right] \\ = A \$ B \end{aligned}$$

Hence the result is proved.

Theorem 5: If $A = [a_{ij}]$, $B = [b_{ij}] \in \text{FSM}_{m \times n}$, then

$$\{(A \diamond B) \cup (A \# B)\} @ \{(A \cdot B) \cap (A \# B)\} = (A @ B).$$

Proof:

$$\begin{aligned} (A \diamond B) \cup (A \# B) &= [a_{ij} + b_{ij} - a_{ij} b_{ij}] \cup \left[\frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right] \\ &= \left[\max \left(a_{ij} + b_{ij} - a_{ij} b_{ij}, \frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right) \right] = [a_{ij} + b_{ij} - a_{ij} b_{ij}] \end{aligned}$$

and

$$\begin{aligned} (A \cdot B) \cap (A \# B) &= [a_{ij} b_{ij}] \cap \left[\frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right] = \left[\min \left(a_{ij} b_{ij}, \frac{2a_{ij} b_{ij}}{a_{ij} + b_{ij}} \right) \right] \\ &= [a_{ij} b_{ij}] \end{aligned}$$

Now

$$\begin{aligned} \{(A \diamond B) \cup (A \# B)\} @ \{(A \cdot B) \cap (A \# B)\} \\ = [a_{ij} + b_{ij} - a_{ij} b_{ij}] @ [a_{ij} b_{ij}] &= \left[\frac{a_{ij} + b_{ij} - a_{ij} b_{ij} + a_{ij} b_{ij}}{2} \right] \\ &= \left[\frac{a_{ij} + b_{ij}}{2} \right] = A @ B \end{aligned}$$

Thus the result is proved.

Theorem 6: If $A = [a_{ij}]$, $B = [b_{ij}] \in \text{FSM}_{m \times n}$, then

$$\{(A \diamond B) \cup (A \$ B)\} @ \{(A \cdot B) \cap (A \$ B)\} = (A @ B)$$

Proof:

$$\begin{aligned} (A \diamond B) \cup (A \$ B) &= [a_{ij} + b_{ij} - a_{ij} b_{ij}] \cup \left[\sqrt{a_{ij} b_{ij}} \right] \\ &= \left[\max \left(a_{ij} + b_{ij} - a_{ij} b_{ij}, \sqrt{a_{ij} b_{ij}} \right) \right] \\ &= [a_{ij} + b_{ij} - a_{ij} b_{ij}] \end{aligned}$$

and

$$\begin{aligned} (A \cdot B) \cap (A \$ B) &= [a_{ij} b_{ij}] \cap \left[\sqrt{a_{ij} b_{ij}} \right] = \left[\min \left(a_{ij} b_{ij}, \sqrt{a_{ij} b_{ij}} \right) \right] \\ &= [a_{ij} b_{ij}] \end{aligned}$$

Now

$$\begin{aligned} \{(A \diamond B) \cup (A \$ B)\} @ \{(A \cdot B) \cap (A \$ B)\} \\ = [a_{ij} + b_{ij} - a_{ij} b_{ij}] @ [a_{ij} b_{ij}] &= \left[\frac{a_{ij} + b_{ij} - a_{ij} b_{ij} + a_{ij} b_{ij}}{2} \right] \\ &= \left[\frac{a_{ij} + b_{ij}}{2} \right] = (A @ B) \end{aligned}$$

Thus the result is proved.

Theorem 7: If $A = [a_{ij}]$, $B = [b_{ij}] \in \text{FSM}_{m \times n}$, then

$$(A \cup B) * (A \cap B) = A * B.$$

Proof:

$$\begin{aligned} (A \cup B) * (A \cap B) &= \left[\max(a_{ij}, b_{ij}) \right] * \left[\min(a_{ij}, b_{ij}) \right] \\ &= \left[\frac{\max(a_{ij}, b_{ij}) + \min(a_{ij}, b_{ij})}{2 \{ \max(a_{ij}, b_{ij}) \cdot \min(a_{ij}, b_{ij}) + 1 \}} \right] \\ &= \left[\frac{a_{ij} + b_{ij}}{2(a_{ij} b_{ij} + 1)} \right] = A * B \end{aligned}$$

Hence the result is proved.

IV. CONCLUSION

Manashm, Jyoti Borah et al. [12] defined some operations on FSM such as union intersection complement etc. and discussed their properties. In the present paper this study has

further been extended by defining some more operations on it. Further work in this direction is required to study whether the notion put forward in this paper yields a fruitful result.

ACKNOWLEDGMENT

The author wishes to express his sincere thanks to the editor and anonymous referee for their valuable suggestions which helped in improving the presentation of the paper.

REFERENCES

- [1] L.A. Zadeh, Fuzzy Sets, Information and Control, 8, 1965, 338-353
- [2] K.T. Atanassov, Intuitionistic fuzzy sets, Fuzzy sets and systems 20(1) 1986, 87-96
- [3] K.T. Atanassov, Intuitionistic fuzzy sets, Springer Physica-Verlag, Heidelberg, 1999
- [4] W.L. Gau and Buehrer, Vague sets IEEE Trans, System Man Cybernet 23(2), 1993, 610-614
- [5] Z. Pawlak, Rough sets, International journal of information and computer sciences 11, 1982, 341-356
- [6] D. Molodtsov, Soft Set Theory-first Results, Computers Math with Applications, 37, 1999, 19-31
- [7] P.K. Maji, R. Biswas and A.R. Roy, Fuzzy Soft Sets, J. Fuzzy Math, 9(3), 2001, 589-602.
- [8] B. Ahmed and A. Kharal, On Fuzzy Soft Sets, advances in Fuzzy System Volume 2009.
- [9] Manash Jyoti Borah, Tridiv Jyoti Neog, Dusmanta Kumar Sut, Relations on fuzzy soft sets, J. Math. Comput. Sci. 2, 2012, no. 3, 515-634.
- [10] Yong Yang and Chen Li Ji, Fuzzy Soft Matrices and their Applications, AICI 2011, Part I INAI 7002, pp. 618-627, 2011.
- [11] B. Chetia and P.K. Das, Journal of the Assam Academy of Mathematics, 2010, 2, 71-83
- [12] Manash Jyoti Borah, Tridiv Jyoti Neog, Dusmanta Kumar Sut, Fuzzy Soft Matrix Theory and its Decision Making, International Journal of Modern Engineering Research (IJMER) Vol. 2 Issue 2, 2012, 121-127.
- [13] Rajkumar Verma, Bhu Dev Sharma, some new equalities connected with Intuitionistic fuzzy sets, Notes on Intuitionistic fuzzy sets Vol. 19, 2, 2013, 25-30
- [14] K.T. Atanassov, New Operation defined over intuitionistic fuzzy sets, Fuzzy sets and systems, 61, 1994, 137-142

D.R. Jain, currently Professor of Mathematics at Jaypee Institute of Information Technology, Noida, UP India, did his PhD from University of Delhi in 1976. He served ARSD College University of Delhi for 44 years in different positions. He has published more than 18 research papers in journals of international repute.