

Amplitude and Latency of P300 Component from Auditory Stimulus in Different Types of Personality: An Event Related Potential Study

Nasir Yusoff, Ahmad Adamu Adamu, Tahamina Begum, Faruque Reza

Abstract—The P300 from Event related potential (ERP) explains the psycho-physiological phenomenon in human body. The present study aims to identify the differences of amplitude and latency of P300 component from auditory stimuli, between ambiversion and extraversion types of personality. Ambivert (N=20) and extravert (N=20) undergoing ERP recording at the Hospital Universiti Sains Malaysia (HUSM) laboratory. Electroencephalogram data was recorded with oddball paradigm, counting auditory standard and target tones, from nine electrode sites (Fz, Cz, Pz, T3, T4, T5, T6, P3 and P4) by using the 128 HydroCel Geodesic Sensor Net. The P300 latency of the target tones at all electrodes were insignificant. Similarly, the P300 latency of the standard tones were also insignificant except at Fz and T3 electrode. Likewise, the P300 amplitude of the target and standard tone in all electrode sites were insignificant. Extravert and ambivert indicate similar characteristic in cognition processing from auditory task.

Keywords—Amplitude, Event Related Potential, P300 Component, Latency

I. INTRODUCTION

PERSONALITY tells many things on characteristics of people and provides benefits in many ways especially for the therapist or counselor to get to know their patient better and planning for intervention. In addition, personality plays a role in determining a person's action or emotion when faced with everyday life event whether pleasant or stressful.

In [1], personality is said to be made up of three important parts: traits, characteristic adaptations and life stories. Meanwhile, it is also suggested that personality is developed through four main sectors – cognitive, conative, affective and somatic [2]. One quote even mentioned [3]:

“There is a growing consensus about the validity of human personality traits as important dispositions toward feelings and behaviors.”

A type of one's personality has been correlated to many other factors such as academics, health problems, employment, social life and even cognitive performance [4]-[6]. Formerly, personality is categorized into three parts which

are emotional stability/instability (later known as neuroticism), extraversion/introversion and psychoticism (aggressiveness/antisocial) [7]. Then, [8] proposed a new theory for personalities which are now commonly applied by researchers called the Big Five personalities trait. These are Openness to experience, Conscientiousness, Extraversion, Agreeableness and Neuroticism. These five personalities are discovered when researchers founded five recurrent factors in analyzing personality ratings in eight different samples from their study [9].

Contradict to introvert, extraversion refers to the tendency toward interpersonal interaction, activism, the need for happiness, and the capacity for joy [10]. Ambiverts are categorised as people who are neither introverts nor extraverts, but are in the middle between the two extremes [11]. It was shown that the risk of cognitive impairments was lower in the ambivert group compared to introvert group or those with high extraversion level [12].

Event related potential (ERP) is a method used in neuroscience to explain neural mechanism associated with human emotion and attention, as well as the neural process of decision making [13]. In ERP, the P300 is among an important component that able to describe or explain the psycho-physiological phenomenon in human body, and it is consistently related to attention, decision making and memory updating [14]. Thus, in this study, we hypothesize that there is some difference in cognitive processing between different type of personality (i.e. ambiversion versus extraversion), as indicated by the P300 component of ERP.

II. METHODOLOGY

A. Participants

Forty undergraduate medical students of Universiti Sains Malaysia were recruited as participants. The participants were grouped into two – ambivert (N=20) and extravert (N=20), based on the score obtained by using personality inventory [15]. A score of 17-32 was considered ambiversion and a score of 33 and above was regarded extraversion [15]. This study has been approved by the Human Ethical Committee of Universiti Sains Malaysia [USM/PPSP/2013/JKP-65[65.3(4)]. Written informed consent was signed up upon agreement to participate in the study. Subjects were excluded from the study if they had hearing impairment, neurosurgical operations history, history of stroke and Other major diseases.

Nasir Yusoff is a Lecturer at the Department of Neurosciences, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia (corresponding author, phone and fax:609-7676300; e-mail: nasiryusoff2310@gmail.com).

Ahmad Adamu Adamu (post-graduate student), Tahamina Begum (medical lecturer) and Faruque Reza (medical lecturer) are with the Department of Neurosciences, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan, Malaysia (e-mail: physioadam@gmail.com, tahamina@usm.my, faruque@usm.my).

B. ERP Procedure

This study was conducted at Event Related Potential/ Magnetoencephalography (ERP/MEG) laboratory of Hospital Universiti Sains Malaysia (HUSM), a sound attenuated room in order to minimize noise. The 128 HydroCel Geodesic Sensor Net was applied, positioned symmetrically on the head (Fig. 1), following the guidelines of standard electrode positions by the 10–20 International system of electrode placement. Auditory stimulus was passed binaurally through a conventional audiometric earphone from the Net station. The electroencephalogram (EEG) data was recorded from nine electrode sites (Fz, Cz, Pz, T3, T4, T5, T6, P3 and P4) scalp sites with Ag/AgCl electrodes placed on the sites. Oddball paradigm was applied in which subjects counted target tones

(60 dB sound pressure level with high pitch at 2000 Hz) and at the same time ignored the standard tones (60 dB sound pressure level with low pitch at 1000 Hz). Tone duration was set up at 100ms with rise/fall time of 10ms and amplifiers with band pass of 0.3 to 30 Hz. Pre-analysis (i.e. filtering, segmentation, artifact detection, bad channel replacement, averaging, montage operation and baseline correction) was done after experiment session. Estimated duration time for ERPs recording was approximately 15 minutes. After pre-analysis, the ERP data was transferred to the Statistical Package for the Social Science version 22 for further analysis. Descriptive statistic (mean and standard deviation) and Independent T-test was used to explain the difference between groups. Significance level was set at 0.05.




Fig. 1 128 HydroCel Geodesic Sensor Net

III. RESULT

A. Socio-Demographic Data

Out of forty respondents, ambivert comprised of twenty subjects, however, one respondent was excluded because the recording was too noisy (8 males and 11 females). Meanwhile, the extravert group comprised of twenty subjects (12 males and 8 females). The mean age and standard deviation of ambivert was 22.1±1.15 and extravert was 22.7±1.41. No significant of age difference was seen between the two groups (p=0.20). Mean and standard deviation of personality score in ambivert was 28.95±2.97 and in extravert was 38.45±3.87 (Table I)

TABLE I
SOCIO-DEMOGRAPHIC DATA

	Ambivert (n=19)	Extravert (n=20)
Sex (male/female)	8/11	12/8
Age (mean years±SD)	22.1±1.15	22.7±1.41
Personality (mean score±SD)	28.9±2.97	38.4±3.87
Handedness (right/left)	18/1	16/4
Race (Malay, Chinese, Indian)	8/8/3	9/9/2

B. P300 Latency

The mean latency of the target tone for the ambivert group was between 503.26 milisecond and 539.21 milisecond and for the standard tone was between 463 milisecond and 511 milisecond. For the extravert group, the mean latency for the

target tone was between 497.7 milisecond and 535.8 milisecond, and between 415.6 milisecond and 469.45 milisecond for the standard tone. The difference of P300 latency of target tone between extravert and ambivert was insignificant at all electrode sites - Fz ($p=0.55$), Cz ($p=0.92$), Pz ($p=0.96$), T3 ($p=0.74$), T4($p=0.55$), T5 ($p=0.36$), T6 ($p=0.53$), P3 ($p=0.56$) and P4 ($p=0.55$) (Table II). For the standard tone, the difference of P300 latency between extravert and ambivert was significant at Fz ($p=0.04$) and T3 ($p=0.05$), however indicated insignificant result at Cz ($p=0.14$), Pz ($p=0.11$), T4 ($p=0.21$), T5 ($p=0.36$), T6 ($p=0.32$), P3 ($p=0.06$) and P4 ($p=0.06$) electrodes (Table III).

TABLE II
P300 LATENCY OF THE TARGET TONES

Electrode site	Ambivert (mean±SD)	Extravert (mean±SD)	p-value	Significance
Fz	539.2±76.7	515.6±139	0.55	NS
Cz	503.2±114	500.9±115	0.92	NS
Pz	513.7±123	520.6±124	0.96	NS
T3	537.0±90.3	525.1±110	0.74	NS
T4	513.2±73.7	497.7±136	0.55	NS
T5	508.2±93.5	535.8±112	0.36	NS
T6	509.6±90.7	533.9±134	0.53	NS
P3	506.8±105	531.8±98.4	0.56	NS
P4	513.1±111	533.6±123	0.55	NS

NS: Not significant; SD: Standard Deviation

TABLE III
P300 LATENCY OF THE STANDARD TONES

Electrode site	Ambivert (mean±SD)	Extravert (mean±SD)	p-value	Significance
Fz	511.0±67.0	468.7±109	0.04	S
Cz	463.0±109	415.6±119	0.14	NS
Pz	485.4±107	432.8±130	0.11	NS
T3	508.7±81.0	466.9±94.7	0.05	S
T4	496.1±81.9	462.7±99.8	0.21	NS
T5	492.2±102	469.1±116	0.36	NS
T6	493.2±77.7	469.4±104	0.32	NS
P3	504.4±97.4	465.5±111	0.06	NS
P4	506.5±80.6	462.1±120	0.06	NS

NS: Not significant; S: Significant; SD: Standard Deviation

C. P300 Amplitude

Mean amplitude for the target tone was between $3.08\mu\text{V}$ and $5.32\mu\text{V}$ in the ambivert group and $2.45\mu\text{V}$ to $5.9\mu\text{V}$ in the extravert group. For the standard tone, it was between $1.08\mu\text{V}$ to $3.30\mu\text{V}$ in the ambivert group and $0.88\mu\text{V}$ to $2.57\mu\text{V}$ in the extravert group. There was insignificant difference in the amplitude for the target tone between ambiverts group and extravert group at all electrodes Fz ($P=0.46$), Cz ($P=0.64$), Pz ($P=0.24$), T3 ($P=0.30$), T4 ($P=0.38$), T5 ($P=0.38$), T6 ($P=0.36$), P3 ($P=0.35$) and P4 ($P=0.72$) (Table IV). Similarly, there was an insignificant difference at all electrode sites for the standard tone between ambivert group and extravert group Fz ($p=0.21$), Cz ($P=0.70$), Pz ($P=0.47$), T3 ($P=0.51$), T4 ($P=0.27$), T5 ($P=0.51$), T6 ($P=0.55$), P3 ($P=0.27$), P4 ($P=0.12$) (Table V).

TABLE IV
P300 AMPLITUDE OF THE TARGET TONE

Electrode site	Ambivert (mean±SD)	Extravert (mean±SD)	p-value	Significance
Fz	5.32±2.53	5.97±3.41	0.46	NS
Cz	3.08±2.78	3.11±2.04	0.64	NS
Pz	4.87±2.60	5.87±2.60	0.24	NS
T3	3.32±2.76	3.63±1.75	0.30	NS
T4	3.26±3.03	2.45±1.92	0.38	NS
T5	4.42±2.94	4.46±1.66	0.38	NS
T6	4.05±2.23	3.55±2.56	0.36	NS
P3	3.64±2.00	4.37±2.01	0.35	NS
P4	3.46±1.73	3.71±1.97	0.72	NS

NS: Not significant; SD: Standard Deviation

TABLE V
P300 AMPLITUDE OF THE STANDARD TONE

Electrode Site	Ambivert (mean±SD)	Extravert (mean±SD)	p-value	Significance
Fz	3.30±1.57	2.57±1.49	0.21	NS
Cz	1.13±0.67	1.26±0.80	0.70	NS
Pz	1.71±1.35	1.91±1.15	0.47	NS
T3	2.01±1.10	1.76±1.14	0.51	NS
T4	2.28±1.27	1.69±1.03	0.27	NS
T5	2.10±1.61	1.78±1.14	0.51	NS
T6	2.38±1.43	2.01±1.08	0.55	NS
P3	1.08±0.62	0.88±0.58	0.27	NS
P4	1.80±0.77	1.42±0.77	0.12	NS

NS: Not significant; SD: Standard Deviation

IV. DISCUSSION

The present study aims to determine auditory cognitive function between extraversion and ambiversion types of personality by using auditory oddball paradigm of the event related potential. Findings of this study showed insignificantly larger P300 amplitude in the extravert group at Fz, Cz, Pz, T3, T5, P3 and P4, whereas ambivert group demonstrated larger amplitude at T4 and T6 for the target tone. Ambivert group showed insignificant larger amplitude for the standard tone at Fz, T3, T4, T5, T6, P3 and P4, while extravert group showed insignificant larger amplitude at Cz and Pz. Additionally, latency for the target tone appeared longer in ambivert group at electrodes Fz, Cz, T3 and T4, while extravert group showed longer latency at Pz, T5, T6, P3 and P4 electrodes. However, the difference was insignificant. For the standard tone, latency was longer in the ambiverts group at all electrode sites in comparison with extravert group, showing significance at Fz and T3 and insignificant at Cz, Pz, T4, T5, T6, P3 and P4 electrodes. It can be seen that the extravert group has larger amplitude, yet the difference is insignificant. In a related study to determine the relation between auditory P300 event related potential and major dimensions of personality, finding yielded a larger amplitude with a positive relation among extraversion, agreeableness, openness and conscientiousness trait, but the amplitude was negatively related to neuroticism. The study did not report statistically significant overall relationship between P300 latency and personality traits [16].

V. CONCLUSION

Extravert and ambivert indicate similar characteristic in cognition processing, as measured by the auditory task, presented in Event Related Potential. This finding might explain the nature of this type of personality. However, the weakness of the respondent selection should be taken into consideration. Future research is suggested to delve further the relation between neuro-cognition and personality, in different age group and population.

ACKNOWLEDGMENT

Author thanks Universiti Sains Malaysia, Kubang Kerian, Kelantan, Malaysia for the financial support (USM Short Term Research Grant – 304/PPSP/61312101) and providing research facilities.

REFERENCES

- [1] D. P. McAdams and J. L. Pals, "A new Big Five: fundamental principles for an integrative science of personality," *American Psychologist*, vol.61, no.3, pp.204, 2006.
- [2] V., Devadoss & M. C. J. Anand, "Dimensions of Personality of Women in Chennai using CETD Matrix," *International Journal of Computer Applications*, vol.50, no.5, pp.0975-8887, 2012.
- [3] G. Matthews, I. J. Deary and M.C. Whiteman, *Personality Traits*, Cambridge University Press, 2003.
- [4] M. C. O'Connor and S.V. Paunonen, "Big Five personality predictors of post-secondary academic performance", *Personality and Individual Differences*, vol.43, no.5, pp.971-990, 2007.
- [5] Di Fabio and D.H. Saklofske, "Comparing ability and self-report trait emotional intelligence, fluid intelligence, and personality traits in career decision", *Personality and Individual Differences*, vol. 64, pp. 174-178, 2014.
- [6] S. Dunkel and D. Van der Linden, "Evidence for the general factor of personality as social-effectiveness", *Personality and Individual Differences*, vol. 64, pp. 147-151, 2014.
- [7] H. J. Eysenck, "Dimensions of personality: 16, 5 or 3?—Criteria for a taxonomic paradigm", *Personality and Individual Differences*, vol.12, no.8, pp. 773-790, 1991.
- [8] P. T. Costa and R.R. MacCrae, *Revised NEO Personality Inventory (NEO PI-R) and NEO Five-Factor Inventory (NEO FFI): Professional Manual: Psychological Assessment Resources*, 1992
- [9] E. C. Tupes and R.E. Christal, "Recurrent Personality Factors Based on Trait Ratings", *Journal of Personality*, vol. 60, no.2, pp 225-251, 1992.
- [10] P. T. Costa Jr & T. A. Widiger, "Personality disorders and the five-factor model of personality", *American Psychological Association*, 1994.
- [11] D. Cohen & J. P. Schmidt, "Ambiversion: characteristics of midrange responders on the Introversion-Extraversion continuum", *Journal of Personality Assessment*, vol. 43, no.5, pp. 514-516, 1979.
- [12] M. Crowe, R. Andel, N. L. Pedersen, L. Fratiglioni, & M. Gatz, "Personality and risk of cognitive impairment 25 years later", *Psychology and Aging*, vol.21, no.3, pp.573, 2006.
- [13] D. S. Goodin, "Clinical utility of long latency 'cognitive' event-related potentials (P3): the pros," *Electroencephalography and Clinical Neurophysiology*, vol. 76, no. 1, pp. 2-5, 1990.
- [14] J. Polich, "Updating P300: an integrative theory of P3a and P3b", *Clinical Neurophysiology*, vol. 118, no.10, pp. 2128-2148, 2007.
- [15] M. Yusoff, A. Rahim and A. R. Esa. The USM personality inventory (USMaP-i) manual (Online). Available: <http://www.medic.usm.my/dme/images/stories/staff/KKMED/2010/manual%20usmap-i.pdf>, 2010. (Retrieved on September 20, 2015).
- [16] R. J. Gurrera, D. F. Salisbury, B. F., O'Donnell, P. G. Nestor & R. W. McCarley, "Auditory P3 indexes personality traits and cognitive function in healthy men and women", *Psychiatry Res*, vol.133, no.2-3, pp. 215-228, 2005.



Dr Mohd Nasir Che Mohd Yusoff was born in Selangor, Malaysia on 23 October 1975. He obtained a degree of Doctor of Philosophy with specialization in Health Psychology in 2009 from University of Malaya, Kuala Lumpur, Malaysia. He is currently working as Senior Lecturer at Department of Neurosciences, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150, Kubang Kerian, Kelantan. Currently, Dr Yusoff is actively participating in Neuroscience Club Association for Kelantan Malaysia Chapter. This association is headed by the United State of America Neuroscience Club. Under this association, many activities have been implemented at society level by the financial support provided by the club.



Ahmad Adamu Adamu was born in Nigeria. He obtained a Bachelor Science of Occupational Therapies from University of Nigeria. He is currently studying Master of Neuroscience at Department of Neurosciences, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150, Kubang Kerian, Kelantan. Currently, Ahmad is actively participating in Neuroscience Club Association for Kelantan Malaysia Chapter. This association is headed by the United State of America Neuroscience Club. Under this association, many activities have been implemented at society level by the financial support provided by the club.



Dr. Tahamina Begum was born in Bangladesh in 1970. She graduated with MBBS in 1996 from Sher-E-Bangla Medical College, Barisal, Bangladesh. In 2006, she was awarded with Doctor of Philosophy from Kyoto University, Japan with specialization in neuroscience. She is currently working as Senior Medical Lecturer at Department of Neurosciences, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150, Kubang Kerian, Kelantan. Currently, Dr Begum is actively participating in Neuroscience Club Association for Kelantan Malaysia Chapter. This association is headed by the United State of America Neuroscience Club. Under this association, many activities have been implemented at society level by the financial support provided by the club.



Dr Mohammed Faruque Reza was born in Bangladesh in 1967. He graduated with MBBS from Rangpur Medical College, Rajshahi University, Bangladesh in 1993. Few years after that, he obtained Master Science in Microbiology from Dhaka University, Bangladesh. In 2005, he was awarded with Doctor of Philosophy from Hokkaido University, Japan with specialization in neuroscience field. He is currently working as Senior Medical Lecturer at Department of Neurosciences, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150, Kubang Kerian, Kelantan. Currently, Dr Reza is actively participating in Neuroscience Club Association for Kelantan Malaysia Chapter. This association is headed by the United State of America Neuroscience Club. Under this association, many activities have been implemented at society level by the financial support provided by the club.