

Morphological Parameters and Selection of Turkish Edible Seed Pumpkins (*Cucurbita pepo* L.) Germplasm

Onder Turkmen, Musa Seymen, Sali Fidan, Mustafa Paksoy

Abstract—There is a requirement for registered edible seed pumpkin suitable for eating in Turkey. A total of 81 genotypes collected from the researchers in 2005 originated from Eskisehir, Konya, Nevşehir, Tekirdağ, Sakarya, Kayseri and Kırşehir provinces were utilized. The used genetic materials were brought to S5 generation by the research groups among 2006 and 2010 years. In this research, S5 stage reached in the genotype given some of the morphological features, and selection of promising genotypes generated scale were made. Results showed that the A-1 (420), A-7 (410), A-8 (420), A-32 (420), B-17 (410), B-24 (410), B-25 (420), B-33 (400), C-24 (420), C-25 (410), C-26 (410) and C-30 (420) genotypes are expected to be promising varieties.

Keywords—Candidate cultivar, edible seed pumpkin, morphologic parameters, selection.

I. INTRODUCTION

PUMPKIN is a kind of vegetable which is used for human nutrition through processing its fruit, flowers and seed. Although pumpkin is generally produced for its fruit [1]. The enculturation processes may be conducted through machines to a large extent, and less affected by diseases and pests. Growing pumpkin seed is important, because it does not require frequent irrigation and may be grown even in completely infertile conditions. The pumpkin seed grown in Turkey are generally made in the type of *Cucurbita pepo* and the type of *Cucurbita moschata* in small amounts [2], [3].

The content of oil and protein are high in the pumpkin seeds with 40-50% oil ratio [4]-[8], 30-40% protein ratio [4], [7]-[9]. Additionally, the pumpkin seed oil is used in human nutrition as salad oil in many countries such as Austria, Slovenia, and Hungary since it has a rich ingredient of carbohydrate, 20-25%, [7] and vitamin E [10]. Similarly, the pumpkin seed oil is consumed in soups and minced meat [11]. However, the use as a cooking oil remain limited due to its color, strong aroma and foaming [7], [12], [13]. It is reported that the pumpkin seed oil prevents the development of prostate cancer and it is heart-friendly [14]. It is also emphasized that consuming 70-80 g of pumpkin seed a day for health [15].

Onder Turkmen and Mustafa Paksoy are with the Selcuk University Agriculture Faculty and Department of Horticulture, Konya, 42075, Turkey (e-mail: turkmen@selcuk.edu.tr, paksoy@selcuk.edu.tr).

Musa Seymen is with the Selcuk University Agriculture Faculty and Department of Horticulture, Konya, 42075, Turkey (phone: 90 543 7958010; fax: 90 332 241 0108; e-mail: mseymen@selcuk.edu.tr).

Sali Fidan is with the Transitional Zone Agricultural Research Institute, Eskişehir, Turkey (e-mail: sfidan@yahoo.com).

Since pumpkin has monoic flower structure, the rate of pseudo-fertilization is rather high. For that reason, it is possible that the lines emerging in the end may be different from the type of the original seed. Therefore, one of the biggest problems in growing pumpkin seed as appetizers is the problem of species [16]-[18]. The pumpkin seed in Turkey and other countries is consumed as edible seed.

In this study, it was aimed to obtain pure lines in order to develop candidates of pumpkin seed. In accordance with this purpose, herbal characteristics of 81 pumpkin genotypes at the level of S5 were determined. Furthermore, scoring was conducted in terms of definition of parameters and the species which may become as a candidate were presented.

II. MATERIAL AND METHODS

Plant material employed in the present research was collected from the regions of Turkey; Eskişehir, Konya, Nevşehir, Tekirdağ, Sakarya, Kayseri and Kırşehir through the utilities of Selçuk University Faculty of Agriculture Horticultural Department and the Institute of Anatolian Agricultural Researches.

The study was established in the field of the Institute of Anatolian Agricultural Researches on 20th of May 2011. The planting distance was 2x1.5 m so that there were 20 plants from each genotype. Sowing was conducted by hand and 2-3 seeds were put into each pit and hand thinning was conducted one week after the germination because of remaining one plant in each pit. The processes such as weeding, middle-breaking, watering and fertilizing were regularly conducted on the plants.

In the plants at the flowering period, the pollination was provided through closing the male and female flowers in the evening, plucking the male flower early in the morning.

Plant observations were conducted during start of flowering period; the seed characteristics of fruit were taken after the harvest on September 12-17 in 2011 and after the seeds were taken out.

Mean of the results obtained from the study were taken and interpreted. In the selection of genotypes, the characteristics with defined forms were considered and the determined characteristics were scored as seen in Table I. The genotypes were evaluated through a total score of 500 and they were classified according to their score.

TABLE I
SELECTION CRITERIA FOR THE CLASS AND GIVEN RELATIVE SCORES [19],
[20]

| Selection criteria | | Sc |
|---------------------------------|--------------------|----|
| Seed shape | Narrow elliptical | 4 |
| | Elliptical | 5 |
| | Large elliptical | 4 |
| Seed color | Light cream | 4 |
| | Cream | 5 |
| | Dark cream | 3 |
| 1000 seed weight | Big (>250 g) | 5 |
| | Middle (200-250 g) | 3 |
| | Minor (<200 g) | 1 |
| Fruit/plant | 2-3 | 3 |
| | 4-5 | 5 |
| | 6-7 | 4 |
| Seed yield/fruit | < 50 g | 2 |
| | 50-80 g | 4 |
| | > 80 g | 5 |
| Seed yield /plant | 100-200 g | 2 |
| | 200-400 g | 4 |
| | >400 g | 5 |
| Easiness of seed coat crackling | Easy | 3 |
| | Difficult | 1 |

III. RESULTS AND DISCUSSION

The lines are collected from different regions of Turkey. Measurements and observations taken from the 81 edible seed pumpkin genotypes were shown in Tables II-IV. Table V shows the total weighted score for each genotype, the rating results.

It was found in the study that 30 genotypes (37%) were vertical, 25 genotypes (31%) were semi-creeping ones, and 26 of them were creeping plants. Among those genotypes, 75

(92%) put forth branches while 6 (7%) did not put branches forth. The 17 of the genotypes (21%) had silver stains on their leaves while 64 (79%) had no stains. When the lobes on the leaves were analyzed, 3 (4%) had few if any lobes, 37 (46%) had few lobes, 32 (39%) had moderate number of lobes and 9 (11%) had many lobes. 30 (37%) of the fruits which are suitable for seeding had no spots, 17 (21%) had yellow spots while 34 (42%) had green spots. Among those genotypes, 11 (14%) had little density of spots, 25 (31%) had intensive and 15 (18%) had higher density of spots. When the color of ripe fruit was analyzed, 21 (24%) were yellow, 2 (2%) were green, 15 (18%) were green-yellow greyish, 22 (27%) were dark yellow-green greyish, 17 (21%) were light yellow and 4 (5%) were dark yellow (Table II).

When the genotypes were analyzed, 59 (72%) of the genotypes had short, 18 (22%) had medium and 4 (5%) had long fruit sizes. When the fruit width of the genotypes was analyzed, 1 (1%) was found narrow, 13 (16%) were medium and 67 (83%) were wide. The fruit index of 70 (86%) genotypes were found round, 8 (10%) were elliptic and 3 (4%) were long. As for the seed index, 1 (1%) was narrow elliptic, 36 (44%) were elliptic and 44 (54%) were wide elliptic. The seeds of 12 (15%) genotypes were found small, 40 (49%) genotypes were found medium and 29 (36%) were found big. When the color of the seeds was analyzed, 7 (9%) of the genotypes were found light cream, 68 (84%) were cream and 6 (7%) were dark cream. Cracking of 68 (84%) genotypes was easy while 13 (16%) were difficult to crack (Table III).

TABLE II
PLANT, LEAF, FLOWER AND FRUIT IN THE S5 STAGE OF EDIBLE SEED PUMPKIN

| Genotype | Appearance of plant | Branch prolongation | Silvery Leaves Stain | Slice of leaf | Spot color | Spot density | Fruit color |
|----------|---------------------|---------------------|----------------------|---------------|------------|--------------|-------------|
| A1 | 1 | 1 | 2 | 2 | 4 | 3 | 5 |
| A3 | 2 | 1 | 2 | 2 | 4 | 2 | 5 |
| A4 | 2 | 1 | 2 | 2 | 2 | 2 | 4 |
| A5 | 3 | 1 | 2 | 3 | 4 | 2 | 2 |
| A6 | 2 | 1 | 2 | 3 | - | - | 6 |
| A7 | 3 | 1 | 1 | 2 | 4 | 1 | 2 |
| A8 | 1 | 2 | 2 | 3 | - | - | 6 |
| A10 | 2 | 1 | 1 | 2 | - | - | 2 |
| A11 | 1 | 2 | 2 | 2 | 2 | 1 | 3 |
| A12-1 | 1 | 1 | 2 | 3 | 4 | 2 | 5 |
| A12-2 | 3 | 1 | 2 | 3 | 4 | 2 | 5 |
| A13 | 3 | 1 | 1 | 2 | 2 | 3 | 4 |
| A14 | 1 | 2 | 2 | 3 | 4 | 3 | 5 |
| A15 | 3 | 1 | 2 | 2 | 4 | 1 | 7 |
| A16 | 2 | 1 | 1 | 2 | 2 | 3 | 4 |
| A18 | 1 | 1 | 2 | 2 | 2 | 2 | 4 |
| A20 | 3 | 1 | 1 | 2 | 4 | 2 | 5 |
| A21 | 3 | 1 | 2 | 1 | 2 | 2 | 4 |
| A23-1 | 2 | 1 | 2 | 3 | 4 | 2 | 5 |
| A23-2 | 2 | 1 | 1 | 3 | 2 | 3 | 4 |
| A24 | 2 | 1 | 2 | 3 | 2 | 2 | 4 |
| A25 | 1 | 1 | 2 | 4 | 2 | 2 | 4 |
| A26 | 2 | 1 | 2 | 3 | - | - | 2 |
| A28 | 3 | 1 | 2 | 4 | - | - | 2 |
| A29 | 3 | 1 | 2 | 3 | - | - | 6 |

| Genotype | Appearance of plant | Branch prolongation | Silvery Leaves Stain | Slice of leaf | Spot color | Spot density | Fruit color |
|----------|---------------------|---------------------|----------------------|---------------|------------|--------------|-------------|
| A31 | 1 | 1 | 1 | 3 | 4 | 1 | 6 |
| A32 | 3 | 1 | 2 | 2 | 4 | 2 | 5 |
| A33 | 1 | 1 | 2 | 3 | 2 | 2 | 4 |
| A34 | 2 | 1 | 2 | 2 | 2 | 2 | 4 |
| B1 | 2 | 1 | 2 | 2 | - | - | 2 |
| B2 | 2 | 1 | 1 | 2 | - | - | 6 |
| B4 | 1 | 1 | 2 | 3 | 2 | 3 | 4 |
| B8 | 3 | 1 | 2 | 3 | 4 | 3 | 5 |
| B11 | 1 | 1 | 1 | 3 | - | - | 2 |
| B12 | 1 | 1 | 2 | 2 | 2 | 2 | 4 |
| B13 | 1 | 1 | 2 | 3 | - | - | 6 |
| B14 | 3 | 1 | 2 | 2 | - | - | 6 |
| B16 | 3 | 1 | 2 | 3 | - | - | 2 |
| B17 | 2 | 1 | 2 | 3 | - | - | 6 |
| B18 | 2 | 1 | 1 | 1 | - | - | 2 |
| B19 | 1 | 1 | 2 | 4 | 4 | 2 | 2 |
| B20 | 2 | 1 | 2 | 2 | - | - | 6 |
| B21 | 2 | 1 | 2 | 2 | - | - | 2 |
| B22 | 1 | 1 | 2 | 2 | 4 | 1 | 6 |
| B23 | 2 | 1 | 1 | 2 | - | - | 6 |
| B24 | 3 | 1 | 2 | 3 | 4 | 2 | 5 |
| B25 | 3 | 1 | 1 | 1 | 4 | 2 | 5 |
| B26 | 3 | 1 | 2 | 3 | - | - | 2 |
| B27-1 | 1 | 1 | 2 | 4 | - | - | 2 |
| B27-2 | 1 | 1 | 2 | 3 | - | - | 2 |
| B28 | 3 | 1 | 1 | 2 | 2 | 2 | 7 |
| B31 | 1 | 1 | 2 | 2 | 4 | 2 | 5 |
| B32 | 3 | 1 | 2 | 2 | 4 | 1 | 2 |
| B33 | 1 | 1 | 2 | 2 | 4 | 1 | 7 |
| B34-1 | 1 | 1 | 2 | 3 | 4 | 2 | 5 |
| B34-2 | 2 | 1 | 2 | 4 | 4 | 2 | 5 |
| C1 | 1 | 1 | 2 | 4 | 4 | 3 | 5 |
| C2 | 2 | 1 | 2 | 3 | - | - | 6 |
| C3 | 2 | 1 | 2 | 3 | - | - | 6 |
| C4 | 3 | 1 | 2 | 2 | 4 | 2 | 5 |
| C5 | 1 | 1 | 2 | 4 | - | - | 6 |
| C7 | 3 | 1 | 2 | 2 | - | - | 6 |
| C8 | 1 | 1 | 2 | 4 | 4 | 2 | 5 |
| C11 | 2 | 1 | 2 | 2 | - | - | 2 |
| C12 | 1 | 2 | 2 | 2 | 4 | 1 | 6 |
| C13 | 2 | 1 | 2 | 2 | 2 | 3 | 4 |
| C16 | 1 | 2 | 1 | 3 | 2 | 3 | 4 |
| C17 | 1 | 1 | 2 | 3 | - | - | 6 |
| C18 | 1 | 1 | 2 | 2 | 4 | 3 | 5 |
| C19 | 3 | 1 | 2 | 2 | - | - | 2 |
| C21 | 1 | 1 | 1 | 3 | 4 | 1 | 7 |
| C24 | 2 | 1 | 1 | 3 | - | - | 2 |
| C25 | 1 | 1 | 2 | 4 | 4 | 3 | 5 |
| C26 | 1 | 2 | 2 | 2 | 2 | 3 | 4 |
| C27 | 3 | 1 | 2 | 2 | 4 | 1 | 2 |
| C28 | 3 | 1 | 2 | 3 | 4 | 2 | 5 |
| C30 | 3 | 1 | 2 | 2 | - | - | 2 |
| C31 | 2 | 1 | 2 | 3 | 4 | 3 | 5 |
| C32 | 2 | 1 | 2 | 3 | - | - | 3 |
| C33 | 3 | 1 | 1 | 3 | 4 | 3 | 5 |
| C34 | 2 | 1 | 2 | 2 | 4 | 1 | 2 |

Note: Appearance of plant: perpendicular-1, semi-wrapping-2, wrapping-3; Branch prolongation: presence-1, absence-2; Silvery Leaves Stain: presence-1, absence-2; Slice of leaf: presence or very few-1, slight-2, medium-3, excess-4, excessive-5; Spot color on the matured fruit: cream-1, yellow-2, orange-3, green-4; J=Density of spot: slight-1, intense-2, excess-3; K=Color on matured fruit: cream-1, yellow-2, green-3, green-yellow-mealy-4, dark yellow-green mealy-5, light yellow-6, dark yellow-7, orange-8.

TABLE III
FRUIT AND SEED CHARACTERISTICS IN THE S5 STAGE OF EDIBLE SEED PUMPKIN

| Genotype | Fruit size | Fruit diameter | Fruit index | Seed index | Seed weight | Seed color | Easiness of seed coat crackling |
|----------|------------|----------------|-------------|-------------------|-------------|-------------|---------------------------------|
| A1 | Middle | Wide | Circular | Elliptical | Big | Cream | Difficult |
| A3 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| A4 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| A5 | Short | Wide | Circular | Elliptical | Big | Dark cream | Easy |
| A6 | Short | Wide | Circular | Large elliptical | Big | Cream | Easy |
| A7 | Short | Wide | Circular | Elliptical | Big | Cream | Easy |
| A8 | Middle | Wide | Circular | Elliptical | Big | Cream | Easy |
| A10 | Short | Wide | Circular | Elliptical | Middle | Cream | Easy |
| A11 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| A12-1 | Short | Wide | Circular | Elliptical | Middle | Cream | Easy |
| A12-2 | Short | Wide | Circular | Large elliptical | Big | Cream | Easy |
| A13 | Short | Wide | Circular | Elliptical | Middle | Light cream | Easy |
| A14 | Long | Middle | Lengthy | Large elliptical | Minor | Cream | Easy |
| A15 | Short | Wide | Circular | Large elliptical | Big | Cream | Easy |
| A16 | Short | Wide | Circular | Elliptical | Big | Cream | Difficult |
| A18 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| A20 | Short | Wide | Circular | Elliptical | Middle | Cream | Easy |
| A21 | Short | Wide | Circular | Elliptical | Minor | Cream | Easy |
| A23-1 | Middle | Wide | Circular | Large elliptical | Big | Dark cream | Easy |
| A23-2 | Middle | Wide | Lengthy | Elliptical | Big | Cream | Easy |
| A24 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| A25 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| A26 | Middle | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| A28 | Short | Middle | Circular | Large elliptical | Big | Cream | Easy |
| A29 | Short | Middle | Circular | Large elliptical | Middle | Cream | Easy |
| A31 | Short | Middle | Circular | Elliptical | Middle | Cream | Easy |
| A32 | Short | Wide | Circular | Large elliptical | Big | Cream | Easy |
| A33 | Short | Wide | Circular | Large elliptical | Big | Cream | Easy |
| A34 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| B1 | Short | Wide | Circular | Elliptical | Middle | Dark cream | Easy |
| B2 | Short | Wide | Circular | Large elliptical | Minor | Cream | Easy |
| B4 | Short | Wide | Circular | Large elliptical | Big | Cream | Easy |
| B8 | Short | Wide | Circular | Large elliptical | Big | Cream | Easy |
| B11 | Short | Wide | Circular | Narrow elliptical | Minor | Light cream | Difficult |
| B12 | Short | Wide | Circular | Large elliptical | Middle | Dark cream | Easy |
| B13 | Middle | Middle | Elliptical | Large elliptical | Middle | Cream | Easy |
| B14 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| B16 | Short | Middle | Circular | Large elliptical | Middle | Cream | Easy |
| B17 | Middle | Wide | Circular | Elliptical | Big | Cream | Easy |
| B18 | Middle | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| B19 | Middle | Wide | Elliptical | Elliptical | Middle | Cream | Easy |
| B20 | Middle | Wide | Circular | Large elliptical | Big | Cream | Easy |
| B21 | Short | Wide | Circular | Large elliptical | Minor | Cream | Difficult |
| B22 | Short | Middle | Elliptical | Elliptical | Middle | Light cream | Easy |
| B23 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| B24 | Middle | Wide | Elliptical | Elliptical | Big | Cream | Easy |
| B25 | Short | Wide | Circular | Elliptical | Big | Cream | Easy |
| B26 | Short | Wide | Circular | Elliptical | Minor | Cream | Easy |
| B27-1 | Short | Middle | Elliptical | Elliptical | Middle | Cream | Difficult |
| B27-2 | Short | Middle | Circular | Elliptical | Minor | Cream | Easy |
| B28 | Short | Wide | Circular | Large elliptical | Minor | Cream | Easy |
| B31 | Middle | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| B32 | Short | Wide | Circular | Large elliptical | Middle | Light cream | Easy |
| B33 | Short | Wide | Circular | Large elliptical | Big | Cream | Easy |
| B34-1 | Middle | Wide | Circular | Large elliptical | Middle | Dark cream | Easy |
| B34-2 | Short | Wide | Circular | Elliptical | Middle | Cream | Easy |
| C1 | Short | Wide | Circular | Large elliptical | Minor | Cream | Easy |
| C2 | Short | Middle | Circular | Large elliptical | Minor | Cream | Easy |

| Genotype | Fruit size | Fruit diameter | Fruit index | Seed index | Seed weight | Seed color | Easiness of seed coat crackling |
|----------|------------|----------------|-------------|------------------|-------------|-------------|---------------------------------|
| C3 | Short | Wide | Circular | Large elliptical | Middle | Cream | Difficult |
| C4 | Short | Wide | Circular | Large elliptical | Middle | Light cream | Easy |
| C5 | Middle | Narrow | Lengthy | Large elliptical | Middle | Cream | Easy |
| C7 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| C8 | Middle | Wide | Circular | Elliptical | Big | Cream | Difficult |
| C11 | Short | Wide | Circular | Elliptical | Minor | Dark cream | Easy |
| C12 | Short | Wide | Circular | Large elliptical | Middle | Cream | Easy |
| C13 | Middle | Wide | Elliptical | Elliptical | Middle | Cream | Difficult |
| C16 | Middle | Wide | Circular | Elliptical | Big | Light cream | Easy |
| C17 | Short | Wide | Circular | Large elliptical | Minor | Cream | Difficult |
| C18 | Middle | Middle | Elliptical | Elliptical | Middle | Cream | Easy |
| C19 | Short | Wide | Circular | Large elliptical | Middle | Cream | Difficult |
| C21 | Short | Middle | Circular | Large elliptical | Middle | Cream | Easy |
| C24 | Middle | Wide | Circular | Elliptical | Big | Cream | Easy |
| C25 | Middle | Wide | Elliptical | Elliptical | Big | Cream | Easy |
| C26 | Short | Wide | Circular | Elliptical | Big | Cream | Easy |
| C27 | Short | Wide | Circular | Elliptical | Middle | Cream | Easy |
| C28 | Short | Wide | Circular | Elliptical | Big | Cream | Difficult |
| C30 | Middle | Wide | Circular | Elliptical | Big | Cream | Easy |
| C31 | Short | Wide | Circular | Elliptical | Middle | Cream | Difficult |
| C32 | Short | Wide | Circular | Large elliptical | Big | Cream | Easy |
| C33 | Short | Middle | Circular | Elliptical | Big | Cream | Difficult |
| C34 | Short | Wide | Circular | Elliptical | Middle | Light cream | Difficult |

When the yield factors of the genotypes were analyzed, the average yield per plant was found 114 g and the biggest yield was obtained from A3 genotype as 282 g. It is followed by the genotypes of C31 (226 g), B21 (214 g), B28 (200 g) and B26 (199 g), respectively. The researchers reported that the yield per plant in edible seed pumpkin was found between 200 g and 500 g [17]-[19], [22], [21].

The highest yield per fruit was taken from the genotype A18 with a yield of 160 g and it was followed by B25 (104 g), A8 (102 g), B33 (95 g) and B23-2 (94 g), respectively. Generally, seed yield per fruit was in the expected level and had shown similarity to the conducted studies [16], [19], [21], [23].

TABLE IV
YIELD AND YIELD COMPONENTS IN THE S5 STAGE OF EDIBLE SEED PUMPKIN

| Genotype | Yield (g/plant) | Yield (g/fruit) | Fruit number (number/plant) | Fruit weight (kg) | Seed diameter (mm) | Seed length (mm) | 1000 seed weight (g) |
|----------|-----------------|-----------------|-----------------------------|-------------------|--------------------|------------------|----------------------|
| A1 | 82 | 91 | 0,90 | 3,13 | 12.33 | 20.47 | 256.8 |
| A3 | 282 | 92 | 3,06 | 2,80 | 10.88 | 20.48 | 233.2 |
| A4 | 59 | 43 | 1,35 | 3,50 | 11.16 | 20.41 | 244.0 |
| A5 | 152 | 55 | 2,79 | 4,13 | 11.78 | 20.05 | 300.0 |
| A6 | 103 | 61 | 1,70 | 5,38 | 10.34 | 20.04 | 269.0 |
| A7 | 142 | 59 | 2,42 | 2,73 | 12.98 | 21.24 | 343.8 |
| A8 | 148 | 102 | 1,45 | 3,92 | 11.54 | 20.08 | 335.6 |
| A10 | 55 | 44 | 1,25 | 2,47 | 11.52 | 20.39 | 206.4 |
| A11 | 15 | 12 | 1,30 | 2,55 | 11.69 | 21.67 | 230.6 |
| A12-1 | 159 | 75 | 2,12 | 3,79 | 10.09 | 16.5 | 243.4 |
| A12-2 | 143 | 63 | 2,25 | 3,60 | 12.15 | 22.54 | 268.0 |
| A13 | 169 | 57 | 2,94 | 3,67 | 11.9 | 19.19 | 208.4 |
| A14 | 55 | 37 | 1,50 | 2,87 | 12.15 | 24.13 | 182.2 |
| A15 | 146 | 63 | 2,30 | 2,33 | 9.23 | 19.97 | 273.8 |
| A16 | 90 | 53 | 1,70 | 3,80 | 12.43 | 20.05 | 265.4 |
| A18 | 40 | 160 | 0,25 | 2,75 | 11.78 | 21.17 | 275.4 |
| A20 | 98 | 45 | 2,16 | 1,87 | 10.12 | 17.73 | 211.6 |
| A21 | 24 | 38 | 0,64 | 4,20 | 12.2 | 20.79 | 64.4 |
| A23-1 | 107 | 85 | 1,25 | 5,33 | 11.13 | 21.66 | 296.0 |
| A23-2 | 122 | 94 | 1,30 | 4,85 | 12.16 | 21.27 | 280.2 |
| A24 | 87 | 56 | 1,55 | 2,27 | 9.59 | 17.39 | 213.4 |
| A25 | 149 | 65 | 2,30 | 2,47 | 10.03 | 20.82 | 213.2 |
| A26 | 116 | 77 | 1,50 | 2,80 | 9.12 | 20.97 | 232.0 |
| A28 | 118 | 25 | 4,80 | 1,53 | 11.77 | 21.78 | 261.2 |

| Genotype | Yield (g/plant) | Yield (g/fruit) | Fruit number (number/plant) | Fruit weight (kg) | Seed diameter (mm) | Seed length (mm) | 1000 seed weight (g) |
|----------|-----------------|-----------------|--------------------------------|-------------------|--------------------|------------------|----------------------|
| A29 | 174 | 67 | 2,58 | 2,53 | 10.89 | 20.33 | 230.0 |
| A31 | 112 | 39 | 2,88 | 2,11 | 11.47 | 18.85 | 217.2 |
| A32 | 79 | 53 | 1,47 | 3,25 | 12.51 | 22.88 | 313.8 |
| A33 | 84 | 54 | 1,55 | 3,87 | 11.59 | 20.98 | 268.6 |
| A34 | 91 | 48 | 1,90 | 2,33 | 11.54 | 21.77 | 245.8 |
| B1 | 95 | 56 | 1,70 | 2,73 | 11.24 | 19.62 | 214.2 |
| B2 | 120 | 37 | 3,21 | 2,80 | 11.24 | 21.42 | 180.4 |
| B4 | 126 | 69 | 1,83 | 3,71 | 7.74 | 19.49 | 256.8 |
| B8 | 128 | 57 | 2,25 | 1,87 | 10.93 | 21.68 | 254.2 |
| B11 | 22 | 28 | 0,80 | 2,50 | 11.08 | 16.28 | 180.4 |
| B12 | 83 | 50 | 1,65 | 2,20 | 10.15 | 18.51 | 217.6 |
| B13 | 135 | 67 | 2,00 | 2,73 | 8.87 | 19.44 | 236.6 |
| B14 | 143 | 51 | 2,78 | 2,87 | 10.16 | 18.86 | 208.4 |
| B16 | 199 | 56 | 3,55 | 2,40 | 10.77 | 20.78 | 246.6 |
| B17 | 115 | 64 | 1,80 | 4,00 | 12.99 | 20.27 | 273.4 |
| B18 | 107 | 68 | 1,56 | 4,62 | 10.55 | 20.97 | 242.2 |
| B19 | 79 | 51 | 1,55 | 3,93 | 10.96 | 17.84 | 213.0 |
| B20 | 57 | 76 | 0,75 | 4,30 | 10.44 | 21.06 | 288.8 |
| B21 | 214 | 71 | 3,00 | 2,93 | 9.55 | 17.25 | 184.0 |
| B22 | 95 | 51 | 1,88 | 2,80 | 10.08 | 17.94 | 230.8 |
| B23 | 111 | 53 | 2,11 | 3,33 | 11.41 | 21.13 | 237.6 |
| B24 | 78 | 47 | 1,67 | 4,27 | 11.52 | 20.44 | 276.0 |
| B25 | 94 | 104 | 0,90 | 4,00 | 11.78 | 20.74 | 327.8 |
| B26 | 78 | 41 | 1,92 | 2,83 | 12.36 | 20.8 | 171.4 |
| B27-1 | 103 | 37 | 2,75 | 2,13 | 11.99 | 18 | 215.4 |
| B27-2 | 156 | 73 | 2,15 | 2,00 | 11.31 | 18.1 | 151.6 |
| B28 | 200 | 61 | 3,30 | 2,47 | 9.84 | 21.97 | 187.8 |
| B31 | 162 | 67 | 2,40 | 3,00 | 9.31 | 19.17 | 227.6 |
| B32 | 177 | 45 | 3,89 | 1,27 | 9.31 | 19.84 | 202.2 |
| B33 | 90 | 95 | 0,95 | 3,80 | 10.53 | 19.52 | 256.0 |
| B34-1 | 120 | 57 | 2,13 | 3,13 | 10.18 | 19.66 | 214.0 |
| B34-2 | 88 | 47 | 1,89 | 2,60 | 11.29 | 19.23 | 244.2 |
| C1 | 100 | 69 | 1,45 | 2,93 | 10.85 | 20.23 | 194.0 |
| C2 | 142 | 38 | 3,74 | 1,27 | 8.3 | 22.21 | 188.0 |
| C3 | 99 | 35 | 2,80 | 1,93 | 8.93 | 22.32 | 213.2 |
| C4 | 68 | 75 | 0,90 | 3,90 | 10.4 | 18.67 | 248.8 |
| C5 | 106 | 69 | 1,55 | 2,67 | 10.32 | 19.43 | 214.0 |
| C7 | 143 | 33 | 4,37 | 1,27 | 9.6 | 17.69 | 205.2 |
| C8 | 101 | 42 | 2,40 | 3,73 | 11.91 | 19.67 | 267.2 |
| C11 | 131 | 51 | 2,55 | 2,20 | 11.42 | 19.77 | 176.0 |
| C12 | 84 | 62 | 1,37 | 2,31 | 10.19 | 18.67 | 209.2 |
| C13 | 65 | 57 | 1,15 | 3,33 | 11.96 | 18.97 | 235.4 |
| C16 | 44 | 71 | 0,61 | 3,43 | 12.52 | 21.69 | 332.8 |
| C17 | 103 | 45 | 2,30 | 2,20 | 10.06 | 21.19 | 197.4 |
| C18 | 90 | 69 | 1,30 | 3,00 | 11.52 | 20.09 | 240.0 |
| C19 | 112 | 55 | 2,05 | 2,47 | 10.89 | 21.2 | 205.6 |
| C21 | 157 | 69 | 2,26 | 2,80 | 9.2 | 16.69 | 225.4 |
| C24 | 129 | 85 | 1,53 | 5,36 | 11.67 | 20.62 | 264.8 |
| C25 | 119 | 68 | 1,75 | 3,07 | 11.8 | 19.95 | 323.4 |
| C26 | 181 | 75 | 2,42 | 2,47 | 10.81 | 19.27 | 269.0 |
| C27 | 67 | 39 | 1,72 | 4,00 | 11.91 | 21.13 | 235.0 |
| C28 | 110 | 47 | 2,35 | 3,27 | 11.45 | 19.89 | 278.2 |
| C30 | 95 | 83 | 1,15 | 6,20 | 11.94 | 20.71 | 290.4 |
| C31 | 226 | 71 | 3,20 | 3,00 | 9.26 | 19.38 | 213.0 |
| C32 | 103 | 50 | 2,05 | 2,33 | 11.28 | 22.49 | 332.8 |
| C33 | 166 | 55 | 3,00 | 2,67 | 11.1 | 18.85 | 266.8 |
| C34 | 123 | 53 | 2,30 | 2,13 | 10.27 | 17.59 | 206.0 |
| Mean | 114 | 59 | 2.02 | 3.06 | 10.92 | 20.05 | 239 |

TABLE V
CLASSIFICATION SCORE IN THE S5 STAGE OF EDIBLE SEED PUMPKIN

| Genotype | Seed shape | Seed color | 1000 seed weight | Fruit/plant | Seed yield/fruit | Seed yield /plant | Easiness of seed coat crackling | Total score |
|----------|------------|------------|------------------|-------------|------------------|-------------------|---------------------------------|-------------|
| A1 | 100 | 75 | 100 | 30 | 50 | 20 | 45 | 420 |
| A3 | 80 | 75 | 60 | 50 | 50 | 40 | 45 | 400 |
| A7 | 100 | 75 | 100 | 30 | 40 | 20 | 45 | 410 |
| A8 | 100 | 75 | 100 | 30 | 50 | 20 | 45 | 420 |
| A32 | 100 | 75 | 100 | 30 | 40 | 20 | 45 | 410 |
| B17 | 100 | 75 | 100 | 30 | 40 | 20 | 45 | 410 |
| B24 | 100 | 75 | 100 | 30 | 40 | 20 | 45 | 410 |
| B25 | 100 | 75 | 100 | 30 | 50 | 20 | 45 | 420 |
| B33 | 80 | 75 | 100 | 30 | 50 | 20 | 45 | 400 |
| C24 | 100 | 75 | 100 | 30 | 50 | 20 | 45 | 420 |
| C25 | 100 | 75 | 100 | 30 | 40 | 20 | 45 | 410 |
| C26 | 100 | 75 | 100 | 30 | 40 | 20 | 45 | 410 |

The average number of fruits per plant was 2.02 fruit /plant. The highest number of fruits was taken from the genotype C7 with 4.37 pieces and it is followed by B32 (3.89 pieces), C2 (3.74 pieces), B16 (3.55 pieces) and B28 (3.3 pieces), respectively. The researchers determined that the number of fruit per plant in the edible seed pumpkin was between 1 and 9 [16], [19], [21], [22]. In our study, the scarcity of yield per plant resulting from the scarcity of fruit per plant comes from uncontrollably picking the inseminated fruit in the plants until the genotypes are appropriated. When the average fruit weight of the genotypes was considered, the heaviest fruit was obtained from C30 genotype with 6.20 kg; it was followed by the genotypes of A6 (5.38 kg), C24 (5.36 kg), A23-1 (5.33 kg) and A23-2 (4.85 kg), respectively.

When the seed width was analyzed, the highest value was obtained from B17 genotype with 12.99 mm and it was followed by the genotypes A7 (12.98 mm), C16 (12.52 mm), A16 (12.43 mm) and B26 (12.36 mm), respectively. The biggest seed height was obtained from the genotype A14 with 24.13 mm and it is followed by the genotypes of A32 (22.88 mm), A12-2 (22.54 mm), C3 (22.32 mm) and C2 (22.21 mm), respectively. The desired shape of the edible seed pumpkin was the seed in the elliptic form and narrow elliptic seeds were also evaluated as appetizers. In this study, it is determined that pumpkin group pumpkin seeds had the length/diameter rate as 1.91 [1]. When the weight of 1000 seeds was analyzed, the best value (343.8 g) was obtained from A7 genotype and then A8 (335.6 g), C32 (332.8 g), C16 (332.8 g) and B25 (327.8 g), (Table IV). In the conducted studies, the weight of 1000 seeds was about 200 g [16], [19], [24] and the results we obtained from our genotypes indicate that the seeds were wiper. As a result of the conducted weighted evaluation, the genotypes having the score over 400 were regarded hopeful. When the Table V was analyzed, 13 genotypes; A1, A3, A7, A8, A32, B17, B24, B25, B33, C24, C25, C26 and C30, receiving a score over 400 were found hopeful.

IV. CONCLUSION

The agriculture of edible seed pumpkin seed has been gradually increasing in Turkey in recent years. Although pumpkin seeds are consumed as dried nuts in Turkey and

some countries, it is used in the production of chocolate, candy, sweets and cakes. Moreover, it draws attention with its quality and high oil content. It is also employed in cooking. When the causes such as they do not require much maintenance, they are easy to store, and they do not have marketing problem are added, the charm of growing pumpkin for their seed can be better understood. Although they have such advantages in production, there are also numerous problems in growing edible seed pumpkin which require solution. The biggest problem in the agriculture of edible seed pumpkin is the need for sort or sorts which are one-to-one in terms of high quality, shape, color, massiveness, thickness of its edges and ease of cracking, and registered.

In the present study, 13 genotypes at the stage of S5 such as A1, A3, A7, A8, A32, B17, B24, B25, B33, C24, C25, C26 and C30 were found candidates. According to the available results, those genotypes will be tested for yield and standard types will be developed in order to provide contribution to the economy of the country. Genotypes were brought to the stage of S5 and the opportunity of developing hybrids may be obtained in the following stage.

ACKNOWLEDGMENTS

This research was funded by TUBITAK (TOVAK), Turkey (Project number:110 O 088).

REFERENCES

- [1] H.S. Paris, "History of the cultivar-groups of *Cucurbita pepo*" In: Janick J. (ed.), Horticultural Reviews, 25: 71–170, 2000.
- [2] R. Yanmaz, B. Düzeltir, "Çekirdek kabağı yetiştiriciliği" *Ekin Dergisi*. Yayın No: 26;22-24, 2003.
- [3] L. Keskin, "Çekirdek Kabak Yetiştiriciliği, Sorunları ve Çözüm Önerileri. Yüksek Lisans Semineri, Selçuk Üniversitesi" Fen Bilimleri Enstitüsü, 33 s., Konya, 2007.
- [4] T.J. Jacks, T.P. Hensarling, L.Y. Yatsu, "Cucurbit seeds: I. Characterizations and uses of oil and proteins" A review, *Econ. Bot.* 26:135-141, 1972.
- [5] M. Murkovic, A. Hillebrand, S. Draxl, J. Winkler, W. Pfannhauser, "Distribution of Fatty Acids and Vitamin E Content in Pumpkin Seeds (*Cucurbita Pepo* L.)" In Breeding Lines. *Acta Hort.* 492: 47– 55, 1999.
- [6] J. Winkler, "The Origin and Breeding of the Hull-less Seeded Styrian Oil-Pumpkin Varieties in Austria" *Cucurbit Genetics Cooperative Report* 23:101-104 (article340), 2000.

- [7] A. Gohari Ardabil, R. Farhoosh, M.H. Haddad Khodaparast, "Chemical Composition and Physicochemical Properties of Pumpkin Seeds (*Cucurbita pepo* Subsp. *pepo* Var. *Styriaca*) Grown in Iran" *J. Agr. Sci. Tech.* (2011) Vol. 13: 1053-1063, 2011.
- [8] M. Yegul, M. Yıldız, Ş. Ellialtınoğlu, K. Abak, "Bazı Kabuksuz Çekirdek Kabağı (*Cucurbita pepo* var. *styrica*) Islah Hatlarında Tohum Verimi ve Kalitesi" *YYÜ Tar. Bil. Derg.* 22 (1):12-19, 2012.
- [9] M.B. Achu, E. Fokou, C. Tchiégang, M. Fotso, F.M. Tchouanguep, "Nutritive value of some Cucurbitaceae oilseeds from different regions in Cameroon" *African Journal of Biotechnology*, Vol. 4 (11), pp. 1329-1334, 2005.
- [10] M. Murkovic, A. Hillebrand, J. Winkler, W. Pfannhauser, "Variability of Vitamin E Content in Pumpkin Seeds (*Cucurbita pepo* L.)" *Z Lebensm Unters Forsch*, 202:275-278, 1996.
- [11] T. Wenzl, E. Prettnner, K. Schweiger, F.S. Wagner, "An Improved Method to Discover Adulteration of Styrian Pumpkin Seed Oil" *J. Biochem. Biophys Methods*, 53: 193-202, 2002.
- [12] D. Diaz Obregon, L. Lloja Lozano, V. Carbajal Zuniga, "Preclinical studies of *Cucurbita maxima* (pumpkin seeds) a traditional intestinal antiparasitic in rural urban areas" *Rev. Gastroenterol Peru* 24(4):323-327, 2004.
- [13] K. Koike, W. Li, L. Liu, E. Hata, T. Nikaido, "New phenolic glycosides from the seeds of *Cucurbita moschata*" *Chem Pharm Bull* 53(2):225-228, 2005.
- [14] M. Ferriol, B. Pico, "Pumpkin and Winter Squash" *Handbook of Plant Breeding Volume 1*, pp 317-349, 2008.
- [15] O. Gilbert, A.H. Fruhwirth, "Seeds and oil of the Styrian oil pumpkin: Components and biological activities" *EUR. J. Lipid Sci. Technol*, 109, 1128-1140, 2007.
- [16] G. Toprakkanıştıran, "Çekirdek kabaklarında seleksiyon ıslahı: 1. Döl kademesinin elde edilmesi" *Yüksek Lisans Tezi, Ankara Üniversitesi, Fen Bilimleri Enstitüsü, Ankara*, 34 s, 1997.
- [17] M. Seymen, Ö. Türkmen, M. Paksoy, "Selection of edible pumpkin seeds (*Cucurbita pepo* L.) genotypes" *Journal of Selçuk University Natural and Applied Science*. 2(4): pp-29-39, 2011.
- [18] M. Seymen, Ö. Türkmen, M. Paksoy, S. Fidan, "Determination of Some morphological characteristics of edible seed pumpkin (*Cucurbita pepo* L.) Genotypes" *Xth EUCARPIA International Meeting on Cucurbitaceae*, pp 739-749 October 15-18, 2012, Antalya-Turkey, 2012.
- [19] B. Düzeltilir, "Çekirdek kabağı (*Cucurbita pepo* L.) hatlarında morfolojik özelliklerine göre tanımlanma ve seleksiyon çalışmaları" *Yüksek Lisans Tezi, Ankara Üniversitesi Fen Bilimleri Enstitüsü, Ankara*, 76 s, 2004.
- [20] M. Seymen, Ö. Türkmen, M. Paksoy, S. Fidan, "Determination of Some morphological characteristics of edible seed pumpkin (*Cucurbita pepo* L.) Genotypes" *Xth EUCARPIA International Meeting on Cucurbitaceae*, pp 739-749 October 15-18, 2012, Antalya-Turkey 2012.
- [21] K. Abak, M. Sakin, Ş. Karakullukçu, "Improvement of pumpkin for naked seeds" *XXIII. International Horticultural Congress, Italy*, Abstracts Book II, 3074, 1990.
- [22] W.A. Warid, J.J. Martinex, J.M. Loaiza, "Productivity of naked seed squash, *Cucurbita pepo* L." *Cucurbit Genetics Cooperative Report* 16. 1993.
- [23] M. Murkoviç, J. Winkler, W. Pfannhauser, "Improvement of the quality of pumpkin seed (*Cucurbita pepo* L.) by use of cluster analysis" *First International Symposium on Cucurbits*. Adana- Turkey, Syf: 41-46, 1997.
- [24] Joshi, D.C., Das, S.K., Mukherjee, R.K., 1993, Physical properties of pumpkin seed, *J. Agric. Engng Res*, 54:219-229.