

# Natural Radioactivity in Foods Consumed in Turkey

E. Kam, G. Karahan, H. Ashyuksek, A. Bozkurt

**Abstract**—This study aims to determine the natural radioactivity levels in some foodstuffs produced in Turkey. For this purpose, 48 different foods samples were collected from different land parcels throughout the country. All samples were analyzed to designate both gross alpha and gross beta radioactivities and the radionuclides' concentrations. The gross alpha radioactivities were measured as below  $1 \text{ Bq kg}^{-1}$  in most of the samples, some of them being due to the detection limit of the counting system. The gross beta radioactivity levels ranged from  $1.8 \text{ Bq kg}^{-1}$  to  $453 \text{ Bq kg}^{-1}$ , larger levels being observed in leguminous seeds while the highest level being in haricot bean. The concentrations of natural radionuclides in the foodstuffs were investigated by the method of gamma spectroscopy. High levels of  $^{40}\text{K}$  were measured in all the samples, the highest activities being again in leguminous seeds. Low concentrations of  $^{238}\text{U}$  and  $^{226}\text{Ra}$  were found in some of the samples, which are comparable to the reported results in the literature. Based on the activity concentrations obtained in this study, average annual effective dose equivalents for the radionuclides  $^{226}\text{Ra}$ ,  $^{238}\text{U}$ , and  $^{40}\text{K}$  were calculated as  $77.416 \mu\text{Sv y}^{-1}$ ,  $0.978 \mu\text{Sv y}^{-1}$ , and  $140.55 \mu\text{Sv y}^{-1}$ , respectively.

**Keywords**—Foods, radioactivity, gross alpha, gross beta, annual equivalent dose, Turkey.

## I. INTRODUCTION

OVER the last century, nuclear materials have found extensive medical, industrial and military applications, which in turn brought about a worldwide concern on radiation exposure of human beings. The nuclear weapon tests and radioactive accidents have also triggered the public fear and as a result a considerable amount of research energy was spent over the last several decades in evaluating the radioactivity content of soil, air, and water [1]. Because the terrestrial or cosmic radioisotopes are always found in the ecosystem and can easily find pathways to enter the metabolisms of plants and animals, such naturally found substances continuously expose people to radiation through the food chain. Therefore, in line with routine measurements of background radiation, a substantial amount of research effort is continually invested in determining how much radioactivity is contained within certain foodstuffs that are also part of the human diet.

A radioactive element in food or water is absorbed in bodies of plants and animals by different mechanisms that are typically dependent on its chemical properties rather than its radioactive characteristics [2]. For example, a plant in need of calcium and potassium will take them through its capillary roots from soil and will not discriminate against the radioactive  $^{45}\text{Ca}$  and  $^{40}\text{K}$  isotopes. Again the distribution patterns of uranium, thorium and their decay products are affected by certain chemical and biochemical interactions. The

amount of these primordial radionuclides in food elements will accordingly depend upon the parent rock and the soil formation along with the transport processes that are involved. Finally, anthropogenic radionuclides such as  $^{137}\text{Cs}$  and  $^{90}\text{Sr}$  exist in the atmosphere as a result of nuclear weapon tests and radioactive accidents starting from 1950s. These manmade radioactive materials will behave in a similar manner when accumulate on plant leaves and grass in the course of time after fallouts.

Although there are many studies in the literature that present data about the radioactivity in soil, water, and air in some Turkish cities, data on radioactive contents of produces are scarce. This study attempts to determine the level of radioactivity in some foodstuffs that are commonly consumed by people living in Turkey. For this purpose, a radiologic survey that includes gross alpha and gross beta radioactivity measurements along with concentrations of naturally occurring radionuclides in 48 food samples was carried out. The results were compared with those from other countries.

## II. MATERIALS AND METHODS

### A. Sample Collection and Preparation

The samples of fruits and vegetables which were raised in different land parcels throughout the country were purchased from different sellers and were taken as individual food items rather than mixed diet samples. The measurement samples were first washed with tap water and then with distilled water, and peeled when necessary. The wet and dried masses were recorded by subtracting the edible portion from the total to account for the loss of moisture from the edible part during the time of preparation.

Almost all foods samples were prepared in two different methods: Home use form and ash form. In home use preparation, solid and liquid samples were first air-dried at room temperature after breaking into small sizes. The samples were then put into 1000 mL Marinelli beakers which were previously treated with dilute HCL to prevent any possible contamination. Finally, each tightly sealed container was stored for a period of 1 month to attain a radioactive equilibrium between  $^{226}\text{Ra}$  and its daughters [3]. The second method involved reducing the samples to ash form where one of two different techniques could be used. In wet ashing, the sample is treated using oxidant nitric acid ( $\text{HNO}_3$ ) whereas dry ashing method is much simpler, takes less time, and is more suitable for radionuclides that do not vaporize at ashing temperature (since some radionuclides in the sample may be lost at high temperatures). In this study, dry ashing method was adopted for all foodstuffs except liquid samples such as water and olive oil that were evaporated at  $55^\circ\text{C}$ . For each dried food crop, a mass of 100 g was taken to ash in high

Erol Kam is with the Science Faculty, Yıldız Technical University, Esenler, Istanbul, Turkey (e-mail: erolkam@hotmail.com).

temperature resistant porcelain plates using an electric furnace. The temperature for dry ashing varied where an upper limit was generally around 450 °C [4]. The ashing time, on the other hand, depended on the type and quantity of the material where it was more than a day for some of the samples. Measurements of the ashed weight were necessary for calculations of radioactivity and radionuclide concentrations and yield. At the end of this process, the samples in ash form were homogenized, weighed and tied up in 100 mL polyethylene beakers, and finally stored for a period of 1 month to achieve an approximate secular equilibrium between  $^{226}\text{Ra}$  and its daughters before performing the radioisotope analyses.

### B. Radioactivity Measurements

Gross alpha and gross beta radioactivity measurements were performed by Krieger method [5] using a proportional gas flow counter (Ortec Telenet Systems). All food samples were counted in counter systems after incineration. The beta counting system had a low background detector with gas-flow window type approximately 5 cm in diameter. The gas for each detector was a mixture of 90% argon and 10% methane. All samples were placed in a 5 cm diameter stainless steel planchette for counting. The alpha counting system was calibrated using a  $^{238}\text{U}$  standard source while the beta system was calibrated with a  $^{40}\text{K}$  standard source. The counting time was 1000 minutes for gross alpha measurements and 100 minutes for gross beta measurements.

Gamma spectrometric measurements were made with a coaxial high purity germanium detector (Canberra GC 1520 model) having 16 % relative efficiency and 1.9 keV energy resolution (FWHM) at 1332 keV gamma transition of  $^{60}\text{Co}$ . To ensure accurate quantitative measurements, the detector was calibrated by two different certified gamma ray standard sources in 1000 mL Marinelli beaker for home use form and mixed radionuclides in 100 mL beaker for ashed form, the same size and type used to count the samples. The counting time for each sample was 50,000 s which produced sufficiently strong peaks and small counting errors. The uncertainties in all measurements were about  $2\sigma$ . The background spectra were also measured under the same conditions used for the reference materials and the samples.

Spectrometric analyses were carried out for the natural radionuclides  $^{226}\text{Ra}$ ,  $^{238}\text{U}$  and  $^{40}\text{K}$ . The activity of each sample was determined using the total net counts under the selected photo-peaks (186 keV for  $^{226}\text{Ra}$ , 1461 keV for  $^{40}\text{K}$ ), the measured photo-peak efficiency, the gamma intensity and the sample weight. The activity concentrations of these radionuclides were obtained for each sample (in units of Bq  $\text{kg}^{-1}$ ) after correcting for background and Compton contributions.

TABLE I A  
GROSS ALPHA AND GROSS BETA RADIOACTIVITIES (BQ  $\text{KG}^{-1}$ ) IN REFINED PRODUCTS

| Food                            | Gross $\alpha$    | Gross $\beta$    |
|---------------------------------|-------------------|------------------|
| Flour                           | $0.352 \pm 0.080$ | $92.22 \pm 3.42$ |
| Olive oil (Bq $\text{l}^{-1}$ ) | ND                | $1.80 \pm 0.02$  |
| Sugar                           | ND*               | $10.31 \pm 0.24$ |

\*not determined

TABLE I B  
GROSS ALPHA AND GROSS BETA RADIOACTIVITIES (BQ  $\text{KG}^{-1}$ ) IN ANIMAL PRODUCTS

| Food                       | Gross $\alpha$    | Gross $\beta$    |
|----------------------------|-------------------|------------------|
| Beef                       | $0.130 \pm 0.033$ | $58.11 \pm 1.74$ |
| Chicken                    | $0.035 \pm 0.001$ | $84.30 \pm 3.08$ |
| Egg                        | $0.008 \pm 0.002$ | $48.12 \pm 2.56$ |
| Fish                       | ND                | $69.50 \pm 1.08$ |
| Milk (Bq $\text{l}^{-1}$ ) | ND                | $43.67 \pm 1.82$ |

TABLE I C  
GROSS ALPHA AND GROSS BETA RADIOACTIVITIES (BQ  $\text{KG}^{-1}$ ) IN FRUITS

| Food       | Gross $\alpha$    | Gross $\beta$    |
|------------|-------------------|------------------|
| Apple      | $0.590 \pm 0.113$ | $25.41 \pm 1.36$ |
| Apricot    | $0.091 \pm 0.002$ | $80.29 \pm 2.56$ |
| Cherry     | $0.074 \pm 0.004$ | $36.91 \pm 3.12$ |
| Grapes     | ND                | $61.79 \pm 1.45$ |
| Olive      | $0.400 \pm 0.082$ | $49.23 \pm 2.87$ |
| Orange     | ND                | $38.69 \pm 1.72$ |
| Peach      | $0.172 \pm 0.060$ | $53.76 \pm 2.77$ |
| Pear       | $0.082 \pm 0.016$ | $29.91 \pm 1.34$ |
| Plum       | $0.363 \pm 0.043$ | $32.44 \pm 1.29$ |
| Strawberry | $0.353 \pm 0.090$ | $32.41 \pm 2.19$ |
| Tangerine  | ND                | $32.02 \pm 1.38$ |
| Watermelon | ND                | $39.12 \pm 2.23$ |

TABLE I D  
GROSS ALPHA AND GROSS BETA RADIOACTIVITIES (BQ  $\text{KG}^{-1}$ ) IN VEGETABLES

| Food            | Gross $\alpha$    | Gross $\beta$     |
|-----------------|-------------------|-------------------|
| Aubergine       | $0.202 \pm 0.035$ | $36.40 \pm 1.89$  |
| Cabbage (dark)  | $0.401 \pm 0.120$ | $75.80 \pm 3.00$  |
| Cabbage (red)   | $0.130 \pm 0.042$ | $68.10 \pm 2.37$  |
| Cabbage (white) | $0.262 \pm 0.067$ | $48.33 \pm 1.92$  |
| Carrot          | $0.442 \pm 0.103$ | $56.29 \pm 2.16$  |
| Cauliflower     | $0.242 \pm 0.074$ | $59.23 \pm 2.28$  |
| Celery          | $0.520 \pm 0.140$ | $70.61 \pm 2.39$  |
| Chard           | ND                | $49.43 \pm 2.82$  |
| Cucumber        | $0.120 \pm 0.041$ | $39.24 \pm 1.67$  |
| Curly           | ND                | $34.70 \pm 3.59$  |
| Green beans     | $0.112 \pm 0.033$ | $61.52 \pm 2.25$  |
| Green pepper    | $0.091 \pm 0.020$ | $43.10 \pm 1.76$  |
| Leek            | $0.150 \pm 0.041$ | $58.33 \pm 2.16$  |
| Marrow          | $0.213 \pm 0.022$ | $50.89 \pm 1.92$  |
| Onion           | $0.262 \pm 0.071$ | $57.02 \pm 2.06$  |
| Pea             | $0.093 \pm 0.002$ | $35.36 \pm 2.51$  |
| Potato          | $0.162 \pm 0.048$ | $74.41 \pm 3.36$  |
| Purslane        | $0.530 \pm 0.122$ | $101.25 \pm 2.82$ |
| Red radish      | $0.123 \pm 0.045$ | $47.65 \pm 1.77$  |
| Romaine lettuce | $0.133 \pm 0.041$ | $79.63 \pm 2.60$  |
| Spinach         | $0.062 \pm 0.011$ | $100.53 \pm 3.08$ |
| Tomato          | $0.173 \pm 0.053$ | $37.39 \pm 1.60$  |

TABLE I E  
GROSS ALPHA AND GROSS BETA RADIOACTIVITIES (BQ KG<sup>-1</sup>) IN LEGUMINOUS PLANTS

| Food           | Gross $\alpha$    | Gross $\beta$      |
|----------------|-------------------|--------------------|
| Chickpea       | 0.470 $\pm$ 0.009 | 197.02 $\pm$ 5.72  |
| Haricot beans  | 0.471 $\pm$ 0.102 | 453.02 $\pm$ 15.64 |
| Lentil (green) | 0.562 $\pm$ 0.013 | 200.12 $\pm$ 9.12  |
| Lentil (red)   | 0.744 $\pm$ 0.190 | 176.03 $\pm$ 3.36  |
| Rice           | ND                | 167.30 $\pm$ 3.31  |
| Wheat          | 0.772 $\pm$ 0.142 | 77.14 $\pm$ 4.63   |

### III. RESULTS

#### A. Gross Alpha and Gross Beta Radioactivity Measurements

Food samples (48 in total) were first analyzed by gathering into five groups: refined products, animal products, fruits, vegetables and leguminous plants. The gross alpha and gross beta radioactivity results in the studied foodstuffs are given in Tables I A-E. The gross alpha radioactivity varies between 0.008 Bq kg<sup>-1</sup> (egg) and 0.772 Bq kg<sup>-1</sup> (wheat). This quantity was the highest in leguminous plants and could not be determined in some foods such as fish, milk, rice, some fruits and some vegetables. In all the samples investigated, gross beta concentrations were found at higher levels than those of alpha and varied between 1.8 Bq kg<sup>-1</sup> (olive oil) and 453 Bq kg<sup>-1</sup> (haricot beans). Particularly, leguminous plants had very high gross beta radioactivities.

#### B. Radionuclide Activity Concentrations

TABLE II A  
RADIONUCLIDE CONCENTRATIONS (BQ KG<sup>-1</sup>) IN REFINED PRODUCTS

| Food                            | <sup>226</sup> Ra | <sup>238</sup> U | <sup>40</sup> K |
|---------------------------------|-------------------|------------------|-----------------|
| Flour                           | 1.15 $\pm$ 0.11   | 0.21 $\pm$ 0.02  | 51.2 $\pm$ 20   |
| Olive oil (Bq l <sup>-1</sup> ) | ND                | ND               | 1.66 $\pm$ 0.36 |
| Sugar                           | ND                | ND               | 7.80 $\pm$ 1.24 |

TABLE II B  
RADIONUCLIDE CONCENTRATIONS (BQ KG<sup>-1</sup>) IN ANIMAL PRODUCTS

| Food                       | <sup>226</sup> Ra | <sup>238</sup> U | <sup>40</sup> K   |
|----------------------------|-------------------|------------------|-------------------|
| Beef                       | ND                | 0.32 $\pm$ 0.16  | 52.17 $\pm$ 5.70  |
| Chicken                    | ND                | ND               | 74.23 $\pm$ 12.61 |
| Egg                        | ND                | ND               | 46.73 $\pm$ 5.21  |
| Fish                       | ND                | ND               | 67.56 $\pm$ 6.98  |
| Milk (Bq l <sup>-1</sup> ) | ND                | ND               | 42.04 $\pm$ 5.00  |

TABLE II C  
RADIONUCLIDE CONCENTRATIONS (BQ KG<sup>-1</sup>) IN FRUITS

| Food       | <sup>226</sup> Ra | <sup>238</sup> U | <sup>40</sup> K  |
|------------|-------------------|------------------|------------------|
| Apple      | ND                | ND               | 19.23 $\pm$ 3.60 |
| Apricot    | ND                | ND               | 59.25 $\pm$ 8.56 |
| Chery      | ND                | ND               | 28.30 $\pm$ 5.90 |
| Grape      | ND                | ND               | 58.50 $\pm$ 3.33 |
| Olive      | 1.29 $\pm$ 0.40   | ND               | 43.25 $\pm$ 2.81 |
| Orange     | ND                | ND               | 23.70 $\pm$ 2.20 |
| Peach      | ND                | ND               | 51.18 $\pm$ 2.96 |
| Pear       | ND                | ND               | 21.86 $\pm$ 1.85 |
| Plum       | ND                | ND               | 25.70 $\pm$ 5.10 |
| Strawberry | ND                | ND               | 27.87 $\pm$ 2.58 |
| Tangerina  | ND                | ND               | 25.60 $\pm$ 7.32 |
| Watermelon | ND                | ND               | 36.28 $\pm$ 2.90 |

TABLE II D  
RADIONUCLIDE CONCENTRATIONS (BQ KG<sup>-1</sup>) IN VEGETABLES

| Food            | <sup>226</sup> Ra | <sup>238</sup> U | <sup>40</sup> K   |
|-----------------|-------------------|------------------|-------------------|
| Aubergine       | ND                | ND               | 32.90 $\pm$ 4.27  |
| Cabbage (dark)  | ND                | ND               | 70.76 $\pm$ 4.82  |
| Cabbage (red)   | 0.21 $\pm$ 0.11   | ND               | 59.70 $\pm$ 8.03  |
| Cabbage (white) | 2.3 $\pm$ 0.97    | ND               | 42.00 $\pm$ 3.24  |
| Carrot          | ND                | ND               | 43.30 $\pm$ 6.00  |
| Cauliflower     | 2.18 $\pm$ 0.88   | ND               | 55.60 $\pm$ 5.02  |
| Celery          | ND                | ND               | 63.02 $\pm$ 7.80  |
| Chard           | ND                | ND               | 36.70 $\pm$ 11.4  |
| Cucumber        | 1.05 $\pm$ 0.24   | 0.14 $\pm$ 0.05  | 34.54 $\pm$ 6.90  |
| Curly           | 1.35 $\pm$ 0.61   | ND               | 28.09 $\pm$ 6.26  |
| Green beans     | 1.17 $\pm$ 0.86   | ND               | 36.61 $\pm$ 6.22  |
| Green pepper    | 3.95 $\pm$ 1.42   | ND               | 39.87 $\pm$ 6.20  |
| Leek            | 0.26 $\pm$ 0.12   | 0.06 $\pm$ 0.02  | 54.82 $\pm$ 980   |
| Marrow          | ND                | ND               | 37.70 $\pm$ 13.20 |
| Onion           | 0.48 $\pm$ 0.25   | ND               | 53.90 $\pm$ 5.31  |
| Pea             | ND                | ND               | 31.80 $\pm$ 3.40  |
| Potato          | 0.94 $\pm$ 0.45   | ND               | 61.30 $\pm$ 12.42 |
| Purslane        | ND                | ND               | 99.65 $\pm$ 11.56 |
| Red radish      | 1.63 $\pm$ 0.62   | ND               | 42.70 $\pm$ 11.00 |
| Romaine lettuce | 1.44 $\pm$ 0.37   | 0.21 $\pm$ 0.04  | 75.30 $\pm$ 7.20  |
| Spinach         | 0.41 $\pm$ 0.20   | ND               | 78.29 $\pm$ 6.42  |
| Tomato          | 0.84 $\pm$ 0.46   | ND               | 32.40 $\pm$ 4.90  |

TABLE II E  
RADIONUCLIDE CONCENTRATIONS (BQ KG<sup>-1</sup>) IN LEGUMINOUS PLANTS

| Food           | <sup>226</sup> Ra | <sup>238</sup> U | <sup>40</sup> K    |
|----------------|-------------------|------------------|--------------------|
| Chickpea       | ND                | 0.58 $\pm$ 0.12  | 177.80 $\pm$ 17.70 |
| Haricot beans  | ND                | ND               | 374.33 $\pm$ 12.56 |
| Lentil (green) | ND                | 0.36 $\pm$ 0.03  | 183.92 $\pm$ 19.20 |
| Lentil (red)   | ND                | ND               | 165.21 $\pm$ 10.61 |
| Rice           | ND                | ND               | 129.50 $\pm$ 12.33 |
| Wheat          | 1.02 $\pm$ 0.52   | 0.68 $\pm$ 0.11  | 67.03 $\pm$ 8.90   |

The amount of <sup>226</sup>Ra, <sup>238</sup>U, and <sup>40</sup>K isotopes were determined in the foodstuffs collected. The activity concentrations of these radionuclides are given in Tables II A-E. As mentioned earlier, the samples prepared in both home use and ashed forms were counted by gamma spectroscopy. Since the measurements from each method were found to be very close, the results were reported in Tables II A-E as the average of both methods.

Because naturally found <sup>40</sup>K emits both gamma and beta rays, the gamma activity concentration of <sup>40</sup>K were found very close to the gross beta activity in all samples. The highest <sup>40</sup>K radioactivity concentrations were measured in leguminous plants as is the case in gross beta measurements. The maximum gamma radioactivity was determined as 374.32 Bq kg<sup>-1</sup> in haricot beans. The smallest <sup>40</sup>K concentration was found as 1.66 Bq kg<sup>-1</sup> in olive oil.

<sup>226</sup>Ra were found only in some of the foodstuffs included in the study, most of them being vegetables. 32 of the 48 samples did not reveal any <sup>226</sup>Ra activity. The maximum activity concentration for the radionuclide was found as 3.95 Bq kg<sup>-1</sup> in green pepper. Only a limited number of samples showed <sup>238</sup>U activities, most of them being leguminous plants or

vegetables. The highest activity concentration for this radioisotope was measured as 0.68 Bq kg<sup>-1</sup> in wheat.

#### IV. DISCUSSION

The radioactivity contents of various foodstuff have been extensively studied in different parts of the world. Table III provides the specific activities of the foods measured in some of these studies along with findings of this work [6-13]. As seen, our results agree pretty well with the literature data.

For all foodstuffs, a close relationship exists between gamma activity concentration of <sup>40</sup>K and gross beta radioactivity. Table IV provides <sup>40</sup>K/T<sub>β</sub> ratios for the samples investigated. The average activity of <sup>40</sup>K and T<sub>β</sub> in foods are 72.36 Bq kg<sup>-1</sup> and 61.99 Bq kg<sup>-1</sup>, respectively. The average <sup>40</sup>K/T<sub>β</sub> ratio is about 0.857 from which one can conclude that a substantial portion of gross beta radioactivity in foods emanate from the natural <sup>40</sup>K isotope. The average concentration of <sup>40</sup>K in adults is 60 Bq kg<sup>-1</sup> [14].

Potassium is a significant element for human health. Like other minerals, amount of potassium in body varies depending on the age of person where 0.012% of natural potassium is <sup>40</sup>K. Any radionuclide that enters the body by means of foods or beverages continuously exposes it in proportion with the element's biological half-life. Therefore, it is important to calculate the annual effective dose equivalent resulting from exposure to natural radionuclides such as <sup>40</sup>K, <sup>226</sup>Ra and <sup>238</sup>U in the foodstuffs that are part of a person's diet.

The annual effective dose equivalent from the radioactive contents in a daily consumption of a mixture of foods can be easily calculated despite the habits of consumption by people may differ in real life. A typical amount of average daily consumption for every foodstuff included in this study is also given in Table IV. These figures were estimated from annual consumption of an ordinary family because there are no such statistical data for Turkish people. The intake amounts for fish, chicken and flour were taken from the corresponding WHO reports [15]. In total, a person was assumed to consume a daily mix of 1274 g of all these foods and as a result takes 276.5 Bq, 21.7 Bq, and 22668.7 Bq of annual total activity from <sup>226</sup>Ra, <sup>238</sup>U, and <sup>40</sup>K, respectively. Using the corresponding dose

conversion factors for these radionuclides (<sup>226</sup>Ra: 2.8\*10<sup>-7</sup> Sv Bq<sup>-1</sup>; <sup>238</sup>U: 4.5\*10<sup>-8</sup> Sv Bq<sup>-1</sup>; <sup>40</sup>K: 6.2\*10<sup>-9</sup> Sv Bq<sup>-1</sup> [16], the annual effective dose equivalents (AEDE) for <sup>226</sup>Ra, <sup>238</sup>U, and <sup>40</sup>K as 77.416 μSv y<sup>-1</sup>, 0.978 μSv y<sup>-1</sup>, and 140.55 μSv y<sup>-1</sup>, respectively from the data obtained in this study.

TABLE III  
RADIONUCLIDE CONCENTRATIONS MEASURED IN DIFFERENT COUNTRIES\*  
FOODSTUFFS

| Food                       | Country       | <sup>226</sup> Ra (Bq kg <sup>-1</sup> ) | <sup>238</sup> U (Bq kg <sup>-1</sup> ) | <sup>40</sup> K (Bq kg <sup>-1</sup> ) |
|----------------------------|---------------|--|---|--|
| Beef                       | Brasil [6]    |  | 0.5                                     |  |
|                            | Germany [7]   | 0.11                                     |   |  |
| Chicken                    | Turkey        | ND                                       | 0.32                                    | 52.17                                  |
|                            | Brasil [6]    | 0.04-0.16                                |   |  |
| Cucumber                   | Turkey        | ND                                       | ND                                      | 74.23                                  |
|                            | Egypt [8]     |  |   | 43-68                                  |
|                            | India [12]    | 0.097                                    |   | 29.67                                  |
| Milk (Bq l <sup>-1</sup> ) | Turkey        | 1.05                                     | 0.14                                    | 34.54                                  |
|                            | Brasil [6]    | 0.029-0.21                               |   |  |
|                            | Egypt [8]     |  |   | 79                                     |
|                            | India [12]    | 2.5                                      |   | 34.35                                  |
| Onion                      | Jordan [13]   |  |   | 40.8                                   |
|                            | Turkey        | ND                                       | ND                                      | 42.04                                  |
|                            | Egypt [8]     |  |   | 44-68                                  |
|                            | Turkey        | 0.48                                     | ND                                      | 53.90                                  |
| Potato                     | Germany [7]   |  |   | 130-180                                |
|                            | Egypt [8]     |  |   | 76-124                                 |
| Rice                       | Turkey        | 0.94                                     | ND                                      | 61.3                                   |
|                            | China [9]     | 0.07-0.23                                |   |  |
|                            | Germany [7]   |  |   | 90-110                                 |
| Tomato                     | India [12]    | 3.07                                     |   | 120.2                                  |
|                            | Turkey        | ND                                       | ND                                      | 129.54                                 |
|                            | Egypt [8]     |  |   | 39-84                                  |
|                            | India [12]    | 0.06                                     |   | 71.92                                  |
| Wheat                      | Turkey        | 0.84                                     | ND                                      | 32.4                                   |
|                            | Germany [7]   |  |   | 97-130                                 |
|                            | India [10]    | >0.5                                     |   |  |
|                            | Pakistan [11] | 0.98                                     |   | 109.27                                 |
|                            | Turkey        | 1.02                                     | 0.68                                    | 67                                     |

TABLE IV  
<sup>40</sup>K/T<sub>β</sub> RATIOS AND YEARLY CONSUMPTION AMOUNTS IN FOODS AND ANNUAL EFFECTIVE DOSE EQUIVALENT (AEDE)

| Food                            | T <sub>β</sub><br>(Bq kg <sup>-1</sup> ) | <sup>40</sup> K<br>(Bq kg <sup>-1</sup> ) | <sup>40</sup> K/T <sub>β</sub><br>(Bq kg <sup>-1</sup> ) | Consumption<br>(kg y <sup>-1</sup> ) | AEDE (μSv y <sup>-1</sup> ) |                  |                 |
|---------------------------------|--|---|--|--------------------------------------|-----------------------------|------------------|-----------------|
|                                 |  |   |  |                                      | <sup>226</sup> Ra           | <sup>238</sup> U | <sup>40</sup> K |
| Flour                           | 92.22                                    | 51.20                                     | 0.56   | 73.2                                 | 23.57                       | 0.692            | 23.24           |
| Olive oil (Bq l <sup>-1</sup> ) | 1.80                                     | 1.66                                      | 0.92   | 4.7                                  |                             |                  | 0.05            |
| Sugar                           | 10.31                                    | 7.80                                      | 0.76   | 14.6                                 |                             |                  | 0.71            |
| Beef                            | 58.11                                    | 52.17                                     | 0.90   | 5.1                                  |                             | 0.07             | 1.65            |
| Chicken                         | 84.30                                    | 74.23                                     | 0.88   | 5.1                                  |                             |                  | 2.35            |
| Egg                             | 48.12                                    | 46.73                                     | 0.97   | 9.8                                  |                             |                  | 2.84            |
| Fish                            | 69.50                                    | 67.56                                     | 0.97   | 5.1                                  |                             |                  | 2.14            |
| Milk (Bq l <sup>-1</sup> )      | 43.67                                    | 42.04                                     | 0.96   | 50                                   |                             |                  | 13.03           |
| Apple                           | 25.41                                    | 19.23                                     | 0.76   | 28.8                                 |                             |                  | 3.43            |
| Apricot                         | 80.29                                    | 59.25                                     | 0.74   | 2.9                                  |                             |                  | 1.07            |
| Cherry                          | 36.91                                    | 28.30                                     | 0.77   | 2.9                                  |                             |                  | 0.51            |
| Grape                           | 61.79                                    | 58.50                                     | 0.95   | 8.1                                  |                             |                  | 2.94            |
| Olive                           | 49.23                                    | 43.25                                     | 0.88   | 8.9                                  | 3.21                        |                  | 2.39            |

| Food            | $T_{\beta}$<br>(Bq kg <sup>-1</sup> ) | <sup>40</sup> K<br>(Bq kg <sup>-1</sup> ) | <sup>40</sup> K/ $T_{\beta}$<br>(Bq kg <sup>-1</sup> ) | Consumption<br>(kg y <sup>-1</sup> ) | AEDE (μSv y <sup>-1</sup> ) |                  |                 |
|-----------------|---------------------------------------|---|--|--------------------------------------|-----------------------------|------------------|-----------------|
|                 |                                       |   |  |                                      | <sup>226</sup> Ra           | <sup>238</sup> U | <sup>40</sup> K |
| Orange          | 38.69                                 | 23.70                                     | 0.61   | 5.8                                  |                             |                  | 0.85            |
| Peach           | 53.76                                 | 51.18                                     | 0.95   | 5.1                                  |                             |                  | 1.62            |
| Pear            | 29.91                                 | 21.86                                     | 0.73   | 5.1                                  |                             |                  | 0.69            |
| Plum            | 32.44                                 | 25.70                                     | 0.79   | 2.9                                  |                             |                  | 0.46            |
| Strawberry      | 32.41                                 | 27.87                                     | 0.86   | 1.8                                  |                             |                  | 0.31            |
| Tangerine       | 32.02                                 | 25.60                                     | 0.80   | 5.8                                  |                             |                  | 0.92            |
| Watermelon      | 39.12                                 | 36.28                                     | 0.93   | 9.8                                  |                             |                  | 2.20            |
| Aubergine       | 36.40                                 | 32.90                                     | 0.90   | 6.9                                  |                             |                  | 1.41            |
| Cabbage (dark)  | 75.80                                 | 70.76                                     | 0.93   | 2.9                                  |                             |                  | 1.27            |
| Cabbage (red)   | 68.10                                 | 59.70                                     | 0.88   | 2.9                                  | 0.17                        |                  | 1.07            |
| Cabbage (white) | 48.33                                 | 42.00                                     | 0.87   | 5.1                                  | 3.28                        |                  | 1.33            |
| Carrot          | 56.29                                 | 43.30                                     | 0.77   | 8.1                                  |                             |                  | 2.17            |
| Cauliflower     | 59.23                                 | 55.60                                     | 0.94   | 3.7                                  | 2.26                        |                  | 1.28            |
| Celery          | 70.61                                 | 63.02                                     | 0.89   | 1.5                                  |                             |                  | 0.59            |
| Chard           | 49.43                                 | 36.70                                     | 0.74   | 1.5                                  |                             |                  | 0.34            |
| Cucumber        | 39.24                                 | 34.54                                     | 0.88   | 8.1                                  | 2.38                        | 0.05             | 1.73            |
| Curly           | 34.70                                 | 28.09                                     | 0.81   | 3.7                                  | 1.40                        |                  | 0.64            |
| Green beans     | 61.52                                 | 36.61                                     | 0.60   | 8.1                                  | 2.65                        |                  | 1.84            |
| Green pepper    | 43.10                                 | 39.87                                     | 0.93   | 8.1                                  | 8.96                        |                  | 2.00            |
| Leek            | 58.33                                 | 54.82                                     | 0.94   | 5.1                                  | 0.37                        | 0.01             | 1.73            |
| Marrow          | 50.89                                 | 37.70                                     | 0.74   | 2.9                                  |                             |                  | 0.68            |
| Onion           | 57.02                                 | 53.90                                     | 0.95   | 6.9                                  | 0.93                        |                  | 2.31            |
| Pea             | 35.36                                 | 31.80                                     | 0.90   | 2.9                                  |                             |                  | 0.57            |
| Potato          | 74.41                                 | 61.30                                     | 0.82   | 50.6                                 | 13.32                       |                  | 19.23           |
| Purslane        | 101.25                                | 99.65                                     | 0.98   | 1.8                                  |                             |                  | 1.11            |
| Red radish      | 47.65                                 | 42.70                                     | 0.90   | 2.9                                  | 1.32                        |                  | 0.77            |
| Romaine lettuce | 79.63                                 | 75.30                                     | 0.95   | 1.8                                  | 0.73                        | 0.02             | 0.84            |
| Spinach         | 100.53                                | 78.29                                     | 0.78   | 5.1                                  | 0.59                        |                  | 2.48            |
| Tomato          | 37.39                                 | 32.40                                     | 0.87   | 50                                   | 11.76                       |                  | 10.04           |
| Chickpea        | 197.02                                | 177.80                                    | 0.90   | 1.8                                  |                             | 0.05             | 1.98            |
| Haricot beans   | 453.02                                | 374.33                                    | 0.83   | 3.7                                  |                             |                  | 8.59            |
| Lentil (green)  | 200.12                                | 183.92                                    | 0.92   | 1.8                                  |                             | 0.03             | 2.05            |
| Lentil (red)    | 176.03                                | 165.21                                    | 0.94   | 1.8                                  |                             |                  | 1.84            |
| Rice            | 167.30                                | 129.50                                    | 0.77   | 8.1                                  |                             |                  | 6.51            |
| Wheat           | 77.14                                 | 67.03                                     | 0.87   | 1.8                                  | 0.51                        | 0.06             | 0.75            |
| Average         | 72.414                                | 61.934                                    | 0.851  |                                      |                             |                  |                 |
| Total           |                                       |   |  | 1.274                                | 1.489                       | 0.019            | 2.703           |

## REFERENCES

- [1] UNSCEAR (2008). Effects of ionizing radiation. United Nations Scientific Committee on the Effects of Atomic Radiation, 2006 Report to the General Assembly, with scientific annexes. United Nations, New York
- [2] Carini F (2001). Radionuclide transfer from soil to fruit. *Journal of Environmental Radioactivity* 52:237-279.
- [3] Karahan G, Bayulken A (2000). Assessment of gamma dose rates around Istanbul, Turkey. *Journal of Environmental Radioactivity* 47:213-221.
- [4] IAEA (1989). Measurement of radionuclides in food and the environment. International Atomic Energy Agency. Technical Reports Series No. 295, Vienna.
- [5] Krieger LH (1975). Interim radiochemical methodology for drinking water. US Environmental Protection Agency, EPA 600/4, 75-008. Cincinnati, Ohio.
- [6] Venturini L, Sordi GAA (1999). Radioactivity in and committed effective dose from some Brazilian foodstuffs. *Health Physics* 76(3):311-313.
- [7] Rassow J (1986). Kernreaktorunfall in Tschernobyl. Risiken der Kernenergie, VCH Publishers, pp. 21-38.
- [8] Badran HM, Sharshar T, Elnimer T (2003). Levels of <sup>137</sup>Cs and <sup>40</sup>K in edible parts of some vegetables consumed in Egypt. *Journal of Environmental Radioactivity* 67:181-190.
- [9] Zhu H (1993). Investigation of food radioactivity and estimation of external dose by ingestion in two Chinese high radiation areas. In M. Sohrabi, J. U. Ahmed, S. A. & Durrani (Eds.), in *Proceedings of the International Conference on High Levels of Natural Radiation*, Rasmar, 1990. IAEA, Vienna.
- [10] Lalit BY, Shukla VK (1982). Natural radioactivity in foodstuffs from high natural radioactivity areas of Southern India. Vohra KG, Mishra UC, Pillai KC, Sadasivan S (Eds.). *Second Special Symposium on Natural Radiation Environment*, held at Bhabha Atomic Research Centre, Bombay 400 085, India, January 19-23, 1981. Halsted Press, pp. 43-49.
- [11] Akhtar N, Tufail M (2007). Natural radioactivity intake into wheat grown on fertilized farms in two districts of Pakistan. *Radiation Protection Dosimetry* 123(1):103-112.
- [12] Shanthi G, Thampi Thanka Kumaran J, Allan Gnana Raj G, Maniyan CG (2010). Natural radionuclides in the South Indian foods and their annual dose. *Nuclear Instruments and Methods in Physics Research A* 619:436-440.
- [13] Ababneh ZQ, Alyassin AM, Aljarrah KM, Ababneh AM (2010). Measurement of natural and artificial radioactivity in powdered milk consumed In Jordan and estimates of the corresponding annual effective dose. *Radiation Protection Dosimetry*, 138(3): 278-283.
- [14] UNSCEAR (1982). Ionizing radiation: sources and biological effects. United Nations Scientific Committee on the Effects of Atomic

Radiation, 1982 Report to the General Assembly, with annexes. United Nations, New York.

- [15] NCRP (1987). Exposure of the population of the United States and Canada from natural radiation. National Council on Radiation Protection and Measurements, Report No. 94. Bethesda, Maryland.
- [16] ICRP (1996). Age dependent doses to members of the public intake of radionuclides: Part 5, Compilation of ingestion and inhalation dose coefficients. International Commission on Radiation Protection, Publication 68. Pergamon Press, Oxford.