# A Hamiltonian Decomposition of 5-star

Walter Hussak and Heiko Schröder

Abstract—Star graphs are Cayley graphs of symmetric groups of permutations, with transpositions as the generating sets. A star graph is a preferred interconnection network topology to a hypercube for its ability to connect a greater number of nodes with lower degree. However, an attractive property of the hypercube is that it has a Hamiltonian decomposition, i.e. its edges can be partitioned into disjoint Hamiltonian cycles, and therefore a simple routing can be found in the case of an edge failure. The existence of Hamiltonian cycles in Cayley graphs has been known for some time. So far, there are no published results on the much stronger condition of the existence of Hamiltonian decompositions. In this paper, we give a construction of a Hamiltonian decomposition of the star graph 5-star of degree 4, by defining an automorphism for 5-star and a Hamiltonian cycle which is edge-disjoint with its image under the automorphism.

Keywords—interconnection networks; paths and cycles; graphs and groups.

#### I. INTRODUCTION

N ETWORKED computer systems have basic requirements such as fast communication and fault tolerance, which are met by an appropriate choice of interconnection topology. The star graph has been proposed as an interconnection network topology that is better than the hypercube for its ability to connect a greater number of nodes with lower degree [1]. On the other hand, an attractive property of the hypercube is that its edges can be partitioned into disjoint Hamiltonian cycles [2]. The presence of edge-disjoint Hamiltonian cycles is desirable for interconnection networks for various reasons. Fault tolerance is easier to achieve as a simple routing can be found in the case of an edge failure. Efficiency can also be improved. An example is the case of all-to-all broadcasting in multiport systems, where a node can send to or receive from all its neighbours in unit time, as messages can be broken down into smaller messages and sent along edge-disjoint Hamiltonian cycles. Edge-disjoint Hamiltonian cycles have been investigated in various interconnection topologies, for example in deBruijn networks [3] and tori [4]. They have also been studied in star graphs and lower bounds for the number of pairwise edge-disjoint Hamiltonian cycles have been given in [5]. However, there has been no significant progress on the optimum case of edge-disjoint cycles, where all the edges in the network topology are partitioned into Hamiltonian cycles, beyond the case of the hypercube.

Star graphs are Cayley graphs of symmetric groups of permutations of finitely many elements and certain restricted sets of transpositions as the generating sets. Properties such as

Walter Hussak is a lecturer in the Department of Computer Science, Loughborough University, Loughborough, UK e-mail: W.Hussak@lboro.ac.uk.

Heiko Schröder is a Professor in the School of Computer Science and Information Technology, RMIT University, Melbourne, Victoria, Australia email: hos@cs.rmit.edu.au. the existence of Hamiltonian paths and cycles [6], Hamiltonian laceability [7] and indeed Hamiltonian decomposability [8], [9] have been considered for various classes of Cayley graphs. To date, it is known that a Cayley graph over a symmetric group and any generating set of transpositions has a Hamiltonian cycle [10]. Thus, this early result demonstrates that star graphs of any degree have a Hamiltonian cycle. To our knowledge, there are no published results on Hamiltonian decompositions of star graphs. The Hamiltonian decompositions of Cayley graphs of degree 4 in [8] concern Cayley graphs over abelian groups. In this paper, we give an a construction of a Hamiltonian decomposition of star graph 5-star of degree 4, by defining a graph automorphism on 5star and a Hamiltonian cycle that has an edge-disjoint image under the automorphism. As 5-star is of degree 4, this gives a Hamiltonian decomposition.

#### **II. PRELIMINARIES**

We give the basic definitions of star graphs, Hamiltonian cycles and automorphisms.

Definition 1: The *n*-star graph  $St_n$  is the simple undirected regular graph of degree *n*-1 whose vertices  $V(St_n)$  are sequences of *n* elements  $\{a_1, \ldots, a_n\}$ 

$$V(St_n) = \{a_{\rho(1)} \dots a_{\rho(n)} : \rho \text{ is a permutation of } \{1, \dots, n\}\}$$

and whose edges  $E(St_n)$  correspond to swapping the positions of the first element with one of the other *n*-1 elements, i.e.  $e \in E(St_n)$  is of the form:

$$e = (a_{\rho(1)} \dots a_{\rho(i-1)} a_{\rho(i)} a_{\rho(i+1)} \dots a_{\rho(n)},$$
  
$$a_{\rho(i)} \dots a_{\rho(i-1)} a_{\rho(1)} a_{\rho(i+1)} \dots a_{\rho(n)})$$
(1)

We define the *distance* between two distinct elements to be:

$$\delta(a_i, a_j) = \min\{|i - j|, n - |i - j|\}$$

Clearly  $\delta(a_i, a_j) = \delta(a_j, a_i)$ . The *length* of the edge *e* above,  $\lambda(e)$ , is defined to be  $\delta(a_{\rho(1)}, a_{\rho(i)})$ .

Definition 2: A Hamiltonian cycle in  $St_n$  is a pair of sequences  $(\underline{v}, \underline{e})$  of vertices  $\underline{v} = v_1 \dots v_{n!+1}$  and edges  $\underline{e} = e_1 \dots e_{n!}$  such that:

(i) 
$$e_i = (v_i, v_{i+1}) \in E(St_n) \ (1 \le i \le n!),$$
  
(ii)  $\{v_1, \dots, v_{n!+1}\} = V(St_n),$ 

(iii) 
$$v_1 = v_{n!+1}$$
.

Thus, a Hamiltonian cycle follows a path along edges visiting each vertex exactly once before returning to the first vertex. A *Hamiltonian decomposition* of  $St_{2k+1}$  where  $k \ge 1$  consists of k Hamiltonian cycles that are edge-disjoint, i.e. no two Hamiltonian cycles have a common edge. Definition 3: Let (V, E) be a graph, where V is a set of vertices and  $E \subseteq V \times V$  a set of edges. Then, a mapping  $\Phi: V \to V$  is an *automorphism* iff:

- (i)  $\Phi$  is bijective
- (ii) for all  $v_1, v_2 \in V$ ,
  - $(v_1, v_2) \in E$  implies  $(\Phi(v_1), \Phi(v_2)) \in E$

#### III. AN AUTOMORPHISM

The following lemma gives a basic class of graph automorphisms preserving Hamiltonian cycles.

Lemma 4: Let  $\phi$  :  $\{a_1, \ldots, a_n\} \rightarrow \{a_1, \ldots, a_n\}$  be a bijection. Then:

- (i)  $\Phi: V(St_n) \to V(St_n)$ , given by  $\Phi(a_{\rho(1)} \dots a_{\rho(n)}) = \phi(a_{\rho(1)}) \dots \phi(a_{\rho(n)})$ , is an automorphism of the graph  $St_n$
- (ii) if  $\underline{v}=v_1 \dots v_{n!+1}$ ,  $\underline{e}=(v_1, v_2) \dots (v_{n!}, v_{n!+1})$  and  $(\underline{v}, \underline{e})$ is a Hamiltonian cycle in  $St_n$ , then  $\Phi_H(\underline{v}, \underline{e}) = (\Phi(v_1) \dots \Phi(v_{n!+1}), (\Phi(v_1), \Phi(v_2)) \dots (\Phi(v_{n!}), \Phi(v_{n!+1})))$ is also a Hamiltonian cycle.

**Proof:** We check that  $\Phi$  is an automorphism. If  $v_1 \neq v_2 \in V(St_n)$ , say  $v_1 = (a_{\rho^1(1)} \dots a_{\rho^1(n)})$ ,  $v_2 = (a_{\rho^2(1)} \dots a_{\rho^2(n)})$ , where  $\rho^1(i) \neq \rho^2(i)$  for some  $1 \leq i \leq n$ , then  $\phi(a_{\rho^1(i)}) \neq \phi(a_{\rho^2(i)})$  as  $\phi$  is injective, and so  $\Phi(v_1)$  and  $\Phi(v_2)$  differ on their respective *i*-th elements  $\phi(a_{\rho^1(i)})$  and  $\phi(a_{\rho^2(i)})$ . Thus,  $\Phi$  is injective. It is surjective as, given  $b_1 \dots b_n \in V(St_n)$ , by surjectivity of  $\phi$  we can choose  $a_{\rho(1)}, \dots, a_{\rho(n)}$ such that  $\phi(a_{\rho(1)}) = b_1, \dots, \phi(a_{\rho(n)}) = b_n$  and therefore  $\Phi(\phi(a_{\rho(1)}) \dots \phi(a_{\rho(n)})) = b_1 \dots b_n$ . To show that Definition 3(ii) holds, let  $(v_1, v_2) \in E(St_n)$ . By Definition 1, for some permutation  $\rho$  and  $1 \leq i \leq n!$ ,

$$e = (a_{\rho(1)} \dots a_{\rho(i-1)} a_{\rho(i)} a_{\rho(i+1)} \dots a_{\rho(n)},$$
$$a_{\rho(i)} \dots a_{\rho(i-1)} a_{\rho(1)} a_{\rho(i+1)} \dots a_{\rho(n)})$$

Then,

$$\begin{split} \Phi(v_1) &= \phi(a_{\rho(1)}) ... \phi(a_{\rho(i-1)}) \phi(a_{\rho(i)}) \phi(a_{\rho(i+1)}) ... \phi(a_{\rho(n)}) \\ \Phi(v_2) &= \phi(a_{\rho(i)}) ... \phi(a_{\rho(i-1)}) \phi(a_{\rho(1)}) \phi(a_{\rho(i+1)}) ... \phi(a_{\rho(n)}) \end{split}$$

As  $\phi : \{a_1, \ldots, a_n\} \to \{a_1, \ldots, a_n\}$  is a bijection, there is a permutation  $\sigma$  of  $\{1, \ldots, n\}$  such that:

$$\phi(a_j) = a_{\sigma(j)} \text{ for } 1 \leq j \leq n$$

Therefore,

 $\Phi(v_1) = a_{\sigma\rho(1)} \dots a_{\sigma\rho(i-1)} a_{\sigma\rho(i)} a_{\sigma\rho(i+1)} \dots a_{\sigma\rho(n)}$ 

 $\Phi(v_2) = a_{\sigma\rho(i)} \dots a_{\sigma\rho(i-1)} a_{\sigma\rho(1)} a_{\sigma\rho(i+1)} \dots a_{\sigma\rho(n)}$ 

and so  $(\Phi(v_1), \Phi(v_2))$  satisfies (1) with  $\sigma \rho$  in place of  $\rho$  and thus  $(\Phi(v_1), \Phi(v_2)) \in E$ . This completes the check of (i) of this lemma, that  $\Phi$  is an automorphism.

To prove (ii) of this lemma, we check that (i), (ii) and (iii) of Definition 2 are satisfied. Put  $(\underline{v}', \underline{e}') = \Phi_H(\underline{v}, \underline{e})$  so that:

$$\underline{v}' = \Phi(v_1) \dots \Phi(v_{n!+1})$$
$$\underline{e}' = (\Phi(v_1), \Phi(v_2)) \dots (\Phi(v_{n!}), \Phi(v_{n!+1}))$$

For (i) of Definition 2, let  $(\Phi(v_i), \Phi(v_{i+1})) \in e'$  where  $1 \leq i \leq n!$ . As  $(\underline{v}, \underline{e})$  is a Hamiltonian cycle in  $St_n$ , we have that  $(v_i, v_{i+1})$  is an edge in  $E(St_n)$ . By (i) of this lemma,  $\Phi$  is an automorphism and so  $(\Phi(v_i), \Phi(v_{i+1}))$  is an edge in  $E(St_n)$  as required. For Definition 2(ii), as  $\Phi$  is an automorphism, it maps the set of all vertices  $\{v_1, \ldots, v_{n!+1}\}$  onto itself. Thus  $V(St_n) = \{v_1, \ldots, v_{n!+1}\} = \{\Phi(v_1), \ldots, \Phi(v_{n!+1})\}$ . For Definition 2(ii), we note that as  $(\underline{v}, \underline{e})$  is a Hamiltonian cycle in  $St_n, v_1 = v_{n!+1}$  and therefore  $\Phi(v_1) = \Phi(v_{n!+1})$ .

The automorphism of interest to us for the graph  $St_5$ , denoted  $\Phi_5$ , corresponds to the bijection  $\phi_5$  on the five elements  $a_1 = a$ ,  $a_2 = b$ ,  $a_3 = c$ ,  $a_4 = d$  and  $a_5 = e$  given by:

$$\phi_5(a) = c, \ \phi_5(b) = a, \ \phi_5(c) = d, \ \phi_5(d) = b, \ \phi_5(e) = e$$

An important property of  $\Phi_5$ , that we shall make use of, is that edges are mapped to edges of a different length.

Lemma 5: If  $(v_1, v_2)$  in an edge in  $St_5$ , then  $\lambda(v_1, v_2) \neq \lambda(\Phi_5(v_1), \Phi_5(v_2))$ .

*Proof:* The length  $\lambda(v_1, v_2)$  of an edge  $(v_1, v_2)$  is the distance between the two symbols  $a_i, a_j \in \{a, b, c, d, e\}$  swapped at that edge. The length of the corresponding edge under  $\Phi_5$  is the distance between  $\phi_5(a_i)$  and  $\phi_5(a_j)$ . There are 10 pairs of symbols  $\{a_i, a_j\}$  to consider:

(i)  $\delta(a, b) = 1$ ,  $\delta(\phi_5(a), \phi_5(b)) = \delta(c, a) = 2$ , (ii)  $\delta(a, c) = 2$ ,  $\delta(\phi_5(a), \phi_5(c)) = \delta(c, d) = 1$ , (iii)  $\delta(a, d) = 2$ ,  $\delta(\phi_5(a), \phi_5(d)) = \delta(c, b) = 1$ , (iv)  $\delta(a, e) = 1$ ,  $\delta(\phi_5(a), \phi_5(e)) = \delta(c, e) = 2$ , (v)  $\delta(b, c) = 1$ ,  $\delta(\phi_5(b), \phi_5(c)) = \delta(a, d) = 2$ , (vi)  $\delta(b, d) = 2$ ,  $\delta(\phi_5(b), \phi_5(d)) = \delta(a, b) = 1$ , (vii)  $\delta(b, e) = 2$ ,  $\delta(\phi_5(b), \phi_5(e)) = \delta(a, e) = 1$ , (viii)  $\delta(c, d) = 1$ ,  $\delta(\phi_5(c), \phi_5(d)) = \delta(d, b) = 2$ , (ix)  $\delta(c, e) = 2$ ,  $\delta(\phi_5(c), \phi_5(e)) = \delta(d, e) = 1$ , (x)  $\delta(d, e) = 1$ ,  $\delta(\phi_5(d), \phi_5(e)) = \delta(b, e) = 2$ .

#### IV. CONSTRUCTION OF A HAMILTONIAN CYCLE

A Hamiltonian cycle for  $St_5$  is constructed by partitioning the vertices of  $St_5$  into 6 pairwise disjoint cycles  $C_1, \ldots, C_6$ , and then producing a 7th cycle  $C_7$  that meets each of the other cycles at exactly two vertices and a common edge. It is clear that the union of the edges in the 7 cycles, excluding edges that  $C_7$  has in common with any of the other 6 cycles, is then a Hamiltonian cycle; we denote it by C. Below, in Lemma 7, we define a cycle  $C_1$  from which 5 further cycles  $C_2$ ,  $C_3$ ,  $C_4$ ,  $C_5$  and  $C_6$  are generated by the 5 length-preserving automorphisms of the following lemma:

Lemma 6: The 5 maps given by:

$$\begin{split} \Psi_2(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}) &= \\ a_{\rho(1)}a_{\rho(2)}a_{\rho(5)}a_{\rho(3)}a_{\rho(4)} \\ \Psi_3(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}) &= \\ a_{\rho(1)}a_{\rho(2)}a_{\rho(4)}a_{\rho(5)}a_{\rho(3)} \\ \Psi_4(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}) &= \\ a_{\rho(1)}a_{\rho(2)}a_{\rho(5)}a_{\rho(4)}a_{\rho(3)} \end{split}$$

$$\begin{split} \Psi_5(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}) &= \\ a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(5)}a_{\rho(4)} \\ \Psi_6(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}) &= \\ a_{\rho(1)}a_{\rho(2)}a_{\rho(4)}a_{\rho(3)}a_{\rho(5)} \end{split}$$

where  $a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}$  is any given point in  $St_5$  with corresponding permutation  $\rho$ , are automorphisms of  $St_5$  which preserve cycles and lengths of edges, i.e.

$$\lambda(v_1, v_2) = \lambda(\Psi_i(v_1), \Psi_i(v_2)) \ (v_1, v_2 \in V(St_5), \ 2 \le i \le 6)$$

(The 5 maps correspond to the 5 possible alternative orders of the last 3 positions in a vertex.)

*Proof:* We check that the lemma holds for  $\Psi_3$  - a very similar check can be performed for  $\Psi_2$ ,  $\Psi_4$ ,  $\Psi_5$  and  $\Psi_6$ . Suppose that  $a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}$  and  $a_{\rho'(1)}a_{\rho'(2)}a_{\rho'(3)}a_{\rho'(4)}a_{\rho'(5)} \in St_5$  differ, i.e.  $\rho$  and  $\rho'$  differ. Then, clearly,

$$\Psi_{3}(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}) =$$

$$a_{\rho(1)}a_{\rho(2)}a_{\rho(4)}a_{\rho(5)}a_{\rho(3)} \neq$$

$$a_{\rho'(1)}a_{\rho'(2)}a_{\rho'(4)}a_{\rho'(5)}a_{\rho'(3)} =$$

$$\Psi_{3}(a_{\rho'(1)}a_{\rho'(2)}a_{\rho'(3)}a_{\rho'(4)}a_{\rho'(5)})$$

Thus,  $\Psi_3$  is injective. If  $a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}$  is any vertex in  $St_n$ , then  $\Psi_3(a_{\rho(1)}a_{\rho(2)}a_{\rho(5)}a_{\rho(3)}a_{\rho(4)}) = a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}$ . Thus,  $\Psi_3$  is surjective. To show that  $\Psi_3$  is an automorphism, suppose that  $v_1 = a_{\rho(1)} \dots a_{\rho(i)} \dots a_{\rho(i)} \dots a_{\nu(i)}$ ,  $v_2 = a_{\rho(i)} \dots a_{\rho(1)} \dots$ , so that  $(v_1, v_2) \in E(St_5)$ , where i = 2, 3, 4 or 5. In the case i = 2,

 $(v_1, v_2) =$ 

 $(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}, a_{\rho(2)}a_{\rho(1)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}),$ 

$$(\Psi_3(v_1), \Psi_3(v_2)) =$$

 $(a_{\rho(1)}a_{\rho(2)}a_{\rho(4)}a_{\rho(5)}a_{\rho(3)}, a_{\rho(2)}a_{\rho(1)}a_{\rho(4)}a_{\rho(5)}a_{\rho(3)})$ 

$$\in E(St_5)$$

In the case i = 3,

 $(v_1, v_2) =$ 

 $(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}, a_{\rho(3)}a_{\rho(2)}a_{\rho(1)}a_{\rho(4)}a_{\rho(5)}),$ 

$$(\Psi_3(v_1), \Psi_3(v_2)) =$$

 $(a_{\rho(1)}a_{\rho(2)}a_{\rho(4)}a_{\rho(5)}a_{\rho(3)}, a_{\rho(3)}a_{\rho(2)}a_{\rho(4)}a_{\rho(5)}a_{\rho(1)})$ 

$$\in E(St_5)$$

In the case i = 4,

 $(v_1, v_2) =$ 

 $(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}, a_{\rho(4)}a_{\rho(2)}a_{\rho(3)}a_{\rho(1)}a_{\rho(5)}),$ 

$$(\Psi_3(v_1), \Psi_3(v_2)) =$$

 $(a_{\rho(1)}a_{\rho(2)}a_{\rho(4)}a_{\rho(5)}a_{\rho(3)}, a_{\rho(4)}a_{\rho(2)}a_{\rho(1)}a_{\rho(5)}a_{\rho(3)})$  $\in E(St_5)$ 

In the case i = 5,

$$(v_1, v_2) =$$

 $(a_{\rho(1)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(5)}, a_{\rho(5)}a_{\rho(2)}a_{\rho(3)}a_{\rho(4)}a_{\rho(1)}),$ 

 $(\Psi_3(v_1), \Psi_3(v_2)) =$ 

 $(a_{\rho(1)}a_{\rho(2)}a_{\rho(4)}a_{\rho(5)}a_{\rho(3)}, a_{\rho(5)}a_{\rho(2)}a_{\rho(4)}a_{\rho(1)}a_{\rho(3)})$ 

 $\in E(St_5)$ 

Thus,  $\Psi_3$  is an automorphism and therefore also preserves cycles by an argument similar to that in Lemma 4(ii). We note from the above cases that, given an edge of the form:

$$(a_{\rho(1)}\ldots a_{\rho(i)}\ldots a_{\rho(i)}\ldots a_{\rho(1)}\ldots) \in E(St_5), \quad (2)$$

we have that

$$(\Psi_3(a_{\rho(1)}\ldots a_{\rho(i)}\ldots),\Psi_3(a_{\rho(i)}\ldots a_{\rho(1)}\ldots))$$

is still of the form (2), albeit  $a_{\rho(i)}$  occurs in a different position in  $\Psi_3(a_{\rho(1)} \dots a_{\rho(i)} \dots)$  and  $a_{\rho(1)}$  occurs in a different position in  $\Psi_3(a_{\rho(i)} \dots a_{\rho(1)} \dots)$ . It follows, by the definition of the length of edges, that  $\Psi_3$  preserves lengths of edges.

Lemma 7: The six cycles  $C_1, C_2, C_3, C_4, C_5, C_6$  formed by starting at vertices *abcde*, *abdec*, *abced*, *abedc*, *abecd*, *abdce* respectively, and progressing along edges of length 1 until cycles are completed, partition  $St_5$  into 6 disjoint cycles. *Proof:* We list the 20 vertices of  $C_1$ :

abcde, bacde, cabde, dabce, eabcd, aebcd, beacd, ceabd, deabc, edabc, adebc, bdeac, cdeab, dceab, ecdab, acdeb, bcdea, cbdea, dbcea, ebcda

As, we have that

 $\Psi_2(abcde) = abecd, \Psi_3(abcde) = abdec, \Psi_4(abcde) = abedc,$ 

 $\Psi_5(abcde) = abced$ , and  $\Psi_6(abcde) = abdce$ ,

it follows by Lemma 6 that the other cycles also contain 20 vertices. The only vertex with ab in the first two positions in  $C_1$  is *abcde*. By Lemma 6, as  $\Psi_2$ ,  $\Psi_3$ ,  $\Psi_4$ ,  $\Psi_5$  and  $\Psi_6$  only reorder the last 3 elements of a vertex, the only vertices with ab in the first two positions in  $C_2, C_3, C_4, C_5$ , and  $C_6$  are *abecd*, *abdec*, *abecd*, *abcde*, *abcde*,

*Lemma 8:* The cycle  $C_7$  given by:

bacde, abcde, cbade, dbace, bdace, adbce,

cdbae, dcbae, bcdae, acdbe, cadbe, dacbe, bacde (3)

meets each  $C_i$   $(1 \le i \le 6)$  at exactly two vertices and a common edge.

Proof: We have that:

$$(bacde, abcde) \in E(C_1)$$

by Lemma 7,

$$(cbade, dbace) = (\Psi_2(cbdea), \Psi_2(dbcea)) \in E(C_2)$$

by Lemmas 7 and 6,

$$(bdace, adbce) = (\Psi_3(bdeac), \Psi_3(adebc)) \in E(C_3)$$

by Lemmas 7 and 6,

$$(cdbae, dcbae) = (\Psi_4(cdeab), \Psi_4(dceab)) \in E(C_4)$$

by Lemmas 7 and 6,

 $(bcdae, acdbe) = (\Psi_5(bcdea), \Psi_5(acdeb)) \in E(C_5)$ 

by Lemmas 7 and 6, and

 $(cadbe, dacbe) = (\Psi_6(cabde), \Psi_6(dabce)) \in E(C_6)$ 

by Lemmas 7 and 6.

## V. EDGE-DISJOINT HAMILTONIAN CYCLES

The Hamiltonian cycle defined by means of  $C_1, \ldots C_7$  in Section IV, produces a Hamiltonian cycle when mapped by the automorphism  $\Phi_5$  of Section III. It remains to show that the 2 Hamiltonian cycles are edge disjoint.

Lemma 9:

- (i) For the set of vertices  $V(C_7)$  of  $C_7$ , we have that  $\Phi_5(V(C_7)) = V(C_7)$ .
- (ii) For the set of edges  $E(C_7)$  of  $C_7$ , we have that  $\Phi_5(E(C_7) \cap \bigcup_{i=1}^6 E(C_i)) = E(C_7) \bigcup_{i=1}^6 E(C_i).$

*Proof:* The vertices of (3) are mapped to

acdbe, cadbe, dacbe, bacde, abcde, cbade,

$$dbace, bdace, adbce, cdbae, dcbae, bcdae, acdbe$$
 (4)

respectively, by  $\Phi_5$ . We see that (4) is just a reordering of the edges in (3) and thus (i) holds. For (ii), consider the edges of  $E(C_7) \cap \bigcup_{i=1}^{6} E(C_i)$  in Lemma 8 shown underlined below:

<u>bacde</u>, <u>abcde</u>, <u>cbade</u>, <u>dbace</u>, <u>bdace</u>, <u>adbce</u>, cdbae, <u>dcbae</u>, <u>bcdae</u>, <u>acdbe</u>, <u>cadbe</u>, <u>dacbe</u>,

We have:

$$(\Phi_{5}(bacde), \Phi_{5}(abcde)) =$$

$$(acdbe, cadbe) \in E(C_{7}) - \bigcup_{i=1}^{6} E(C_{i})$$

$$(\Phi_{5}(cbade), \Phi_{5}(dbace)) =$$

$$(dacbe, bacde) \in E(C_{7}) - \bigcup_{i=1}^{6} E(C_{i})$$

$$(\Phi_{5}(bdace), \Phi_{5}(adbce)) =$$

$$(abcde, cbade) \in E(C_{7}) - \bigcup_{i=1}^{6} E(C_{i})$$

$$(\Phi_{5}(cdbae), \Phi_{5}(dcbae)) =$$

$$(dbace, bdace) \in E(C_{7}) - \bigcup_{i=1}^{6} E(C_{i})$$

$$(\Phi_{5}(cdbae), \Phi_{5}(acdbe)) =$$

$$(adbce, cdbae) \in E(C_7) - \bigcup_{i=1}^{6} E(C_i)$$
  
 $(\Phi_5(cadbe), \Phi_5(dacbe)) =$   
 $(dcbae, bcdae) \in E(C_7) - \bigcup_{i=1}^{6} E(C_i)$ 

Theorem 10: The Hamiltonian cycles C and  $\Phi_5(C)$  are edge-disjoint.

*Proof:* Let  $v \in V(C) - V(C_7)$ ,  $v \in V(C_i)$  say, where  $1 \le i \le 6$ . Then, there exist  $u_1, u_2 \in V(C_i)$  such that the edges incident at v in C,  $(u_1, v)$  and  $(v, u_2)$ , belong to  $C_i$  and so, by the definition of  $C_i$  in Lemma 7,

$$\lambda(u_1, v) = \lambda(v, u_2) = 1 \tag{5}$$

Consider the edges  $(v_1, v)$  and  $(v_2, v)$  incident at v in  $\Phi_5(C)$ . As  $v \notin V(C_7)$ , by Lemma 9(i)  $v = \Phi_5(v')$  for some  $v' \in V(C) - V(C_7)$ , say  $v' \in C_j$  where  $1 \leq j \leq 6$ . Then, there exist edges  $(v'_1, v')$ ,  $(v', v'_2)$  in  $C_j$  such that  $\Phi_5(v'_1) = v_1$  and  $\Phi_5(v'_2) = v_2$ . By the definition of  $C_j$  in Lemma 7,

$$\lambda(v_1', v') = \lambda(v', v_2') = 1 \tag{6}$$

By (6) and Lemma 5,

$$\lambda(\Phi_5(v_1'), \Phi_5(v')) = \lambda(\Phi_5(v'), \Phi_5(v_2')) = 2$$

i.e.

$$\lambda(v_1, v) = \lambda(v, v_2) = 2 \tag{7}$$

By (5) and (7), different edges are incident at the vertex  $v \in C - C_7$  in C and  $\Phi_5(C)$ . Hence, we have shown that an edge in C, which has a vertex not in  $C_7$ , cannot belong to  $\Phi_5(C)$ . It follows that an edge common to both C and  $\Phi_5(C)$  must be an edge in  $C_7$ . So, let e be an edge of C belonging to  $C_7$ . For  $1 \le i \le 6$ , e cannot be an edge in  $C_i$  as C does not contain edges common to  $C_7$  and  $C_i$ . By Lemma 9(ii),  $\Phi_5$  maps an edge in  $C_7$  and some  $C_i$ , where  $1 \le i \le 6$  to e. Thus,  $\Phi_5$  maps an edge not in C to e. Therefore  $e \notin \Phi_5(C)$ . We conclude that C and  $\Phi_5(C)$  have no common edges.

# VI. CONCLUSIONS

We have given a Hamiltonian decomposition of 5-star, based on a graph automorphism relating the two Hamiltonian cycles. Our further work will investigate properties of similar automorphisms in higher degree star graphs to determine whether they can be used to establish or refute the existence of Hamiltonian decompositions there.

## ACKNOWLEDGEMENTS

We are grateful to the late Professor Ondrej Sykora for introducing us to the problem.

# International Journal of Information, Control and Computer Sciences ISSN: 2517-9942 Vol:4, No:1, 2010

# REFERENCES

- [1] S.B.Akers, D.Harel, and B.Krishnamurthy, "The star graph: an attractive alternative to the n-cube," Int. Conf. on Parallel Processing, Chicago, USA, 1987, 393-400.
- [2] K.Okuda and S.W.Song, "Revisiting Hamiltonian decomposition of the hypercube," Proc. 13th Symp. on Integrated Circuits and Systems Design, Manaus, Brazil, 2000, 55-60.
- [3] R.Rowley and B.Bose, "On the number of disjoint Hamiltonian cycles in De Bruijn Graphs," Oregon State University Technical Report 93-80-09, 1993
- [4] M.M.Bae and B.Bose, "Edge disjoint Hamiltonian cycles in k-ary n-cubes and hypercubes," IEEE Transactions on Computers, 52(10), 2003, 1271-1284
- [5] R.Cada, T.Kaiser, M.Rosenfeld, and Z.Ryjacek, "Disjoint Hamiltonian cycles in the star graph," Information Processing Letters, 110(1), 2009, 30-35.
- [6] S.J.Curran, and J.A.Gallian, "Hamiltonian cycles and paths in Cayley graphs and digraphs a survey," *Discrete Mathematics*, 156, 1996, 1-18.
  [7] T.-K.Li, J.J.M.Tan, L.-H.Hsu, "Hyper Hamiltonian laceability on edge fault star graph," *Information Sciences*, 165, 2004, 59-71.
- [8] J.-C.Bermond, O.Favaron, and M.Maheo, "Hamiltonian decomposition of Cayley graphs of degree 4," Journal of Combinatorial Theory, 46, 1989, 142-153.
- [9] J.-H.Park, "Hamiltonian decomposition of recursive circulants," Proc. 9th Int. Symp. of Algorithms and Computation ISAAC'98, Taejon, Korea, 1998, 297-306.
- [10] V.L.Kompel'makher, and V.A.Liskovets, "Sequential generation of arrangements by means of a basis of transpositions," Kibernetica, 3, 1975, 17-21.