

# A Numerical Method for Diffusion and Cahn-Hilliard Equations on Evolving Spherical Surfaces

Jyh-Yang Wu, Sheng-Gwo Chen

**Abstract**—In this paper, we present a simple effective numerical geometric method to estimate the divergence of a vector field over a curved surface. The conservation law is an important principle in physics and mathematics. However, many well-known numerical methods for solving diffusion equations do not obey conservation laws. Our presented method in this paper combines the divergence theorem with a generalized finite difference method and obeys the conservation law on discrete closed surfaces. We use the similar method to solve the Cahn-Hilliard equations on evolving spherical surfaces and observe stability results in our numerical simulations.

**Keywords**—Conservation laws, diffusion equations, Cahn-Hilliard Equations, evolving surfaces.

## I. INTRODUCTION

FINDING numerical methods to compute partial differential equations on evolving surfaces is an interesting and difficult problem. These methods have many important applications in fluid dynamics, magnetohydrodynamics, image processing, and so on. See more details in [1], [2], [9]. In this note, we shall introduce a new numerical method for solving the diffusion equation on evolving closed surfaces that we proposed in 2016. This method is an intrinsic geometric method to deal with the discrete conservation law on evolving regular surfaces, and we shall improve the Cahn-Hilliard equation on evolving spherical surfaces by this method.

The Cahn-Hilliard equation with a variable mobility on a regular closed surface  $\Sigma$  takes the form:

$$\frac{\partial u(x,t)}{\partial t} = \nabla \cdot [M(u(x,t)) \nabla \mu(u(x,t))],$$

$$x \in \Sigma, 0 < t \leq T. \text{ and } \mu(u(x,t)) = F'(u(x,t)) - \varepsilon^2 \Delta_{\Sigma} u(x,t)$$

where the quantity  $u(x,t)$  is the difference between the mole fractions of binary mixtures. The function  $F(u) = \frac{1}{4}(u^2 - 1)^2$  is the Helmholtz free energy per unit volume of a homogeneous fluid, and  $\varepsilon$  is a positive constant. See [10], [11] for more details.

## II. PRELIMINARIES

First, we introduce a discrete Laplace-Beltrami operator on a stationary surface that we proposed in 2013 and 2014 [3], [4].

J.-Y. Wu is with the Department of Information Management, Yuan Ze University, Taiwan (e-mail: jywu@saturn.yzu.edu.tw).

S.-G. Chen is with the Department of Applied Mathematics, National Chiayi University, Taiwan (corresponding author; e-mail: csg@mail.ncyu.edu.tw).

### A. The Local Tangential Method [3]

Let  $S = (V, F)$  be a triangular mesh with  $V = \{p_i \mid 1 \leq i \leq n_v\}$  the list of vertices and  $F = \{T_k \mid 1 \leq k \leq n_f\}$  the list of triangles. We introduce the approximating tangent plane  $TS(p)$  at the vertex  $p$  of as:

1. The unit normal vector  $N_A(p)$  at the vertex  $p$  in  $S$  is given by the weighted normal vector

$$N_A(p) = \frac{\sum_{k \in I(p)} w_{T_k} N_{T_k}}{\left\| \sum_{k \in I(p)} w_{T_k} N_{T_k} \right\|}, \quad (1)$$

where  $N_{T_k}$  is the unit normal vector of the triangle  $T_k$  including a vertex  $p$ . We refer to [3] for more details.

2. The approximating tangent space  $TS(p)$  of  $S$  at  $p$  is now determined by

$$TS(p) = \{w \in \mathbf{R}^3 \mid w \perp N_A(p)\}. \quad (2)$$

We can choose an orthonormal basis  $e_1, e_2$  for the tangent plane  $TS(p)$  of  $S$  at  $p$ . Hence,  $\{e_1, e_2, N_A(p)\}$  forms an orthonormal basis for  $\mathbf{R}^3$  and every  $q \in \Sigma$  around  $p$  can be assigned a new  $xyz$ -coordinate by

$$x(q)e_1 + y(q)e_2 = (q - p) - ((q - p) \cdot N_A(p))N_A(p) \quad (3)$$

and

$$z(q) = h(x(p), y(p)) = (q - p) \cdot N_A(p). \quad (4)$$

Obviously, the new coordinate of  $p$  is  $(0,0,0)$ .

### B. Discrete Tangential Gradient Vector

Since the gradient  $\nabla_{\Sigma} f$  of a smooth function  $f$  on a regular surface  $\Sigma$  with a parametrization  $\mathbf{x}(u, v)$  is given by

$$\nabla_{\Sigma} f = \frac{f_u G - f_v F}{EG - F^2} \mathbf{x}_u + \frac{f_v E - f_u F}{EG - F^2} \mathbf{x}_v, \quad (5)$$

where  $E, F,$  and  $G$  are the coefficients of the 1<sup>st</sup> fundamental form of  $\Sigma$  and  $f_u = \frac{\partial}{\partial u} f(\mathbf{x}(u, v))$  and  $f_v = \frac{\partial}{\partial v} f(\mathbf{x}(u, v))$ , we

need to approximate the local parametrization of  $\Sigma$  around  $p$  and the differential quantities of the function  $f$ .

We shall construct a local parametrization by representing the regular surface  $\Sigma$  as locally a graph surface around the vertex  $p$ . Consider the triangular mesh  $S = (V, F)$  of a closed surface  $\Sigma$  with mesh size  $r > 0$ . Given a vertex  $p \in V$ , let  $p_j, j = 0, 1, \dots, n$  be the neighboring vertices of  $p$  with  $p_0 = p$ . Suppose that the new coordinate of  $p_j$  is  $(x_j, y_j, z_j)$ , and we denote  $h(x_j, y_j) = z_j$ . We use the polynomial fitting for the height function  $h$  of  $\Sigma$  around  $p$ . By the Taylor expansion, one has

$$h(x_j, y_j) - h(0,0) = x_j h_x(0,0) + y_j h_y(0,0) + \frac{1}{2}(x_j^2 h_{xx}(0,0) + 2x_j y_j h_{xy}(0,0) + y_j^2 h_{yy}(0,0)) + \dots + O(r_j^m) \quad (6)$$

for each  $j = 1, \dots, n$ .

Set  $A = \begin{pmatrix} x_1 & \dots & x_n \\ y_1 & \dots & y_n \\ x_1^2 & \dots & x_n^2 \\ x_1 y_1 & \dots & x_n y_n \\ y_1^2 & \dots & y_n^2 \end{pmatrix}$  and  $\alpha_j, j = 1, \dots, n$  is a set of

real numbers. Then, we have

$$h_u(0,0) = \sum_{j=1}^n \alpha_j (h(u_j, v_j) - h(0,0)) + O(r^3) \quad (7)$$

if  $A \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix} = (1 \ 0 \ 0 \ 0 \ 0)$  and

$$h_v(0,0) = \sum_{j=1}^n \alpha_j (h(u_j, v_j) - h(0,0)) + O(r^3) \quad (8)$$

if  $A \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix} = (0 \ 1 \ 0 \ 0 \ 0)$ .

Similarly, we can also approximate the differential quantities of the function  $f$  on  $\Sigma$ . Therefore, the gradient vector  $\nabla_\Sigma f$  at  $p$  can be approximated by

$$\nabla_A f(p) = \frac{1}{1+h_u^2+h_v^2} \begin{pmatrix} f_u(1+h_u^2) - f_v h_u h_v \\ f_v(1+h_v^2) - f_u h_u h_v \\ f_u + f_v \end{pmatrix} \quad (9)$$

We refer to [5] for the detail of the high-order approach.

**Theorem 1.** Using above notations, one has

$$\nabla_\Sigma f(p) = \nabla_A f(p) + O(r^2) \quad (10)$$

where

$$\nabla_A f(p) = \frac{1}{1+h_u^2+h_v^2} \begin{pmatrix} f_u(1+h_u^2) - f_v h_u h_v \\ f_v(1+h_v^2) - f_u h_u h_v \\ f_u + f_v \end{pmatrix} \quad (11)$$

*C. Discrete Laplace-Beltrami Operator*

If  $X$  is a local vector field  $X = A\mathbf{x}_u + B\mathbf{x}_v$  on  $\mathbf{x}(U) \subset \Sigma$ . The divergence  $Div_\Sigma X$ , of  $X$  is defined as a function  $Div_\Sigma X : \mathbf{x}(U) \rightarrow \mathbf{R}$  given by the trace of the linear map  $Y(p) \rightarrow \nabla_{Y(p)} X$  for  $p$  in  $\Sigma$ . A direct computation yields

$$Div_\Sigma X = \frac{1}{\sqrt{EG-F^2}} \left[ \frac{\partial}{\partial u} (A\sqrt{EG-F^2}) + \frac{\partial}{\partial v} (B\sqrt{EG-F^2}) \right] \quad (12)$$

Note that the divergence theorem gives

$$\int_U Div_\Sigma X = \int_{\partial U} X \cdot \mathbf{n} \quad (13)$$

where  $\mathbf{n}$  is the unit outward normal vector on  $\partial U$ .

The Laplace-Beltrami operator  $\Delta_\Sigma f$  acting on the function  $f$  is defined by  $\Delta_\Sigma f = Div_\Sigma(\nabla_\Sigma f)$  and has the local representation

$$\Delta_\Sigma f = \frac{1}{\sqrt{EG-F^2}} \left[ \frac{\partial}{\partial u} \left( \frac{G}{\sqrt{EG-F^2}} f_u \right) - \frac{\partial}{\partial u} \left( \frac{F}{\sqrt{EG-F^2}} f_v \right) + \frac{\partial}{\partial v} \left( \frac{E}{\sqrt{EG-F^2}} f_v \right) - \frac{\partial}{\partial v} \left( \frac{F}{\sqrt{EG-F^2}} f_u \right) \right] \quad (14)$$

See [7], [8] for details.

We use the divergence theorem to give a discrete approximation of the divergence of a vector field  $X$  defined on a triangular surface mesh  $S = (V, F)$ . Consider a vertex  $p \in V$  and let  $p_j, j = 0, 1, \dots, n$ , be the neighboring vertices of  $p$  with  $p_0 = p$ . These vertices  $p_j$  are labeled counterclockwise about the normal vector  $N_A(p)$ . Let  $T_j$  be the triangle with vertices  $p, p_j$ , and  $p_{j+1}$ . We define the approximating outer normal vectors  $n(T_j, p_j)$  and  $n(T_j, p_{j+1})$  of the triangle  $T_j$  at the vertex  $p_j$  and  $p_{j+1}$  in  $\Sigma$  by

$$n(T_j, p_j) = \frac{(p_{j+1} - p_j) \wedge N_A(p_j)}{\|(p_{j+1} - p_j) \wedge N_A(p_j)\|} \quad (15)$$

and

$$n(T_j, p_{j+1}) = \frac{(p_{j+1} - p_j) \wedge N_A(p_{j+1})}{\|(p_{j+1} - p_j) \wedge N_A(p_{j+1})\|}. \quad (16)$$

Since

$$\int_{\partial U} X \cdot n = \sum_{j=0}^{n-1} \left\{ \frac{\|p_{j+1} - p_j\|}{6} (1 + O(r^2)) [2X(p_j) \cdot n(T_j, p_j) + X(p_{j+1}) \cdot n(T_j, p_j) + 2X(p_{j+1}) \cdot n(T_j, p_{j+1}) + X(p_j) \cdot n(T_j, p_{j+1})] (1 + O(r)) \right\} \quad (17)$$

and

$$\int_U \text{Div}_\Sigma X = \left( \sum_{j=0}^{n-1} |T_j| (1 + O(r^2)) \right) (\text{Div}_\Sigma X(p) + O(r)). \quad (18)$$

We can now define the discrete divergence  $\text{Div}_A X$  of a vector field on the triangular surface mesh  $\mathbf{S}$  by

$$\text{Div}_A X(p) = \frac{1}{\sum_{k=0}^{n-1} |T_k|} \left[ \sum_{j=0}^{n-1} \frac{\|p_{j+1} - p_j\|}{6} ((2X(p_j) + X(p_{j+1})) \cdot n(T_j, p_j) + (X(p_j) + 2X(p_{j+1})) \cdot n(T_j, p_{j+1})) \right] \quad (19)$$

where  $|T_k|$  denotes the area of the triangle  $T_k$ .

**Theorem 2.** Let  $f$  be a smooth function defined on a regular surface  $\Sigma$  and the vector field  $\nabla_A f$  on  $S$  satisfy

$$\nabla_\Sigma f = \nabla_A f + O(r^2) \quad (20)$$

where  $\nabla_\Sigma f$  is the gradient vector field of  $f$  on  $\Sigma$ . Then, we have

$$\Delta_\Sigma f(p) = \text{Div}_A(\nabla_A f)(p) + O(r). \quad (21)$$

### III. DISCRETE ALGORITHMS OF PDES ON EVOLVING SURFACES

Next, we discuss the discrete algorithms about the diffusion equation and Cahn-Hilliard equation on evolving surfaces.

#### A. Diffusion Equations on Evolving Surfaces

Let  $\{\Sigma(t)\}$ ,  $t \in [0, T]$ , denote a moving oriented regular surfaces in  $\mathbf{R}^3$ . Suppose that these regular surfaces are moving with prescribed velocity field  $X(t)(p(t))$ ,  $p(t) \in \Sigma(t)$ . We want to solve the surface diffusion equation:

$$\partial_t^* h + h \nabla_{\Sigma(t)} \cdot X - \Delta_{\Sigma(t)} h = 0 \quad (22)$$

on evolving surfaces  $\Sigma(t)$ . Here,  $\partial_t^* = \partial_t + X \cdot \nabla$  is the material derivative. See Elliott and Ranner [11] for more details.

After some direct computations, one has

$$\frac{dh(t)}{dt}(u, v) + \frac{1}{2} h(t)(u, v) [\text{div}_{\Sigma(t)} V(t)(u, v) - 4c(t)(u, v) H(t)(u, v)] = \Delta_{\Sigma(t)} h(t)(u, v), \quad (23)$$

where  $H(t)(u, v)$  is the mean curvature on the surface  $\Sigma(t)$ .

This is the equation for the conservation law on the evolving surface  $\Sigma(t)$  with the velocity field  $X(t)(u, v) = V(t)(u, v) + c(t)(u, v)N(t)(u, v)$ . See [6] about these computations.

Since the divergence theorem, integrating (22) on a portion  $U(t) \subset \Sigma(t)$ , one has

$$\int_{U(t)} \partial_t^* h + h \nabla_{\Sigma(t)} \cdot X = \int_{U(t)} \Delta_{\Sigma(t)} h \quad (24)$$

and this is equivalent to

$$\frac{d}{dt} \int_{U(t)} h = \int_{U(t)} \Delta_{\Sigma(t)} h = \int_{\partial U(t)} \nabla_{\Sigma(t)} h \cdot \mathbf{n}(t). \quad (25)$$

After an explicit time discretization, (25) becomes

$$\frac{1}{\tau^n} \left( \int_{U^{n+1}} h^{n+1} - \int_{U^n} h^n \right) = \int_{U^n} \Delta_{\Sigma^n} h^n = \int_{\partial U^n} (\nabla_{\Sigma^n} h^n) \cdot \mathbf{n}(t) \quad (26)$$

or

$$\left( \int_{U^{n+1}} h^{n+1} \right) = \int_{U^n} h^n + \tau^n \int_{U^n} \Delta_{\Sigma^n} h^n \quad (27)$$

After space discretization, (27) yields

$$h^{n+1}(p^{n+1}) = \frac{\sum_{p^j \in T_k^n} |T_k^n|}{\sum_{p^{n+1} \in T_k^{n+1}} |T_k^{n+1}|} \left[ h^n(p^n) + \tau^n \Delta_{\Sigma^n} h^n(p) \right] \quad (28)$$

Now our two-step algorithm can be stated as follows.

Step 1: For each  $p$ , use the above method to compute the Laplace-Beltrami operator  $\Delta_A h^j(p^j)$  and set

$$\bar{h}^j(p^j) = h^j(p^j) + \tau^j \Delta_A h^j(p^j) \quad (29)$$

Step 2: Let  $p_k^j$ ,  $k = 0, 1, \dots, n$ , be the neighboring vertices of  $p^j$

with  $p_0^j = p^n$ . These vertices  $p_k^j$  are labeled counterclockwise about the normal vector  $N_A(p^j)$  of the surface  $\Sigma^j$  at the vertex  $p^j$  in the space  $\mathbf{R}^3$ . Let  $T_k^j$  be the triangle with vertices  $p^j$ ,  $p_k^j$  and  $p_{k+1}^j$ ,  $j \geq 0$ . We set

$$h^{j+1}(p^{j+1}) = \frac{\sum_k |T_k^j|}{\sum_k |T_k^{j+1}|} \bar{h}^j(p^j). \quad (30)$$

One can find the proofs of the convergence and conservation law of this method in [6].

*B. Cahn-Hilliard Equation on Evolving Surfaces*

We solve the Cahn-Hilliard equation

$$u_t = \Delta_{\Sigma(t)}(u^3 - u - \varepsilon^2 \Delta_{\Sigma(t)} u) \quad (31)$$

on an evolving surface  $\Sigma(t)$ .

Set  $\phi = u^3 - u - \varepsilon^2 \Delta_{\Sigma(t)} u$ , then (31) can be rewritten as

$$\begin{cases} u_t = \Delta_{\Sigma(t)} \phi, \\ \phi = u^3 - u - \varepsilon^2 \Delta_{\Sigma(t)} u. \end{cases} \quad (32)$$

Using the explicit time discretization, the first equation in (28) becomes

$$\frac{u^{n+1} - u^n}{\tau^n} = \Delta_{\Sigma^n} \phi^n \quad (33)$$

or

$$u^{n+1} = u^n + \tau^n \Delta_{\Sigma^n} \phi^n. \quad (34)$$

The function  $\phi^n$  is given by

$$\phi^n = (u^n)^3 - u^n - \varepsilon^2 \Delta_{\Sigma^n} u^n. \quad (35)$$

IV. SIMULATIONS

Now we solve (32) with  $\varepsilon = 0.1$  on two different evolving surfaces. All initial conditions of our simulations are the same random numbers in  $[-0.5, 0.5]$ , and the time step is  $\frac{r^2}{100}$ .

We solve (32) on the unit sphere moving with a prescribed velocity

$$\mathbf{X}(t, p) = \begin{pmatrix} 0.5s\pi \cos(st\pi) \sin(0.5z\pi) \\ 0 \\ 0 \end{pmatrix} \quad (36)$$

and

$$\mathbf{Y}(t, p) = \begin{pmatrix} 0.5s \cos(st\pi) \cos(0.5z\pi) \\ 0 \\ 0 \end{pmatrix} \quad (37)$$

where  $p = (x, y, z)$  in the unit sphere and  $s$  is a nonzero real number. We show all solution of (34) at times are 0.75 and 1.

Fig. 1 shows the solution of (32) on a stationary unit sphere. Figs. 2-7 present the numerical solutions of (32) with  $s = 1, 2, 10, 50, 250$ , and  $500$ . Fig. 8 is the solution with  $\mathbf{X}(t, p) = (500\pi \cos(500t\pi) \sin(0.5z\pi) \ 0 \ 0)$ . Fig. 9 shows the Ginzberg-Landau free energy of (32) on the moving surfaces with different  $s$  in (36).

Figs. 10-15 show the numerical solution on a unit sphere with velocity field  $\mathbf{Y}$ .

V. CONCLUSION

Conservation laws play important key roles in the partial differential equation on surfaces. An efficient numerical method should at least obey the conservation law. Our proposed method obeys the discrete conservation law. Furthermore, the Ginzberg-Landau free energy is decreasing in our simulations.




Fig. 1 The solution of (26) on a unit sphere




Fig. 2 The solution of (26) with  $s = 1$  in the velocity field  $X$




Fig. 3 The solution of (26) with  $s = 2$  in the velocity field  $X$




Fig. 4 The solution of (26) with  $s = 10$  in the velocity field  $X$




Fig. 5 The solution of (26) with  $s = 50$  in the velocity field X




Fig. 6 The solution of (26) with  $s = 250$  in the velocity field X




Fig. 7 The solution of (26) with  $s = 500$  in the velocity field X




Fig. 8 The solution of (26) with  $X(1) = 500\pi \cos(500t\pi) \sin(0.5z\pi)$




Fig. 9 The Ginzberg-Landau free energy with different  $s$




Fig. 10 The solution of (26) with  $s = 1$  in the velocity field  $Y$




Fig. 11 The solution of (26) with  $s = 2$  in the velocity field  $\mathbf{Y}$




Fig. 12 The solution of (26) with  $s = 10$  in the velocity field  $\mathbf{Y}$




Fig. 13 The solution of (26) with  $s = 50$  in the velocity field  $\mathbf{Y}$






Fig. 14 The solution of (26) with  $s = 250$  in the velocity field  $Y$




Fig. 15 The solution of (26) with  $s = 500$  in the velocity field  $Y$

ACKNOWLEDGMENT

This work was partially supported by MOST, Taiwan.

[11] C. M. Elliott and T. Ranner, "Evolving surface finite element method for the Cahn–Hilliard equation", *Numer. Math.* vol.3, 2015, pp483-534.

REFERENCES

[1] M. Burger, L. He and C.-B. Schonlieb, "Cahn-Hilliard inpainting and a generalization for grayvalue images", *SIAM J. Imaging Sciences*, vol. 2, 2009, pp. 1129-1167.  
 [2] J. Cahn, J. Hilliard, "Free energy of a non-uniform system. I. interfacial free energy", *J. Chem. Phys.* vol. 28, 1958, pp. 258–267.  
 [3] S.-G. Chen and J.-Y. Wu, "Discrete conservation laws on curved surfaces", *SIAM J. Sci. Comput.*, vol. 35, 2013, pp. A719-A739.  
 [4] S.-G. Chen and J.-Y. Wu, "Discrete conservation laws on curved surfaces II: A dual approach", *SIAM J. Sci. Comput.*, vol. 36, 2014, pp. A1813-A1830.  
 [5] S.-G. Chen, M.-H. Chi, and J.-Y. Wu, "High-Order Algorithms for Laplace–Beltrami Operators and Geometric Invariants over Curved Surfaces", vol 65, 2015, pp.839-865.  
 [6] S.-G. Chen and J.-Y. Wu, "Discrete conservation laws on evolving surfaces", *SIAM J. Sci. Comput.*, vol. 38, 2016, pp. A1725-A1742.  
 [7] M. DoCarmo, "Differential geometry of curves and surfaces", Prentice-Hall, London, 1976.  
 [8] M. DoCarmo, "Riemannian geometry", Birkhauser, Boston, 1992.  
 [9] I. Dolcetta, S. Vita, R. March, "Area-preserving curve-shortening flows: from phase separation to image processing", *Interfaces Free Bound.* vol. 4, 2002, pp. 325-343.  
 [10] Q. Dua, L. Ju and Li Tian, "Finite element approximation of the Cahn–Hilliard equation on surfaces", *Comput. Methods Appl. Mech. Engrg.*, vol. 200, 2011, pp. 2458-2470.