

# An Accurate Computation of Block Hybrid Method for Solving Stiff Ordinary Differential Equations

A. M. Sagir

**Abstract**—In this paper, self-starting block hybrid method of order (5,5,5,5)<sup>T</sup> is proposed for the solution of the special second order ordinary differential equations with associated initial or boundary conditions. The continuous hybrid formulations enable us to differentiate and evaluate at some grids and off – grid points to obtain four discrete schemes, which were used in block form for parallel or sequential solutions of the problems. The computational burden and computer time wastage involved in the usual reduction of second order problem into system of first order equations are avoided by this approach. Furthermore, a stability analysis and efficiency of the block method are tested on stiff ordinary differential equations, and the results obtained compared favorably with the exact solution.

**Keywords**—Block Method, Hybrid, Linear Multistep Method, Self – starting, Special Second Order.

## I. INTRODUCTION

LET us consider the numerical solution of the special second order ordinary differential equation of the form

$$y'' = f(x, y), \quad a \leq x \leq b \quad (1)$$

with associated initial or boundary conditions. The mathematical models of most physical phenomena especially in mechanical systems without dissipation leads to special second order initial value problem of type (1). Solutions to initial value problem of type (1) according to Fatunla [1], [2] are often highly oscillatory in nature and thus, severely restrict the mesh size of the conventional linear multistep method. Such system often occurs in mechanical systems without dissipation, satellite tracking, and celestial mechanics.

Lambert [3] and several authors such as Onumanyi *et al* [4], Awoyemi [5], Yahaya and Adegboye [6], and Fudziah *et al*. [7], have written on conventional linear multistep method

$$\sum_{j=0}^k \alpha_j y_{n+j} = h^2 \sum_{j=0}^k \beta_j f_{n+j}, \quad k \geq 2 \quad (2)$$

or compactly in the form

$$\rho(E)y_n = h^2 \delta(E)f_n \quad (3)$$

where  $E$  is the shift operator specified by  $E^j y_n = y_{n+j}$  while  $\rho$  and  $\delta$  are characteristics polynomials and are given as

$$\rho(\xi) = \sum_{j=0}^k \alpha_j \xi^j, \quad \delta(\xi) = \sum_{j=0}^k \beta_j \xi^j \quad (4)$$

$y_n$  is the numerical approximation to the theoretical solution  $y(x)$  and  $f_n = f(x_n, y_n)$ .

In the present consideration, our motivations for the study of this approach is a further advancement in efficiency, i.e obtaining the most accuracy per unit of computational effort, that can be secured with the group of methods proposed in this paper over Taparki and Odekunle [8]; and Adeboye [9].

*A. Definition : Consistent, Lambert [3]*

The linear multistep method (2) is said to be consistent if it has order  $p \geq 1$ , that is if

$$\sum_{j=0}^k \alpha_j = 0 \quad \text{and} \quad \sum_{j=0}^k j \alpha_j - \sum_{j=0}^k \beta_j = 0 \quad (5)$$

Introducing the first and second characteristics polynomials (4), we have from (5) LMM type (2) is consistent if

$$\rho(1) = 0, \quad \rho^1(1) = \delta(1)$$

*B. Definition: Zero Stability, Lambert [3]*

A linear multistep method type (2) is zero stable provided the roots  $\xi_j, j = 0(1)k$  of first characteristics polynomial  $\rho(\xi)$  specified as  $\rho(\xi) = \det[\sum_{j=0}^k A(i)\xi^{k-i}] = 0$  satisfies  $|\xi_j| \leq 1$  and for those roots with  $|\xi_j| = 1$  the multiplicity must not exceed two. The principal root of  $\rho(\xi)$  is denoted by  $\xi_1 = \xi_2 = 1$ .

*C. Definition: Convergence, Lambert [3]*

The necessary and sufficient conditions for the linear multistep method type (2) is said to be convergent if it is consistent and zero stable.

*D. Definition: Order and Error Constant, Lambert [3]*

The linear multistep method type (2) is said to be of order  $p$  if  $c_0 = c_1 = \dots = c_{p+1} = 0$  but  $c_{p+2} \neq 0$  and  $c_{p+2}$  is called the error constant, where

$$c_0 = \sum_{j=0}^k \alpha_j = \alpha_0 + \alpha_1 + \alpha_2 + \dots + \alpha_k$$

A.M. Sagir is with Department of Basic Studies, College of Basic and Remedial Studies, Hassan Usman Katsina Polytechnic, P.M.B. 2052 Katsina, Katsina State, Nigeria (phone: 08039474856, e-mail: amsagir@yahoo.com).

$$\begin{aligned}
 c_1 &= \sum_{j=0}^k j \alpha_j = (\alpha_1 + 2 \alpha_2 + 3 \alpha_3 + \dots + k \alpha_k) \\
 &\quad - (\beta_0 + \beta_1 + \beta_2 + \dots + \beta_k) \\
 c_2 &= \sum_{j=0}^k \frac{1}{2!} j^2 \alpha_j - \sum_{j=0}^k \beta_j \\
 &= \left\{ \begin{aligned} &\frac{1}{2!} (\alpha_1 + 2^2 \alpha_2 + 3^2 \alpha_3 + \dots + k^2 \alpha_k) \\ &- (\beta_1 + 2\beta_2 + 3\beta_3 + \dots + k\beta_k) \end{aligned} \right\} \\
 &\quad \cdot \\
 &\quad \cdot \\
 c_q &= \sum_{j=1}^k \left\{ \frac{1}{q!} j^q \alpha_j - \frac{1}{(q-2)!} j^{q-2} \beta_j \right\} \\
 &= \left\{ \begin{aligned} &\frac{1}{q!} (\alpha_1 + 2^q \alpha_2 + 3^q \alpha_3 + \dots + k^q \alpha_k) \\ &- \frac{1}{(q-1)!} (\beta_1 + 2^{(q-1)} \beta_2 + 3^{(q-1)} \beta_3 + \dots + k^{(q-1)} \beta_k) \end{aligned} \right\} \quad (6)
 \end{aligned}$$

E. Theorem: Lambert, [3]

Let  $f(x, y)$  be defined and continuous for all points  $(x, y)$  in the region  $D$  defined by  $\{(x, y) : a \leq x \leq b, -\infty < y < \infty\}$  where  $a$  and  $b$  finite, and let there exist a constant  $L$  such that for every  $x, y, y^*$  such that  $(x, y)$  and  $(x, y^*)$  are both in  $D$ :

$$|f(x, y) - f(x, y^*)| \leq L |y - y^*| \quad (7)$$

Then if  $\eta$  is any given number, there exist a unique solution  $y(x)$  of the initial value problem (1), where  $y(x)$  is continuous and differentiable for all  $(x, y)$  in  $D$ . The inequality (7) is known as a Lipschitz condition and the constant  $L$  as a Lipschitz constant.

## II. DERIVATION OF THE PROPOSED METHOD

We proposed an approximate solution to (1) in the form

$$y(x) = \sum_{j=0}^{t+m-1} a_j x^j = y_{n+j}, i = 0(1)m + t - 1 \quad (8)$$

$$y''(x) = \sum_{j=0}^{t+m-1} j(j-1) a_j x^{j-2} = f_{n+j}, \quad (9)$$

$i = 2(3)m + t - 1$  with  $m = 5, t = 2$  and  $p = m + t - 1$  where the  $a_j, j = 0, 1, (m + t - 1)$  are the parameters to be determined,  $t$  and  $m$  are points of interpolation and collocation respectively. Where  $P$ , is the degree of the polynomial interpolant of our choice.

Specifically, we collocate (9) at  $x = x_{n+j}, j = 0(1)k$  and

interpolate (8) at  $x = x_{n+j},$

$j = 0(1)k - 2$  using the method described above. Putting in the matrix equation form and then solved to obtain the values of parameters  $a_j$ 's,  $j = 0, 1, \dots$  which is substituted in (8) yields, after some algebraic manipulation, the new continuous form for the solution.

$$y(x) = \sum_{j=0}^{k-2} \alpha_j(x) y_{n+j} + \sum_{j=0}^k \beta_j(x) f_{n+j} \quad (10)$$

We set  $\xi = (x - x_{n+1})$

If we let  $k = 3$ , after some algebraic manipulations we obtain a continuous form of solution

$$\begin{aligned}
 y(x) &= \{-(\xi)\} y_n + \left\{ \left( \frac{h + \xi}{h} \right) \right\} y_{n+1} \\
 &+ \left\{ \frac{6(\xi)^6 - 30h(\xi)^5 + 45h^2(\xi)^4 - 20h^3(\xi)^3 + 101h^5(\xi)}{1440h^4} \right\} f_n \\
 &+ \left\{ \frac{-6(\xi)^6 + 21h(\xi)^5 + 5h^2(\xi)^4 - 70h^3(\xi)^3 + 60h^4(\xi)^2 + 108h^5(\xi)}{120h^4} \right\} f_{n+1} \\
 &+ \left\{ \frac{-6(\xi)^6 + 12h(\xi)^5 + 25h^2(\xi)^4 - 20h^3(\xi)^3 + 27h^5(\xi)}{240h^4} \right\} f_{n+2} \\
 &+ \left\{ \frac{54(\xi)^6 - 162h(\xi)^5 - 135h^2(\xi)^4 + 540h^3(\xi)^3 - 459h^5(\xi)}{800h^4} \right\} f_{n+\frac{4}{3}} \\
 &+ \left\{ \frac{6(\xi)^6 - 3h(\xi)^5 - 15h^2(\xi)^4 + 10h^3(\xi)^3 - 16h^5(\xi)}{1800h^4} \right\} f_{n+3} \\
 &+ \left\{ \frac{9(\eta)^5 + 10h(\eta)^4 - 10h^2(\eta)^3 + 11h^4(\eta)}{240h^3} \right\} f_{n+2} \quad (11)
 \end{aligned}$$

Evaluating (11) at  $x = x_{n+4/3}, x = x_{n+2}$  and  $x = x_{n+3}$ , yield the following schemes:

$$\begin{aligned}
 (a) \quad &y_{n+\frac{4}{3}} - \frac{4}{3} y_{n+1} + \frac{1}{3} y_n \\
 &= \frac{h^2}{437400} \{10135f_n + 146580f_{n+1} \\
 &\quad + 15690f_{n+2} - 73953f_{n+4/3} - 1252f_{n+3}\} \\
 (b) \quad &y_{n+2} - 2y_{n+1} + y_n \\
 &= \frac{h^2}{1200} \{85f_n + 1180f_{n+1} + 190f_{n+2} \\
 &\quad - 243f_{n+4/3} - 12f_{n+3}\} \\
 (c) \quad &y_{n+3} - 3y_{n+1} + 2y_n \\
 &= \frac{h^2}{1200} \{155f_n + 2640f_{n+1} + 1470f_{n+2} \\
 &\quad - 729f_{n+\frac{4}{3}} + 64f_{n+3}\} \quad (12)
 \end{aligned}$$

Taking the first derivative of (11), thereafter, evaluate the resulting continuous polynomial solution at

$x = x_0$  yields

$$\begin{aligned}
 (d) \quad &hz_0 - y_{n+1} + y_n = \frac{h^2}{7200} \{-1625f_n - 6060f_{n+1} - \\
 &1110f_{n+2} + 5103f_{n+\frac{4}{3}} + 92f_{n+3}\} \quad (13)
 \end{aligned}$$

(12) and (13) constitute the member of a zero stable block integrators of order  $(5,5,5,5)^T$  with

$c_7 = \left( \frac{2351}{3936600}, \frac{7}{3600}, \frac{1}{600}, -\frac{143}{50400} \right)$ . The application of the block integrators with  $n = 0$  gives the accurate values of unknown as shown in tables I and II of forth section of this paper. To start the IVP integration on the sub interval  $[X_0, X_3]$ , we combine (12) and (13), when  $n = 0$  i.e the 1-block 4-point method as given in (14). Thus produces simultaneously values for  $y_1, y_2, y_3$  and  $y_{\frac{4}{3}}$  without recourse to any predictor like

Awoyemi [5] and Tafarki [8] to provide  $y_1$  and  $y_2$  in the main method. Hence, this is an improvement over their cited works. Though, this does not becloud the contribution of these authors.

III. STABILITY ANALYSIS

Recall, that, it is a desirable property for a numerical integrator to produce solution that behave similar to the theoretical solution to a problem at all times. Thus, several definitions, which call for the method to posses some “adequate” region of absolute stability, can be found in several literatures. See Lambert [3], Fatunla [1, 2] etc.

Following Fatunla [1, 2], the four integrator proposed in this report in (12) and (13) are put in the matrix equation form and for easy analysis the result was normalized to obtain;

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} y_{n+1} \\ y_{n+\frac{4}{3}} \\ y_{n+2} \\ y_{n+3} \end{bmatrix} = \begin{bmatrix} 0 & 0 & -\frac{1}{3} & 1 \\ 0 & 0 & -1 & 3 \\ 0 & 0 & -2 & 6 \\ 0 & 0 & -1 & 3 \end{bmatrix} \begin{bmatrix} y_{n-2} \\ y_{n-\frac{5}{3}} \\ y_{n-1} \\ y_n \end{bmatrix} + h^2 \left\{ \begin{bmatrix} -\frac{913}{5400} & \frac{523}{14580} & -\frac{313}{109350} & -\frac{14}{81} \\ \frac{81}{400} & \frac{19}{120} & -\frac{1}{100} & -1 \\ -\frac{243}{400} & \frac{49}{40} & \frac{4}{75} & -4 \\ \frac{567}{800} & -\frac{37}{240} & \frac{23}{1800} & \frac{1}{6} \end{bmatrix} \begin{bmatrix} f_{n+1} \\ f_{n+\frac{4}{3}} \\ f_{n+2} \\ f_{n+3} \end{bmatrix} + \begin{bmatrix} 0 & 0 & 0 & \frac{2027}{87480} \\ 0 & 0 & 0 & \frac{17}{240} \\ 0 & 0 & 0 & \frac{31}{240} \\ 0 & 0 & 0 & -\frac{65}{288} \end{bmatrix} \begin{bmatrix} f_{n-2} \\ f_{n-\frac{5}{3}} \\ f_{n-1} \\ f_n \end{bmatrix} \right. \quad (14)$$

with  $y_0 = \begin{pmatrix} y_0 \\ hz_0 \end{pmatrix}$  usually giving along the initial value problem. (14) is the 1- block 4 – point method. The first characteristics polynomial of the proposed 1- block 4 – point method is given

$$\text{by } \rho(\lambda) = \det [\lambda I - A_1^{(1)}] \quad (15)$$

$$\rho(\lambda) = \det \begin{bmatrix} \lambda & 0 & \frac{1}{3} & -1 \\ 0 & \lambda & 1 & -3 \\ 0 & 0 & \lambda + 2 & -6 \\ 0 & 0 & 1 & \lambda - 3 \end{bmatrix} \quad (16)$$

Solving the determinant of (13), yields  $\rho(\lambda) = \lambda^3(\lambda - 1)$ , which implies,  $\lambda_1 = \lambda_2 = \lambda_3 = 0$  or  $\lambda_4 = 1$

By definition of zero stable and (16), the 1 - block 4 - point method is zero stable and is also consistent as its order  $(5,5,5,5)^T > 1$ , thus, it is convergent following Henrici [10] and Fatunla [2].

IV. IMPLEMENTATION OF THE METHOD

This section deal with numerical experiments by considering the derived discrete schemes in block form for solution of stiff differential equations of second order initial value problems. The idea is to enable us see how the proposed methods performs when compared with exact solutions. The results are summarized in Table I & II.

A. Numerical Experiment

From Taparki and Odekunle [8]; Consider the IVP  $y'' = y, x \in [0,1], y_0 = 1, y'_0 = 1, h = 0.1$ , whose exact solution is  $y = e^x$

TABLE I  
RESULTS FOR THE PROPOSED METHOD

x	Exact Solution	Approximate Value	Error of Proposed Method	Taparki and Odekunle [8]
0.1	1.105170918	1.105170918	0.0000E+00	4.920292 x 10 <sup>-3</sup>
0.2	1.221402758	1.221402758	0.0000E+00	2.037513 x 10 <sup>-2</sup>
0.3	1.349858808	1.349858807	5.7600E-10	4.7477416 x 10 <sup>-2</sup>
0.4	1.491824698	1.491824696	1.6413E-09	8.7788535 x 10 <sup>-2</sup>
0.5	1.648721271	1.648721269	1.7001E-09	1.4235215 x 10 <sup>-1</sup>
0.6	1.822118800	1.822118798	2.3905E-09	2.1268728 x 10 <sup>-1</sup>
0.7	2.013752707	2.013752704	3.4705E-09	3.0047789 x 10 <sup>-1</sup>
0.8	2.225540928	2.225540924	4.4925E-09	4.07590071 x 10 <sup>-1</sup>
0.9	2.459603111	2.459603107	4.1569E-09	5.3609119 x 10 <sup>-1</sup>
1.0	2.718281828	2.718281824	4.4590E-09	6.88271136 x 10 <sup>-1</sup>

B. Numerical Experiment

From Adeboye [9]; Consider the BVP  $y'' - y = 4x - 5$ ;  $y(0) = y(1) = 0$ ,  $h = 0.1$ , whose exact solution is

$$y = \frac{7}{4(e^2 - e^{-2})} [e^{2x} - e^{-2x}] - \frac{3}{4}x$$

TABLE II  
RESULTS FOR THE PROPOSED METHOD

x	Exact Solution	Approximate Value	Error of Proposed Method	Adeboye [9]
0.0	0.0000000000	0.0000000000	0.000000000E+00	0.000000000E+00
0.1	0.14735784232	0.1473578284	1.390000000E-08	6.598600000E-06
0.2	0.25015214537	0.2501521164	2.890000000E-08	9.454000000E-06
0.3	0.31341504348	0.3134150000	4.340000000E-08	1.156300000E-05
0.4	0.34178302747	0.3417825591	4.680000000E-07	1.204180000E-05
0.5	0.33954334810	0.3395424500	8.981000000E-07	8.902600000E-06
0.6	0.31067692433	0.3106755871	1.337200000E-06	1.922800000E-06
0.7	0.25889818576	0.2588965200	1.665700000E-06	2.803580000E-05
0.8	0.18769224781	0.1876902363	2.011500000E-06	8.259870000E-05
0.9	0.10034979197	0.1003474152	2.376700000E-06	1.870490000E-04
1.0	0.0000000000	-0.0000023895	2.389500000E-06	3.379500000E-06

V. CONCLUSION

In this paper, a new block method with uniform integrators of order  $(5,5,5,5)^T$  was developed. The resultant numerical integrators possess the following desirable properties:

- i. Zero stability i.e. stability at the origin
- ii. Convergent schemes
- iii. An addition of equation from the use of first derivative
- iv. Being self – starting as such it eliminates the use of predictor – corrector method
- v. Facility to generate solutions at 4 points simultaneously
- vi. Produce solution over sub intervals that do not overlap
- vii. Apply uniformly to both IVP<sub>s</sub> and BVP<sub>s</sub> with adjustment to the boundary conditions

In addition, the new schemes compare favourably with the theoretical solution and the results are more accurate than Taparki and Odekunle [8], and Adeboye [9], see table I and II respectively. Hence, this work is an improvement over other cited works.

REFERENCES

[1] S.O. Fatunla, Block Method for Second Order Initial Value Problem. International Journal of Computer Mathematics, England. Vol. 4, 1991, pp 55 – 63.

[2] S.O. Fatunla, Higher Order parallel Methods for Second Order ODEs. Proceedings of the fifth international conference on scientific computing, 1994, pp 61 – 67.

[3] J.D.Lambert, Computational Methods in Ordinary Differential Equations (John Wiley and Sons, New York, USA, 1973).

[4] P. Onumanyi, D.O. Awoyemi, S.N. Jator and U.W. Sirisena, New linear Multistep with Continuous Coefficient for first order initial value problems. Journal of Mathematical Society, 13, 1994, pp 37 – 51.

[5] D.O. Awoyemi, A class of Continuous Stormer – Cowell Type Methods for Special Second Order Ordinary Differential Equations. Journal of Nigerian Mathematical Society Vol. 5, Nos. 1 & 2, 1998, pp100 – 108.

[6] Y.A. Yahaya and Z.A. Adegboye, A family of 4 – step Block Methods for Special Second Order in Ordinary Differential Equations. Proceedings Mathematical Association of Nigeria, 2008, pp 23 – 32.

[7] I. Fudziah L. K. Yap. and O. Mohammad, Explicit and Implicit 3 – point Block Methods for Solving Special Second Order Ordinary Differential Equations Directly. International Journal of math. Analysis, Vol. 3, 2009, pp 239 – 254.

[8] R. Taparki and M.R. Odekunle, An Implicit Runge – Kutta Method for Second Order Initial Value Problem in Ordinary Differential Equations. International Journal Num. Mathematics, Vol. 5 No. 2, 2010, pp 222 – 234.

[9] K.R. Adeboye, A superconvergent H2 – Galerkin method for the solution of Boundary Value Problems. Proceedings National Mathematical Centre, Abuja, Nigeria. Vol. 1 no.1, 2000, pp 118 – 124, ISBN 978 – 35488 – 0 – 2 .

[10] P. Henrici, Discrete Variable Methods for ODEs. (John Wiley New York U.S.A, 1962).