

Analysis of the Genetic Sequences of PCV2 Virus in Mexico

Robles F^A, Chevez J^A, Angulo R^B, Díaz E^B and González C^A.

Abstract—These All pig-producing countries from around the world report the presence of Postweaning multisystemic wasting syndrome (PMWS.) In America, PCV2 has been recognized in Canada, United States and Brazil. Knowledge concerning the genetic sequences of PMWS has been very important. In Mexico, there is no report describing the genetic sequences and variations of the PCV2 virus present around the country. For this reason, the main objective was to describe the homology and genetic sequences of the PCV2 virus obtained from different regions of Mexico. The results show that in Mexico are present both subgenotypes \"a\" and \"b\" of this virus and the homologies are from 89 to 99%. Regarding with the aminoacid sequence, three major heterogenic regions were present in the position 59-91, 123-136 and 185-210. This study presents the results of the first genetic characterization of PCV2 in production herds from Mexico.

Keywords—PCV-2; sequencing analysis; Mexico

I. INTRODUCTION

THE porcine circoviruses (PCV) are members of the genus *Circovirus*, family *Circoviridae*, it was originally isolated as a non-cytopathic contaminant of the porcine kidney cell line PK15, up to now no PCV infections has been consistently detected in the mammal species tested other than swine and mice [26]. These viruses are very small non-enveloped icosahedral viruses with a single stranded circular DNA genome of about 1.7 kb organized in an ambisense direction; into this family exist other 2 virus that are important for the Veterinary Medicine, chicken anemia virus and psittacine Beak and Feather Disease virus; also there are three plant virus and one that affects to human. [10], [8], [6], [30]. The PCV virus have at least six major Open Reading Frames (ORFs); and to two have been well recognized, ORF1, called the rep gene, which encodes a protein of 35.7 kDa involved in virus replication, and ORF2, called the cap gene, which encodes the major immunogenic capsid protein of 27.8 kDa [19], [12], [15], [20], [22].

Currently, two types of porcine circovirus have been described type 1 (PCV1) and type 2 (PCV2). In terms of the overall DNA sequence homology within the PCV1 or PCV2

isolates is greater than 90%, whereas the homology between PCV1 and PCV2 isolates is only 68 to 76% [5]. According with its importance, PCV1 is widespread, although it apparently caused no symptoms in pigs [8]. The other, porcine circovirus (PCV2) is considered to be an important emerging pathogen associated with a number of different syndromes and diseases in pigs as post-weaning multisystemic wasting syndrome (PWMS); dermatitis and nephropathy syndrome (PDNS), porcine respiratory disease complex, reproductive failure, granulomatous enteritis, exudative epidermitis, necrotizing lymphadenitis, congenital tremor [6] and actually has been described the important impact of adverse effect on the immune system in the form of compromising the development of a protective immune response to commercial vaccines commonly used on pigs around the time of PCV2 infection [23].

On the other hand, phylogenetic researches had divided PCV2 into two major genotypic groups: PCV2- group 1 (with three clusters A, B and C) and PCV2- group 2 with 5 clusters A, B, C, D and E); the principal differences between both are the size of the genome (group-1 1767 nucleotides and group-2 1768 nucleotides, the deleted nucleotide is located at nt1040) and a difference in aminoacid sequence that may serve as a signature motif [7]. Although the group-1 had been associated in all diseased animals have not been identified molecular markers specific to a pathogenic state [2], [10]. Recently, this classification has changed to PCV2-a and PCV2-b in base on a signature motif that encodes a distinct stretch of amino acid in the viral capsid protein [33]. Also third genotype (PCV2-c) was detected in Denmark with only three sequences to date [9], [13].

Even though the reason for the variety of clinical manifestations associated with PCV2 is not well understood, currently it has been shown that the ORF3 encodes a protein that is not essential for the replication *in vitro* but is important for the viremia process and is involved in the apoptosis (activating caspase-8 and caspase-3 pathways) induced by PCV2 [15], and there is a report that describe that B lymphocytes are more important that mononuclear cells for PCV2 replication [36].

Some studies on PCV2 and syndromes causing by it have been directed in order to obtain a better knowledge focused on the dynamics and behavior of PCV2 infections within farms. Major clinical signs evident are wasting and respiratory or digestive disorders, the low weight is evident at 7 and 12 weeks of age, and there is high mortality [29]. The

A. Authors are with Boehringer Ingelheim Vetmedica S.A. de C.V. Research and Development Department (corresponding author to provide phone: +52 3336687552; fax: +52 333668080; e-mail: francisco.robles@boehringer-ingelheim.com).

B. Authors are with Boehringer Ingelheim Vetmedica Inc. (corresponding author to provide phone: +1 (816) 236-2778; fax: +1 (816) 236-2778.

seroconversion pattern is equivalent to that of typical antibody dynamics for most viral agents that affect swine and both humoral immune response and cell-mediated immune response are required for full protection [21]; and has been established (by Real Time quantitative PCR) that the peak of the virus load is between 14 and 21 days PI [28]; as well is documented that the injection directly into tissue of SPF pigs of cloned PCV2 genomic DNA is capable of developing and infection and disease resembling those induced by infection via intranasal route, with gross and histopathological lesions in multiple tissues [11].

Nowadays researches around the world have developed molecular diagnostic techniques more sensible and sensitive in order to identify the herds infected [3], [14], [25], [4], [16], and based in serology the porcine circovirus type 2 has been described as a important virus in Europe (Germany, United Kingdom, Spain, France, Czech Republic, Sweden, and Denmark), Asia (China, Japan and Taiwan) and America (Canada, Costa Rica, Mexico, Brazil and the United States) with seroprevalences between 50% to 100% of positive farms [37], [1], [5], [33], [34], [17], [32]; additionally, in all these countries the knowledge of the genetic sequences have been very important in order to determinate the level of the homology between the different isolations and with this to learn more about of the disease. In Mexico pigs with antibodies against PCV2 was identified in 2001 [32], also it has been showed the presence of antibodies against PCV2 in 98% of backyard pigs from Mexico, this is indicative that PCV is present around all the Mexican territory [27]; but at own knowledge there is not a preview report that described the epidemiological and genetic behavior of PCV2 in Mexico, because of this the main objective of this study was to contribute to the understanding of the genetic characterization of PCV2 virus in Mexico.

II. MATERIALS AND METHODS

A. Collection and type of samples

This study was conducted in 34 swine farms distributed in 10 different regions of Mexico, which are important for its pig production; in 20 of these farms the pigs showed clinical PMWS besides that histopathological findings related with PMWS was founded in pigs from these farms. The sampling period was from January to September of 2006 collecting serum (n=3094), organs (n=122) including lymph nodes, lungs, ileum, semen (n=22), and fecal swaps (n=437); the samples were pooled according with the type of sample to a total of 888 pools (Table 2).

B. DNA isolation

The DNAeasy kit (QIAGEN) was used for the DNA extraction from the tissue and serum samples according with the manufacturer's instructions. For tissue 5-25 mg were weighed and incubated at 55°C over night with lysis buffer plus Proteinase K (20mg/ml), while each 200 µl of serum

samples was used for the process. For both, DNA was eluted with 60 µl buffer and 1 µl was used for the PCR reaction.

C. PCR and cloning of ORF2 fragment

A set of PCV2 specific primers (the sequence was kindly provided by Li Wang- VIDO University) was directed to amplify a fragment of 506 nucleotides. These primers amplify the sequences between the position 1570 - 1100 of the ORF2 (*cap* gene), the PCR was performed in a 25 µl reaction using 10- 20 ng template DNA, 10 µMol each primer, 0.5 Units of Taq polymerase Recombinant (INVITROGEN), 1.5 mM magnesium chloride, 0.2 mM each (10mM) dNTP's and the solution buffer recommended by the manufacture. The PCR thermocycling conditions were: initial denaturation 2 min at 94°C, then 35 cycles of denaturation 40 s 94°C, annealing 30 s 55°C, elongation 30s 72°C and finally one terminal extension of 72°C 10 min. The PCR products were analyzed by electrophoresis on a 1% agarose gel dye with Ethidium Bromide at 100 volts/ 45 min.

Ten positive samples (corresponding at different geographical regions) were purified with PCR Kit purify (INVITROGEN) and the amplified was cloned into TOPO TA cloning for sequencing (INVITROGEN) according with the manufacture instructions and the plasmidic DNA was extracted with MiniPrepKit SNAP (INVITROGEN), finally the samples were sequenced using an ABI PRISM 377 DNA sequencer (APPLIED). In order to get the percentages of sequences identity among different PCV2 sequences around the world; the sequences were compared pair-wise at both the nucleotide and amino acid levels using GENE BANK Blast program, multiple alignments and the phylogenetic tree were performed using the CLUSTAL W program using the sequence of the capsid protein (ORF2).

III. RESULTS

The number of samplings carried out in all these farms was 89 with 72 of the 89 samplings found to be positive (80.89%). The analysis for each the farms showed that 33 of 34 farms were PCV2 PCR positive; which is equal to 97.06%.

In addition, samples from each one of the regions were further characterized by sequencing. Ten distinct sequences were identified, and the results showed that the homologies of nucleotides from Mexican PCV2 virus are between 89% until 99%. The PCV2 sequence obtained in Guanajuato (MX LEON) differs most compared to the the other Mexican sequences (89 to 91%) just has 99% the homology with PCV2 from Queretaro (MX QRO) that is the second sequence with different homology (89 to 91%). The others sequences apparently are more closely related (it is showed in Table I).

TABLE I
GENETIC HOMOLOGY BETWEEN THE MEXICAN PCV2.

homo-logy, %	PCV2MX LEON 1	PCV2MX JAL 2	PCV2MX MICH 3	PCV2MX NL 4	PCV2MX PBC_SC 5	PCV2MX PBC_C 6	PCV2MX QRO 7	PCV2MX SON_N 8	PCV2MX SON_G 9
PCV2MX JAL 2	90 (4.57/506)								
PCV2MX MICH 3	89 (4.55/506)	90 (304/306)							
PCV2MX NL 4	90 (4.57/506)	90 (304/306)	99 (502/506)						
PCV2MX PBC_SC 5	91 (4.61/506)	94 (479/506)	94 (477/506)	94 (477/506)					
PCV2MX PBC_C 6	89 (4.55/506)	90 (304/306)	99 (502/506)	99 (502/506)	94 (477/506)				
PCV2MX QRO 7	99 (504/506)	90 (4.57/506)	89 (4.55/506)	90 (4.57/506)	91 (4.61/506)	89 (4.53/506)			
PCV2MX SON_N 8	89 (4.55/506)	90 (304/306)	99 (502/506)	99 (504/506)	94 (477/506)	90 (303/306)	89 (4.55/506)		
PCV2MX SON_G 9	90 (4.58/506)	90 (303/306)	99 (503/506)	99 (505/506)	94 (479/506)	90 (303/306)	90 (4.58/506)	99 (505/506)	
PCV2MX YUC II	90 (4.56/506)	90 (303/306)	99 (503/506)	99 (503/506)	94 (478/506)	90 (303/306)	90 (4.56/506)	99 (503/506)	99 (304/306)

The phylogenetic analysis (Figure 1) shows that the Mexican PCV2 isolates are divided in 3 principal clusters: the first has 2 sub-clusters that correspond to the Central and Southeast regions with 3 sequences; the second is formed only by one sequence of the Central Region; and the third cluster includes the majority of the sequences, and is sub-divided in 3 sub-clusters with a subcluster with two sequences from the North of Mexico; one more subcluster with just one sequence from the north of Mexico too; and one cluster with the 2 sequences that should the greatest heterologies obtained from the low laying area region.




Fig. 1 Phylogenetic tree of Mexican PCV2 isolates.

The analysis of the amino acids sequences shows that there was a range from 88% until 100% amino acid identify among Mexican PCV2 isolates. Again, the sequences obtained from Guanajuato y Queretaro were the most different compared to those obtained in the others regions (see Table II). When the Mexican virus amino acids sequences were compared with the PCV2 isolates recovered around the world (Figure 2), it was possible to identify that in Mexico there are virus of PVC2- a and PCV2- b; in the phylogenetic tree (Figure 3) the sequences Guanajuato (MX LEON) and Queretaro (MX QRO) are into the group a and the others belong to group b, this is the reason why there is a big differences. Also the majority of sequences are closely related with European virus, except for Puebla (MXPBC) that is more related with one virus from China and Yucatan (MXYUC) that is closely related with Canadian virus.

TABLE II
AMINO ACIDS HOMOLOGY BETWEEN THE MEXICAN PCV2 ISOLATES.

homo-logy, %	PCV2MX LEON 1	PCV2MX JAL 2	PCV2MX MICH 3	PCV2MX NL 4	PCV2MX PBC_SC 5	PCV2MX PBC_C 6	PCV2MX QRO 7	PCV2MX SON_N 8	PCV2MX SON_G 9
PCV2MX JAL 2	88								
PCV2MX MICH 3	88	100							
PCV2MX NL 4	88	100	100						
PCV2MX PBC_SC 5	89	94	94	94					
PCV2MX PBC_C 6	88	99	99	99	94				
PCV2MX QRO 7	100	88	88	88	89	88			
PCV2MX SON_N 8	88	100	100	100	94	99	88		
PCV2MX SON_G 9	88	99	99	99	94	98	88	99	
PCV2MX YUC II	88	100	100	100	94	99	88	100	99




Fig. 3 Phylogenetic tree of Mexican PCV2 isolates and others PCV2 virus around the world. In the red boxes are indicated the sequences from Mexico. Subgenotype A and Subgenotype B.

The analysis of the main changes between amino acid sequences of both PCV2 genogroups found in Mexico shows that there are 20 changes in the capsid protein; the most of these changes are in the domain region situated in the position from aa59 to aa91. In the table III is showed in detail.

The couple of Mexican sequences that are into genogroup a do not show changes in the amino acid sequence and their sequences are closely related with European sequences. Regarding genogroup b, the sequences obtained from Puebla region show change from R₆₃ to K₆₃ in PCV2MX PB_C; and

MOST FREQUENT CHANGES OF AMINO ACIDS IN PCV2 SEQUENCES FROM MEXICO

Sample	amino acid position																				
	69	62	72	76	77	86	88	90	93	91	121	122	131	136	141	156	160	191	204	210	
PCV232QXQRO	A	S	L	L	D	V	T	K	I	I	S	I	I	Q	P	M	S	R	K	D	Genogroup 2
PCV232XQRO	A	S	L	L	D	V	T	K	I	I	S	I	I	Q	P	M	S	R	K	D	
PCV232AL	R	R	M	I	N	L	S	P	R	V	S	V	T	L	T	L	A	G	I	E	Genogroup 3
PCV232Q22H	R	R	M	I	N	L	S	P	R	V	S	V	T	L	T	L	A	G	I	E	
PCV232QL	R	R	M	I	N	L	S	P	R	V	S	V	T	L	T	L	A	G	I	E	
PCV232XQ23N	R	R	M	I	N	L	S	P	R	V	S	V	T	L	T	L	A	G	I	E	
PCV232XQ2VC	R	R	M	I	N	L	S	P	R	V	S	V	T	L	T	L	A	G	I	E	
PCV232XQ2Q	R	R	M	I	N	L	S	P	R	V	S	V	T	L	T	L	A	G	I	E	
PCV232Q22C	R	K	M	I	N	L	S	P	R	V	S	V	T	L	T	L	A	G	I	E	
PCV232Q22B2_C	R	K	M	I	N	L	S	P	R	V	S	V	T	L	T	L	A	G	I	E	

Other important observation is that the majority of Mexican virus sequences are closely related with European PCV2s, for the PCV2- group a with Austria, Germany, France, Hungarian

In conclusion, these results have demonstrated that both subgenotypes of PCV2 are present in at least in 90% of the principal zones for swine industry in Mexico and also has contributed to the understanding of PCV2 epidemiology in Mexico. Is important to conduct future research in order to understand the economic impact of this pathogen in the Mexican swine industry and demonstrate if one subgenotype is more pathogenic than other (as has been reported recently in USA and Canada by Cheung *et al.*, 2007). One more time it was confirmed that PCV2 virus is present around the entire world.

- [1] C. Boisseson, V. Beven, L. Bigarre, R. Thiery, N. Rose, E. Eveno, F. Madec and A. Jestin. "Epidémiologie moléculaire de circovirus porcin (PCV2) en relation avec la maladie d'amairgissement du porcelet". *Journées Recherche Porcine*. 36, pp. 333-338. 2004.
- [2] C. Boisseson, V. Beven, L. Bigarre, R. Thiery, N. Rose, E. Eveno, Madec, F. and Jestin, A. "Molecular characterization of porcine circovirus type 2 isolates from post-weaning multisystemic wasting syndrome-affected and non-affected pigs". *J. Gen. Virol.* 85, pp. 293-304. 2004.
- [3] I. Brunborg, T. Modal and C. Jonassen. "Quantification of porcine circovirus type 2 isolated from serum/plasma and tissue samples of healthy pigs and pigs with postweaning multisystemic wasting syndrome using a TaqMan-based real-time PCR". *J. Viro. Met.* 122, pp. 171-178. 2004.
- [4] S. Cao, H. Chen, J. Zhao, L. Lu, S. Xiao, A. Guo, B. Wu and Q. He. "Detection of porcine circovirus type 2, porcine parvovirus and porcine pseudorabies virus from pigs with postweaning multisystemic wasting syndrome by multiplex PCR". *Vet. Res. Commun.* 29, pp. 263-269. 2005.
- [5] V. Celer, and P. Carasova. "First evidence of porcine circovirus type 2 (PCV2) infection of pig in the Czech Republic by semi-nested PCR". *J. Vet. Med.* 49, pp. 155-159. 2002.

- [6] C. Chae. "A review of porcine circovirus 2-associated syndromes and diseases". *Vet. J.* 169, pp. 326-336. 2005.
- [7] A. Cheung, K. Lager, K. Kohutnyuk, A. Vicent, S. Henry, R. Baker, R. Rowland and A. Dunham. "Detection of two porcine circovirus type 2 genotypic groups in United States swine herds". *Arch. Virol.* 152, pp. 1035-1044. 2007.
- [8] A. Crowther, J. Berriman, W. Curran, M. Allan and D. Todd. "Comparison of the structures of three Circoviruses: Chicken anemia virus, Porcine Circovirus type 2, and Beak and feather disease virus". *J. Virol.* 77, pp. 13036-13041. 2003.
- [9] K. Dupont, E. Nielsen, P. Bækbo and L. Larsen. "Genomic analysis of PCV2 isolates from Danish archives and a current PMWS case-control study supports a shift in genotypes with time". *Vet. Microbiology.* 128, pp. 56-64. 2008.
- [10] M. Fenaux, P. Halbur, M. Gill, T. Toth and X. Meng. "Genetic characterization of type 2 porcine Circovirus (PCV-2) from pigs with Postweaning Multisystemic Wasting Syndrome in different geographic regions of north American and development of a different PCR-Restriction Fragment Length Polymorphism assay to detect and differentiate between infections with PCV-1 and PCV-2". *J. Clin. Microbiol.* 38, pp. 2494-2503. 2000.
- [11] M. Fenaux, P. Halbur, G. Haqshenas, R. Royer, P. Thomas, P. Nawagitgul, M. Gill, T. Toth and P. Meng. "Cloned genomic DNA of type 2 porcine circovirus is infectious when injected directly into the liver and lymph nodes of pigs: characterization of clinical disease, virus distribution, and pathologic lesions". *J. Virol.* 76, pp. 541-551. 2002.
- [12] M. Fenaux, T. Opriessning, P. Halbur and X. Meng. "Immunogenicity and pathogenicity of chimeric infectious clones of pathogenic porcine circovirus type 2 (PCV2) and nonpathogenic PCV1 in weanling pigs". *J. Virol.* 77, pp. 11232-1143. 2003.
- [13] L. Grau-Roma, E. Crisci, M. Sibila, S. Lopez-Soria, M. Nofrarias, M. Cortey, L. Fraile, A. Olvera and J. Segales. "A proposal on porcine circovirus type 2 (PCV2) genotype definition and their relation with postweaning multisystemic wasting syndrome (PMWS) occurrence". *Vet. Microbiology.* 128, pp. 23-35. 2008.
- [14] J. Kim and C. Chae. "Multiplex nested PCR compared with in situ hybridization for the differentiation of porcine circovirus and porcine parvovirus from pigs with postweaning multisystemic wasting syndrome". *Can. J. Vet. Res.* 67, pp. 133-137. 2003.
- [15] J. Liu, I. Chen, Q. Du, H. Chua and J. Kwang. "The ORF3 protein of porcine circovirus type 2 is involved in viral pathogenesis in vivo". *J. Virol.* 80, pp. 5065-5073. 2006.
- [16] Q. Liu, L. Wang, P. Willson and L. Babiuk. "Quantitative, competitive PCR analysis of porcine circovirus DNA in serum from pigs with postweaning multisystemic wasting syndrome". *J. Clin. Microbiol.* 38, pp. 3474-3477. 2000.
- [17] Liu, Q.; Wang, L.; Willson, P.; O'Connor, B.; Keenlside, J.; Chirini-Trejo, M.; Melendez, R. and Babiuk, L. 2002. Seroprevalence of porcine circovirus type 2 in swine populations in Canada and Costa Rica. *Can. J. Vet. Res.* 66: 225-231.
- [18] D. Mahé, P. Blanchard, C. Truong, C. Claire Arnauld, P. Le-Cann, R. Cariolet, F. Madec, E. Albina, and A. Jestin. "Differential recognition of ORF2 protein from type 1 and type 2 porcine circoviruses and identification of immunorelevant epitopes". *J. Gen. Virol.* 81, pp. 1815-1824. 2000.
- [19] A. Mankertz and B. Hillenbrand. "Analysis of transcription of porcine circovirus type 1". *J. Gen. Virol.* 83, pp. 2743-2751. 2002.
- [20] A. Mankertz, J. Mankertz, K. Wolf and J. Buhk. "Identification of a protein essential for replication of porcine circovirus". *J. Gen. Virol.* 79, pp. 381-383. 1998.
- [21] N. McKeown, T. Opriessning, P. Thomas, D. Guenette, F. Elvinger, M. Fenaux, P. Halbur and X. Meng. "Effects of porcine circovirus type 2 (PCV2) maternal antibodies on experimental infection of piglets with PCV2". *Clin. Diagn. Lab. Immunol.* 12, pp. 1347-1352. 2005.
- [22] P. Nawagitgul, R. Morozov, R. Bolin, P. Harms and D. Sorden. "Open Reading Fragment 2 of porcine circovirus type 2 encodes a major capsid protein". *J. Gen. Virol.* 81, pp. 2281-2287. 2000.
- [23] T. Opriessning, N. McKeown, K. Harmon, X. Meng and P. Halbur. "Porcine circovirus type 2 infection decreases the efficacy of modified live porcine reproductive and respiratory syndrome virus vaccine". *Clin. Vaccine Immunol.* 1, pp. 923-929. 2006.
- [24] T. Opriessnig, S. Ramamoorthy, D. Madson, A. Patterson, N. Pal, S. Carman, X. Meng and P. Halbur. "Differences in virulence among porcine circovirus type 2 isolates are unrelated to cluster type 2a or 2b and prior infection provides heterologous protection". *J. Gen. Virol.* 89, pp. 2482-2491. 2008.
- [25] M. Ouardani, L. Wilson, R. Jette, C. Montpetit and S. Dea. "Multiplex PCR for detection and typing of porcine circoviruses". *J. Clin. Microbiol.* 37, pp. 3917-3924. 1999.
- [26] J. Quintana, M. Balasch, J. Segales, M. Calsamiglia, G. Rodriguez-Arriola, J. Plana-Duran and M. Domingo. "Experimental inoculation of porcine circoviruses type 1 (PCV1) and type 2 (PCV2) in rabbits and mice". *Vet. Res.* 33, pp. 229-237. 2002.
- [27] H. Ramirez-Mendoza, C. Martinez; C. Mercado, H. Castillo-Juarez, J. Hernandez and J. Segales. "Porcine circovirus type 2 antibody detection in backyard pigs from Mexico city". *Res. Vet. Sci.* 83, pp. 130-132. 2007.
- [28] A. Resendes, J. Segales, M. Balasch, M. Calsamiglia, M. Sibila, H. Ilerbrok, E. Mateu, J. Plana-Duran, A. Mankertz, and M. Domingo. "Lack of an effect of a commercial vaccine adjuvant on the development of postweaning multisystemic wasting syndrome (PMWS) in porcine circovirus type 2 (PCV2) experimentally infected conventional pigs". *Vet. Res.* 35, pp. 83-90. 2004.
- [29] G. Rodriguez-Arriola, J. Segales, M. Calsamiglia, A. Resendes, M. Balasch, J. Plana-Duran, J. Casal, and M. Domingo. "Dynamics of porcine circovirus type 2 infection in herd of pigs with postweaning and multisystemic wasting syndrome". *AJVR.* 6, pp.: 354-357. 2002.
- [30] S. Staebler, T. Sydler, E. Buergi, K. McCullough, F. McNeilly, G. Allan, and A. Pospischil. "PMWS: an emerging disease identified in archived porcine tissues". *Vet. J.* 170, pp. 132-134. 2005.
- [31] S. Timmusk, P. Wallgren, I. Marit, F. Hasslung, G. Allan, B. Meehan, M. McMenamy, F. McNeilly, L. Fuxler, K. Belak, D. Podersoo, T. Saar, M. Berg and C. Fossum. "Phylogenetic analysis of porcine circovirus type 2 (PCV2) pre- and post-epizootic postweaning multisystemic wasting syndrome (PMWS)". *Virus Genes.* 36, pp. 509-520. 2008.
- [32] M. Trujano, G. Iglesias, J. Segales and J. Palacios. "PCV2 from emaciated pigs in Mexico". *Vet. Record* 148, pp.792. 2001.
- [33] J. Vicente, J. Segales, U. Hofle, M. Balasch, J. Plana-Duran, M. Domingo and C. Gortazar. "Epidemiological study on porcine circovirus type 2 (PCV2) infection in the European wild boar (*Sus scrofa*)". *Vet. Res.* 35, pp. 243-253. 2004.
- [34] P. Wallgren, F. Hasslung, G. Bergstrom, A. Linder, K. Belak, C. Hard, M. Stampe; B. Molander, T. Bjornberg, E. Norregard, C. Ehlorsson, M. Tornquist, C. Fossum, G. Allan and J. Robertsson. "Postweaning multisystemic wasting syndrome -PMWS. The first year with the disease in Sweden". *Vet. Quarterly.* 26, pp. 170-187. 2004.
- [35] D. Wiederkehr, T. Sydler, E. Buergi, M. Haessig, D. Zimmermann, A. Pospischil, E. Brugnera and X. Sidler. "A new emerging genotype subgroup within PCV-2b dominates the PMWS epizooty in Switzerland". *Vet. Microbiology.* 136, pp. 27-35. 2009.
- [36] S. Yu, T. Opriessnig, P. Kitikoon, D. Nilubol, P. Halbur and E. Thacker. "Porcine circovirus type 2 (PCV2) distribution and replication in tissue and immune cells in early infected pigs". *Vet. Immuno. Immunopathol.* 115, pp. 261-272. 2007.
- [37] J. Zhou, Q. Chen, J. Ye, H. Shen, T. Chen and S. Shang. "Serological investigation and genomic characterization of PCV2 isolates from different geographical regions of Zhejiang province in China". *Vet. Res. Commun.* 3, pp. 205-220. 2006.