

Best Coapproximation in Fuzzy Anti- n -Normed Spaces

J. Kavikumar, N. S. Manian, M. B. K. Moorthy

Abstract—The main purpose of this paper is to consider the new kind of approximation which is called as t -best coapproximation in fuzzy n -normed spaces. The set of all t -best coapproximation define the t -coproximal, t -co-Chebyshev and F -best coapproximation and then prove several theorems pertaining to this sets.

Keywords—Fuzzy- n -normed space, best coapproximation, co-proximal, co-Chebyshev, F -best coapproximation, orthogonality

I. INTRODUCTION

THE concept of best coapproximation was introduced by Franchetti and Furi [2], in order to study some characteristic properties of real Hilbert spaces, and such problems were considered further by Papini and Singer, [12] and Rao and Saravanan [13]. The concept of n -norm on a linear space has been introduced and developed by Gähler in [3], [4]. Following Misiak [10], Malčeski [9] and Gunawan and Mashadi [5] developed the theory of n -normed space. The concept of fuzzy norm was initiated by Katsaras in [7] and further, Narayanan and Vijayabalaji [11] introduced the concept of fuzzy n -normed linear space. Moreover, Vijayabalaji and Thillaigovindan [17] introduced the notion of convergent sequence and Cauchy sequence in fuzzy n -normed linear space. In [6] Iqbal H. Jebril and Samanta introduced fuzzy anti-norm on a linear space depending on the idea of fuzzy anti-norm was introduced by Bag and Samanta [1] and investigated their important properties. In [8] Kavikumar et. al. introduced the notion of fuzzy anti- n -normed linear space. Further, Surender Reddy [15] introduced the notion of convergent sequence and Cauchy sequence in fuzzy anti- n -normed linear space. The set of all t -best approximations on fuzzy normed linear spaces was initiated and studied by Vaezpour and Karimi [16]. The set of all t -best approximations on fuzzy anti- n -normed linear space was introduced in [14]. In this paper we consider the set of all t -best coapproximation in fuzzy anti- n -normed spaces and then prove several theorems pertaining to this set.

II. PRELIMINARIES

Definition 1: [17]. Let $n \in \mathbb{N}$ (natural numbers) and X be a real linear space of dimension $d \geq n$. (Here we allow d to be infinite). A real valued function $\|\bullet, \bullet, \dots, \bullet\|$ on $X \times X \times \dots \times X$ (n times) $= X^n$ satisfying the following four properties:

- $\|x_1, x_2, \dots, x_n\| = 0$ if and only if x_1, x_2, \dots, x_n are linearly dependent.

J. Kavikumar is with the Department of Mathematics and Statistics, Faculty of Science, Technology and Human Development, Universiti Tun Hussein Onn Malaysia, 86400 Parit Raja, Johor, Malaysia, (e-mail:kavi@uthm.edu.my).

N. S. Manian is with the Department of Mathematics, Angel College of Engineering and Technology, Tirupur 641665, Tamilnadu, India.

M. B. K. Moorthy is with the Department of Mathematics, Institute of Road and Transport, Erode, 638316, Tamilnadu, India.

- $\|x_1, x_2, \dots, x_n\|$ is invariant under any permutation of x_1, x_2, \dots, x_n .
- $\|x_1, x_2, \dots, cx_n\| = |c| \|x_1, x_2, \dots, x_n\|$, for any real c .
- $\|x_1, x_2, \dots, x_{n-1}, y + z\| \leq \|x_1, x_2, \dots, x_{n-1}, y\| + \|x_1, x_2, \dots, x_{n-1}, z\|$

is called an n -norm on X and the pair $(X, \|\bullet, \dots, \bullet\|)$ is called an n -normed linear space.

Definition 2: [17]. Let X be a linear space over a real field \mathbb{F} . A fuzzy subset N of $X^n \times [0, \infty)$ is called a fuzzy n -norm on X if and only if:

- $N(x_1, x_2, \dots, x_n, t) > 0$.
- $N(x_1, x_2, \dots, x_n, t) = 1 \Leftrightarrow x_1, x_2, \dots, x_n$ are linearly dependent.
- $N(x_1, x_2, \dots, x_n, t)$ is invariant under any permutation of x_1, x_2, \dots, x_n .
- $N(x_1, x_2, \dots, cx_n, t) = N(x_1, x_2, \dots, x_n, \frac{t}{|c|})$ if $c \neq 0, c \in \mathbb{F}(\text{field})$
- $N(x_1, x_2, \dots, x_n + x'_n, s + t) \geq N(x_1, x_2, \dots, x_n, t) * N(x_1, x_2, \dots, x'_n, s)$ for all $s, t \in \mathbb{R}$
- $N(x_1, x_2, \dots, x_n, \cdot)$ is left continuous and non-decreasing function of \mathbb{R} such that $\lim_{t \rightarrow \infty} N(x_1, x_2, \dots, x_n, t) = 1$.

Then (X, N) is called a fuzzy n -normed linear space.

Definition 3: [8] Let X be a linear space over a real field \mathbb{F} . A fuzzy subset N of $X^n \times [0, \infty)$ is called a fuzzy anti n -norm on X if and only if:

- for all $t \in \mathbb{R}$ with $t \leq 0$, $N(x_1, x_2, \dots, x_n, t) = 1$.
- for all $t \in \mathbb{R}$ with $t > 0$, $N(x_1, x_2, \dots, x_n, t) = 0 \Leftrightarrow x_1, x_2, \dots, x_n$ are linearly dependent.
- $N(x_1, x_2, \dots, x_n, t)$ is invariant under any permutation of x_1, x_2, \dots, x_n .
- for all $t \in \mathbb{R}$ with $t > 0$, $N(x_1, x_2, \dots, cx_n, t) = N(x_1, x_2, \dots, x_n, \frac{t}{|c|})$ if $c \neq 0, c \in \mathbb{F}(\text{field})$
- for all $s, t \in \mathbb{R}$, $N(x_1, x_2, \dots, x_n + x'_n, s + t) \leq \max\{N(x_1, x_2, \dots, x_n, s), N^*(x_1, x_2, \dots, x'_n, t)\}$
- $N(x_1, x_2, \dots, x_n, \cdot)$ is right continuous and non-increasing function of \mathbb{R} such that

$$\lim_{t \rightarrow \infty} N(x_1, x_2, \dots, x_n) = 0$$

Then (X, N) is called a fuzzy anti n -normed linear space.

To strengthen the above definition, we present the following example.

Example 1: [8] Let $(X, \|\bullet, \bullet, \dots, \bullet\|)$ be a n -normed linear space

Define,

$$N(x_1, x_2, \dots, x_n, t) =$$

$$\begin{cases} 1 - \frac{t}{t + \|x_1, x_2, \dots, x_n\|} & \text{when } t(> 0) \in \mathbb{R}, \forall x \in X \\ 1 & \text{when } t(\leq 0) \in \mathbb{R}, \forall x \in X \end{cases}$$

Then (X, N) is a fuzzy anti n -normed linear space.

Definition 4: [15] A sequence $\{x_k\}$ in a fuzzy anti- n -normed linear space (X, N) is said to be convergent to $x \in X$ if given $t > 0, 0 < r < 1$, there exists an integer $n_0 \in \mathbb{N}$ such that

$$N(x_1, x_2, \dots, x_{n_1}, x_k - x, t) < r, \forall k \geq n_0.$$

Theorem 1: [15] In a fuzzy anti- n -normed linear space (X, N) , a sequence $\{x_k\}$ converges to $x \in X$ if and only if

$$\lim_{k \rightarrow \infty} N(x_1, x_2, \dots, x_{n_1}, x_k - x, t) = 0, \forall t > 0.$$

Definition 5: [15] Let (X, N) be a fuzzy anti- n -normed linear space. Let $\{x_k\}$ be a sequence in X then $\{x_k\}$ is said to be a Cauchy sequence if

$$\lim_{k \rightarrow \infty} N(x_1, x_2, \dots, x_{n_1}, x_{k+p} - x_k, t) = 0, \forall t > 0$$

and $p = 1, 2, 3, \dots$. A fuzzy anti- n -normed linear space (X, N) is said to be complete if every Cauchy sequence in X is convergent. A complete fuzzy anti- n -normed space (X, N) is called a fuzzy anti- n -Banach space. The open ball $B(x, r, t)$ and the closed ball $B[x, r, t]$ with the center $x \in X$ and radius $0 < r < 1, t > 0$ are defined as follows:

$$B(x, r, t) = \{y \in X : N(x_1, x_2, \dots, x_{n_1}, x - y, t) < r\},$$

$$B[x, r, t] = \{y \in X : N(x_1, x_2, \dots, x_{n_1}, x - y, t) \leq r\}.$$

A subset A of X is said to be open if there exists $r \in (0, 1)$ such that $B(x, r, t) \subset A$ for all $x \in A$ and $t > 0$. A subset A of X is said to be closed if for any sequence $\{x_k\}$ in A converges to $x \in A$. i.e., $\lim_{k \rightarrow \infty} N(x_1, x_2, \dots, x_{n_1}, x_k - x, t) = 0$, for all $t > 0$ implies that $x \in A$.

Corollary 1: [15] Let (X, N) be a fuzzy anti- n -normed linear space. Then N is a continuous function on

$$\underbrace{X \times X \times \dots \times X}_n \times \mathbb{R}.$$

III. T-BEST COAPPROXIMATION

Definition 6: Let A be a nonempty subset of fuzzy anti- n -normed space (X, N) and $t > 0$. For $x \in X$, an element $y_0 \in A$ is said to be a t -best coapproximation of x from A if $N(x_1, x_2, \dots, x_{n-1}, y_0 - y, t) \leq N(x_1, x_2, \dots, x_{n-1}, x - y, t)$, for all $y \in A$. The set of all elements of t -best coapproximation of x from A is denoted by $R_A^t(x)$; i.e., $R_A^t(x) = \{y_0 \in A : N(x_1, x_2, \dots, x_{n-1}, y_0 - y, t) \leq N(x_1, x_2, \dots, x_{n-1}, x - y, t), \forall y \in A\}$.

For $t > 0$ putting

$$\begin{aligned} \check{A}_x^t &= \{x \in X; N(x_1, x_2, \dots, x_{n-1}, y, t) \\ &\leq N(x_1, x_2, \dots, x_{n-1}, y - x, t) \forall y \in A\} \\ &= (R_A^t)^{-1}(\{0\}). \end{aligned}$$

It is clear $y_0 \in R_A^t(x)$ if and only if $x - y_0 \in \check{A}_x^t$.

Definition 7: Let A be a nonempty subset of a fuzzy anti- n -normed space (X, N) . If for $t > 0$ and each $x \in X$ has at least (respectively exactly) one t -best coapproximation in A , then A is called a t -best coproximal (respectively t -co-Chebyshev) set. Also A is called t -quasi-co-Chebyshev set if $R_A^t(x)$ is a compact set.

Theorem 2: Let (X, N) be a fuzzy anti- n -normed space and A be a subspace of X and $t > 0$. Then for each $x \in X$

- (a) A is a t -coproximal if and only if $X = A + \check{A}_x^t$.
- (b) A is a t -co-Chebyshev subspace if and only if $X = A \oplus \check{A}_x^t$.

Proof: (a)(\Rightarrow) Assume that A is t -coproximal, $x \in X$ and $y_0 \in R_A^t(x)$. Then, $x - y_0 \in \check{A}_x^t$. Now, $x = y_0 + (x - y_0) \in A + \check{A}_x^t$. Hence $X = A + \check{A}_x^t$.

(\Leftarrow) Let $x \in X = A + \check{A}_x^t$. $x = y_0 + \bar{y}$, $y_0 \in A$, $\bar{y} \in \check{A}_x^t$ and so $0 \in R_A^t(\bar{y}) = R_A^t(x - y_0)$. Since, $N(x_1, x_2, \dots, x_{n-1}, 0 - (x - y_0), t) \leq N(x_1, x_2, \dots, x_{n-1}, y - (x - y_0), t)$, so $N(x_1, x_2, \dots, x_{n-1}, y_0 - x, t) \leq N(x_1, x_2, \dots, x_{n-1}, (y + y_0) - x, t)$ where $y + y_0 \in A$; hence $y_0 \in R_A^t(x)$. Therefore A is t -coproximal.

(b)(\Rightarrow) Suppose that A is t -co-Chebyshev subspace, $x \in X$, and $x = y_1 + \bar{y}_1 = y_2 + \bar{y}_2$, where $y_1, y_2 \in A$ and $\bar{y}_1, \bar{y}_2 \in \check{A}_x^t$. We show that $y_1 = y_2$ and $\bar{y}_1 = \bar{y}_2$. Since $x = y_1 + \bar{y}_1 = y_2 + \bar{y}_2$, then $x - y_1 = \bar{y}_1, x - y_2 = \bar{y}_2$, this implies that $y_1, y_2 \in R_A^t(x)$. Therefore $y_1 = y_2$, it follows that $\bar{y}_1 = \bar{y}_2$. Thus $X = A \oplus \check{A}_x^t$.

(\Leftarrow) Let $X = A \oplus \check{A}_x^t$, and suppose for $x \in X$, there exist $y_1, y_2 \in R_A^t(x)$. Then $x - y_1, x - y_2 \in \check{A}_x^t$ and therefore, $x = y_1 + \bar{y}_1 = y_2 + \bar{y}_2$, where $\bar{y}_1 = x - y_1, \bar{y}_2 = x - y_2$. It follows that $y_1 = y_2$ and $\bar{y}_1 = \bar{y}_2$. ■

Theorem 3: Let A be a nonempty subset of a fuzzy anti- n -normed space (X, N) . The for $t > 0$ and for each $x \in X$.

- (a) $R_{A+y}^t(x+y) = R_A^t(x) + y$, for every $x, y \in X$.
- (b) $R_{\alpha A}^{|\alpha|t}(\alpha x) = \alpha R_A^t(x)$ for every $x \in X$ and $\alpha \in \mathbb{R} \setminus \{0\}$.
- (c) A is t -coproximal (respectively t -co-Chebyshev) if and only if $A + y$ is t -coproximal (respectively t -co-Chebyshev), for any $y \in X$.
- (d) A is t -coproximal (respectively t -co-Chebyshev) if and only if αA is $|\alpha|t$ -coproximal (respectively $|\alpha|t$ -co-Chebyshev), for any $y \in X$, for any given $\alpha \in \mathbb{R} \setminus \{0\}$.

Proof: (i) For any $x, y \in X, t > 0, y_0 \in R_{A+y}^t(x+y)$ if and only if

$$N(x_1, x_2, \dots, x_{n-1}, y_0 - (a + y), t) \leq N(x_1, x_2, \dots, x_{n-1}, x + y - (a + y), t) \text{ for all } (a + y) \in A + y$$

$$N(x_1, x_2, \dots, x_{n-1}, (y_0 - y) - a, t) \leq N(x_1, x_2, \dots, x_{n-1}, x - a, t) \text{ for all } a \in A, \text{ if and only if, } (y_0 - y) \in R_A^t(x), \text{ i.e., } y_0 \in R_A^t(x) + y.$$

(ii) For any $x \in X, \alpha \in \mathbb{R} \setminus \{0\}$, and $t > 0, y_0 \in R_{\alpha A}^{|\alpha|t}(\alpha x)$ if and only if,

$$N(x_1, x_2, \dots, x_{n-1}, (y_0 - \alpha a, |\alpha|t) \leq N(x_1, x_2, \dots, x_{n-1}, \alpha x - \alpha a, \alpha t) \text{ for all } a \in A$$

$$\text{if and only if } N(x_1, x_2, \dots, x_{n-1}, (\frac{1}{\alpha} y_0 - a, |t|) \leq N(x_1, x_2, \dots, x_{n-1}, x - a, t) \text{ for all } a \in A \text{ if and only if, } \frac{1}{\alpha} y_0 \in R_A^t(x) \text{ if and only if, } y_0 \in \alpha R_A^t(x). \text{ Therefore } R_{\alpha A}^{|\alpha|t}(\alpha x) = \alpha R_A^t(x)$$

(iii) Is an immediate consequence of (i)

(iv) Is an immediate consequence of (ii). ■

Corollary 2: Let M be a nonempty subspace of a fuzzy anti- n -normed space X . Then for $t > 0$ and each $x \in X$.

- (a) $R_M^t(x+y) = R_M^t(x) + y$, for every $x, y \in X$.
- (b) $R_M^{|\alpha|t}(\alpha x) = \alpha R_M^t(x)$ for every $x \in X$ and $\alpha \in \mathbb{R} \setminus \{0\}$.

Proof: The proof is an immediate consequence of theorem 3 and this fact that $M + y = M$ and $\alpha M = M$ for all $y \in M$ and $\alpha \in \mathbb{R} \setminus \{0\}$. ■

Definition 8: For $x \in X, a \in A, 0 < r < 1$, and $t > 0$, define

$$e_a^t(x) = N(x_1, x_2, \dots, x_{n-1}, x - a, t)$$

Theorem 4: Let (X, N) be a fuzzy anti n -normed space, A be a subset of X , $x \in X \setminus \bar{A}$ and $t > 0$. Then we have

$$R_A^t(x) = \left[\bigcap_{a \in A} B[a, e_a^t(x), t] \right] \cap A.$$

Proof: By definition of $R_A^t(x)$ for each $a \in A$ we have

$$R_A^t(x) \subseteq [B[a, e_a^t(x), t] \cap A]$$

Therefore

$$R_A^t(x) \subseteq \left[\bigcap_{a \in A} B[a, e_a^t(x), t] \right] \cap A.$$

Conversely, let $y \in \left[\bigcap_{a \in A} B[a, e_a^t(x), t] \right] \cap A$, then we have $y \in A$, and for each $a \in A$, $N(x_1, x_2, \dots, x_{n-1}, a - y, t) \leq e_a^t = N(x_1, x_2, \dots, x_{n-1}, x - a, t)$, which implies that $y \in R_A^t(x)$. So $\left[\bigcap_{a \in A} B[a, e_a^t(x), t] \right] \cap A \subseteq R_A^t(x)$, which completes the proof. ■

Corollary 3: Let (X, N) be a fuzzy anti- n -normed space, A be a subset of X , $x \in X \setminus \bar{A}$ and $t > 0$. Then

- (a) The set $R_A^t(x)$ is t -bounded.
- (b) If A is t -closed, then $R_A^t(x)$ is t -closed.

Theorem 5: Let (X, N) be a fuzzy anti- n -normed space. For each $x \in X$ and $t > 0$, if A is a convex subset of X , then $R_A^t(x)$ is a convex subset of A (for $R_A^t(x) \neq \emptyset$).

Proof: Let $z_1, z_2 \in R_A^t$, then for $t > 0$ and each $x \in X$, $N(x_1, x_2, \dots, x_{n-1}, y - z_1, t) \leq N(x_1, x_2, \dots, x_{n-1}, x - y, t)$ and $N(x_1, x_2, \dots, x_{n-1}, y - z_2, t) \leq N(x_1, x_2, \dots, x_{n-1}, x - y, t)$ for all $y \in A$. Now for each $\lambda \in (0, 1)$ we have

$$\begin{aligned} & N(x_1, x_2, \dots, x_{n-1}, y - (\lambda z_1 + (1 - \lambda)z_2), t) \\ &= N(x_1, x_2, \dots, x_{n-1}, \lambda y - \lambda z_1 + y - \lambda y - z_2 + \lambda z_2, t) \\ &= N(x_1, x_2, \dots, x_{n-1}, \lambda(y - z_1) + (1 - \lambda)(y - z_2), \\ &\quad \lambda t + (1 - \lambda)t) \\ &\leq \max\{N(x - 1, x_2, \dots, x_{n-1}, y - z_1, \frac{\lambda t}{\lambda}), \\ &\quad N(x_1, x_2, \dots, x_{n-1}, y - z_2, \frac{(1 - \lambda)t}{(1 - \lambda)})\} \\ &\leq \max\{N(x - 1, x_2, \dots, x_{n-1}, x - y, \frac{\lambda t}{\lambda}), \\ &\quad N(x_1, x_2, \dots, x_{n-1}, x - y, \frac{(1 - \lambda)t}{(1 - \lambda)})\} \\ &\leq N(x_1, x_2, \dots, x_{n-1}, x - y, t) \end{aligned}$$

So $\lambda z_1 + (1 - \lambda)z_2 \in R_A^t(x)$ and $R_A^t(x)$ is convex. ■

Theorem 6: For $t > 0$ and each $x \in X$. let A be a t -coproximinal subspace of a fuzzy anti- n -normed space (X, N) . Then

- (a) If \check{A}_x^t is a t -compact set then A is t -quasi-co-Chebyshev.
- (b) If \check{A}_x^t is a t -closed set then $R_A^t(x)$ is t -closed for every $x \in X$.

Proof: (i) Suppose $x \in X$ and $\{y_n\}$ is a sequence in $R_A^t(x)$. Since $x - y_n \in \check{A}_x^t$ and \check{A}_x^t is a t -compact set, there exists a subsequence $\{x - y_{n_k}\}$ that t -convergence to $x - y_0 \in \check{A}_x^t$. Consequently, $\{y_n\}$ has a subsequence $y_{n_k} \xrightarrow{t} y_0 \in R_A^t(x)$ and hence A is t -quasi-co-Chebyshev.

(ii) The proof is similar to (i). ■

Definition 9: A subset A of a fuzzy anti- n -normed space (X, N) is called to be t -boundedly compact if every t -bounded

sequence in A has a subsequence t -converging to an element of X .

Theorem 7: Suppose for some $t > 0$ and each $x \in X$, A is a t -boundedly compact and t -closed subset of a fuzzy anti- n -normed space (X, N) , then, A is t -quasi-co-Chebyshev.

Proof: Let $\{y_n\}$ be any sequence in $R_A^t(x)$. Then $N(x_1, x_2, \dots, x_{n-1}, y_n - y, t) \leq N(x_1, x_2, \dots, x_{n-1}, x - y, t)$ for every $y \in A$. Since $R_A^t(x)$ is t -bounded, $\{y_n\}$ is a t -bounded sequence in A , and so $\{y_n\}$ has a t -convergent subsequence $\{y_{n_k}\}$, let $y_{n_k} \xrightarrow{t} y_0 \in A$, as A is t -closed. Consider

$$\begin{aligned} & N(x - 1, x_2, \dots, x_{n-1}, y_0 - y, t) \\ &= \lim_k N(x_1, x_2, \dots, x_{n-1}, y_{n_k} - y, t) \\ &\leq N(x_1, x_2, \dots, x_{n-1}, x - y, t) \end{aligned}$$

for every $y \in A$. So $y_0 \in R_A^t(x)$, which implies that A is t -quasi-co-Chebyshev. ■

Definition 10: Let (X, N) be a fuzzy anti- n -normed space and A be a subset of X . For $t > 0$ and an element $x \in X$ is said to be t -orthogonal to an element $y \in X$, and we denote it by $x \perp_x^t y$, if $N(x_1, x_2, \dots, x_{n-1}, x + \lambda y, t) \geq N(x_1, x_2, \dots, x_{n-1}, x, t)$ for all scalar $\lambda \in \mathbb{R}$, $\lambda \neq 0$. We say $A \perp_x^t y$ if $x \perp_x^t y$ for every $x \in A$.

Theorem 8: For $t > 0$ and each $x \in X$ and $y_0 \in A$, let (X, N) be a fuzzy anti- n -normed space and A be a subspace of X . If $A \perp_x^t x - y_0$ then $y_0 \in R_A^t(x)$.

Proof: Suppose $t > 0$, $x \in X$ and $A \perp_x^t x - y_0$. Then $N(x_1, x_2, \dots, x_{n-1}, a + \lambda(x - y_0), t) \geq N(x_1, x_2, \dots, x_{n-1}, a, t)$ for all $a \in A$ and all scalar $\lambda \in \mathbb{R}$, $\lambda \neq 0$. Then $N(x_1, x_2, \dots, x_{n-1}, x - y_0 + \lambda^{-1}a, \frac{t}{|\lambda|}) \geq N(x_1, x_2, \dots, x_{n-1}, \lambda^{-1}a, \frac{t}{|\lambda|})$. Hence $N(x_1, x_2, \dots, x_{n-1}, x - a', \frac{t}{|\lambda|}) \geq N(x_1, x_2, \dots, x_{n-1}, y_0 - a', \frac{t}{|\lambda|})$, where $a' = y_0 - \lambda^{-1}a$. Now if $\lambda = 1$ then, $N(x_1, x_2, \dots, x_{n-1}, y - y_0, t) \geq N(x_1, x_2, \dots, x_{n-1}, x - y, t)$ for all $y \in A$, and so $y_0 \in R_A^t(x)$. ■

IV. F-BEST COAPPROXIMATION

Definition 11: Let A be a nonempty subset of a fuzzy anti- n -normed space (X, N) . An element $y_0 \in A$ is said to be an F -best coapproximation of $x \in X$ from A if it is a t -best coapproximation of x from A , for every $t > 0$, i.e.,

$$y_0 \in \bigcap_{t \in (0, \infty)} R_A^t(x).$$

The set of all elements of F -best coapproximation of X from A is denoted by $FR_A^t(x)$, i.e.,

$$FR_A^t(x) = \bigcap_{t \in (0, \infty)} R_A^t(x).$$

If each $x \in X$ has at least (respectively exactly) one F -best coapproximation in A , then A is called F -coproximinal (respectively F -co-Chebyshev) set.

Example 2: Let $X = \mathbb{R}^3$. Define $N : X \times X \times X \times [0, \infty) \rightarrow [0, 1]$ by

$$\begin{aligned} N(x_1, x_2, x_3, t) &= \frac{\|x_1, x_2, x_3\|}{t}, \quad \text{if } t > 0, t \in \mathbb{R}, x_1, x_2, x_3 \in X \\ &= 1, \quad \text{if } t \leq 0, t \in \mathbb{R}, x_1, x_2, x_3 \in X. \end{aligned}$$

where $\|x_1, x_2, x_3\| = \min_{1 \leq i \leq 3} \sum_{j=1}^3 |x_{ij}|$. Then (X, N) is a fuzzy anti-3-normed linear space. Let $A = \{(a, b, c) \in \mathbb{R}^3 : a^2 + b^2 \leq 1, 0 \leq c \leq a^2 + b^2\}$ and $x_1 = (1, 0, 0)$, $x_2 = (0, 1, 0)$, $x_3 = (0, 0, 4)$ are in X . Let $a_0 = (0, -1, 1)$ and $a_1 = (0, 1, 1)$ are in A . Then $(0, -1, 1), (0, 1, 1) \in FR_A^t(0, 0, 4)$. So A is not a F -co-Chebyshev set.

Theorem 9: Let $\{\|\cdot, \cdot, \dots, \cdot\|_\alpha^* : \alpha \in (0, 1]\}$ be a descending family of α - n -norm on X corresponding to the fuzzy anti- n -norm on X . Then $y_0 \in A$ is a best coapproximation to $x \in X$ in the descending family of α - n -norm on X corresponding to the fuzzy anti- n -norm on X if and only if y_0 is a F -best coapproximation to x in the fuzzy anti- n -normed space (X, N) .

Proof: For each $x \in X$, y_0 is a best coapproximation to $x \in X$ in the descending family of α - n -norm on X corresponding to the fuzzy anti- n -norm on X . if and only if $\|x_1, x_2, \dots, y - y_0\|_\alpha^* \leq \|x_1, x_2, \dots, x - y\|_\alpha^*$, for every $y \in A$, if and only if $\frac{t}{t + \|x_1, x_2, \dots, y - y_0\|_\alpha^*} \geq \frac{t}{t + \|x_1, x_2, \dots, x - y\|_\alpha^*}$ for every $y \in A$ and $t \in (0, \infty)$, if and only if $N(x_1, x_2, \dots, x_{n-1}, y - y_0, t) \leq N(x_1, x_2, \dots, x_{n-1}, x - y, t)$ for every $y \in A$ and $t \in (0, \infty)$, if and only if $y_0 \in FR_A^t(x)$. ■

Definition 12: Let (X, N) be a fuzzy anti- n -normed space and A be a subset of X . For each element $x \in X$ is said to be F -orthogonal to an element $y \in X$ and we denote it by $x \perp_x^F y$, if for every $t > 0$, $x \perp_x^t y$. We say $A \perp_x^F y$ if $x \perp_x^F y$ for every $x \in A$.

Theorem 10: Let $\{\|\cdot, \cdot, \dots, \cdot\|_\alpha^* : \alpha \in (0, 1]\}$ be a descending family of α - n -norm on X corresponding to the fuzzy anti- n -norm on X . Then $x \in X$ is Brikhoff orthogonal to $y \in X$ in the descending family of α - n -norm on X corresponding to the fuzzy anti- n -norm on X if and only if x is a F -orthogonal to y in the fuzzy anti- n -normed space (X, N) .

Proof: For each $x \in X$, x is a Brikhoff orthogonal to $y \in X$ in the descending family of α - n -norm on X corresponding to the fuzzy anti- n -norm on X . if and only if $\|x_1, x_2, \dots, x_{n-1}, x\|_\alpha^* \leq \|x_1, x_2, \dots, x_{n-1}, x + \lambda y\|_\alpha^*$, for every scalar $\lambda \in \mathbb{R} \setminus \{0\}$, if and only if $\frac{t}{t + \|x_1, x_2, \dots, x_{n-1}, x\|_\alpha^*} \geq \frac{t}{t + \|x_1, x_2, \dots, x_{n-1}, x + \lambda y\|_\alpha^*}$ for every scalar $\lambda \in \mathbb{R} \setminus \{0\}$ and $t > 0$, if and only if $N(x_1, x_2, \dots, x_{n-1}, x, t) \leq N(x_1, x_2, \dots, x_{n-1}, x + \lambda y, t)$ for every scalar $\lambda \in \mathbb{R} \setminus \{0\}$ and $t > 0$, if and only if $x \perp_x^F y$. ■

V. CONCLUSION

In this paper we introduced the concept of t -best coapproximation in and F -best coapproximation in fuzzy anti- n -normed spaces and also introduced t -coproximinal and t -co-Chebyshev in fuzzy anti- n -normed spaces. Then prove several theorems pertaining to this sets illustrate with example.

REFERENCES

- [1] T. Bag and T. K. Samanta, A comparative study of fuzzy norms on a linear space, Fuzzy Sets and Systems. 159 (2008) 670 -684.
- [2] C. Franchetti and M. Furi, Some characteristic properties of real Hilbert spaces, Rev. Roumaine Math. Pures Appl. 17 (1972) 1045 - 1048.
- [3] S. Gähler, Lineare 2-normierte Räume, Math. Nachr. 28 (1964) 1-43.
- [4] S. Gähler and Unter Suchyngen Über Verallgemeinerte m -metrische Räume I, Math. Nachr. 40 (1969) 165 - 189.
- [5] H. Gunawan and M. Mashadi, On n -normed spaces, Int. J. Math & Math. Sci. 27(10) (2001) 631 - 639.

- [6] Iqbal H. Jebril and T. K. Samanta, Fuzzy anti-normed space, J. Mathematics and Technology, February (2010) 66 -77.
- [7] A. K. Katsaras, Fuzzy topological vector spaces, Fuzzy Sets and Systems, 12 (1984) 143 - 154.
- [8] J. Kavikumar, Y. B. Jun and Azme Khamis, The Riesz theorem in fuzzy n -normed linear spaces, J. Appl. math & Informatics. 27 (2009) (2-4) 841 - 555.
- [9] R. Malčeski, Strong n -convex n -normed spaces, Mat. Bilten. 21 (47) (1997) 81 - 102.
- [10] A. Misiak, n -inner product spaces, Math. Nachr. 140 (1989) 299 - 319.
- [11] AL. Narayanan and S. Vijayabalaji, Fuzzy n -normed linear space, Int. J. Math. & Math. Sci. 24 (2005) 3963 - 3977.
- [12] Papini, P. L., and Singer, I. Best coapproximation in normed linear spaces, Monatshefte für Mathematik, 88 (1979) 27 - 44.
- [13] G.S. Rao and R. Saravanan, Characterization of best uniform coapproximation, Approximation Theory and its Applications. 15 (1999) 23 - 37.
- [14] B. Surender Reddy, Some results on t -best approximation in fuzzy anti- n -normed linear spaces, Int. J. Math. Archive. 2 (9) (2011) 1747 -1757.
- [15] B. Surender Reddy, Fuzzy anti- n -normed linear space, J. Mathematics and Technology. 2(1) February (2011) 14 - 26.
- [16] S. M. Vaezpour and F. Karimi, t -Best approximation in fuzzy normed spaces. Iranian J. Fuzzy Systems. 5 (2) (2008) 93 - 98.
- [17] S. Vijayabalaji and N. Thillaigovindan, Complete fuzzy n -normed linear space, Journal of Fund. Sci.3 007) 119 - 126.