

# Differential Sandwich Theorems with Generalised Derivative Operator

<sup>1</sup>Maslina Darus and <sup>2</sup>Khalifa Al-Shaqsi

**Abstract**—In this paper, a generalized derivatives operator  $\mathfrak{D}_{\lambda,\beta}^n f$  introduced by the authors will be discussed. Some subordination and superordination results involving this operator for certain normalized analytic functions in the open unit disk will be investigated. Our results extend corresponding previously known results.

**Keywords**—Analytic functions, Univalent functions, Derivative operator, Differential subordination, Differential superordination.

## I. INTRODUCTION

Denote by  $\mathbb{U}$  the unit disk of the complex plane:

$$\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}.$$

Let  $\mathcal{H}(\mathbb{U})$  be the space of analytic function in  $\mathbb{U}$ . Let

$$\mathcal{A}_n = \{f \in \mathcal{H}(\mathbb{U}), f(z) = z + a_{n+1}z^{n+1} + \dots,\}$$

for  $(z \in \mathbb{U})$  with  $\mathcal{A}_1 = \mathcal{A}$ .

For  $a \in \mathbb{C}$  and  $n \in \mathbb{N}$  we let

$$\mathcal{H}[a, n] = \{f \in \mathcal{H}(\mathbb{U}), f(z) = a + a_n z^n + a_{n+1} z^{n+1} + \dots,\} \\ (z \in \mathbb{U}).$$

If functions  $f$  and  $F$  are analytic in  $\mathbb{U}$ , then we say that  $f$  is subordinate to  $F$ , and write  $f \prec F$ , if there exists a Schwarz function  $w$  analytic in  $\mathbb{U}$  with  $|w(z)| < 1$  and  $w(0) = 0$  such that  $f(z) = F(w(z))$  in  $\mathbb{U}$ . Furthermore, if the function  $F(z)$  is univalent in  $\mathbb{U}$ , then  $f(z) \prec F(z) (z \in \mathbb{U}) \Leftrightarrow f(0) = F(0)$  and  $f(\mathbb{U}) \subset F(\mathbb{U})$ .

A function  $f$ , analytic in  $\mathbb{U}$ , is said to be convex if it is univalent and  $f(\mathbb{U})$  is convex.

Let  $p, h \in \mathcal{H}(\mathbb{U})$  and let  $\psi(r, s, t; z) : \mathbb{C}^3 \times \mathbb{U} \rightarrow \mathbb{C}$ . If  $p$  and  $\psi(p(z), zp'(z), z^2p''(z); z)$  are univalent and if  $p$  satisfies the (second-order) differential superordination

$$h(z) \prec \psi(p(z), zp'(z), z^2p''(z); z), \quad (z \in \mathbb{U}) \quad (1)$$

then  $p$  is called a solution of the differential superordination (1). (If  $f$  subordinate to  $F$ , then  $F$  is superordinate to  $f$ ).

An analytic function  $q$  is called a subordinator of the differential superordination, or more simply a subordinator if  $q \prec p$  for all  $p$  satisfying (1). A univalent subordinator  $\tilde{q}$  that satisfies  $q \prec \tilde{q}$  for all subordinants  $q$  of (1) is said to be the best subordinator. (Note that the best subordinator is unique up to a rotation of

$\mathbb{U}$ ). Recently Miller and Mocanu [5] obtained conditions on  $h, q$  and  $\psi$  for which the following implication holds:

$$h(z) \prec \psi(p(z), zp'(z), z^2p''(z); z) \Rightarrow q(z) \prec p(z) \quad (z \in \mathbb{U}).$$

We now state the following definition.

**Definition 1.1:** [3] Let function  $f$  in  $\mathcal{A}$ , then for  $n, \lambda \in \mathbb{N}_0$  and  $\beta > 0$ , we define the following differential operator

$$\mathfrak{D}_{\lambda,\beta}^n f(z) = z + \sum_{k=2}^{\infty} [1 + \beta(k-1)]^n C(k, \lambda) a_k z^k, \quad (z \in \mathbb{U}),$$

where  $C(\lambda, k) = \binom{k+\lambda-1}{\lambda}$ .

Special cases of this operator includes the Ruscheweyh derivative operator  $\mathfrak{D}_{\lambda,1}^0 f(z) \equiv D_\lambda$  [6], the Salagean derivative operator  $\mathfrak{D}_{0,1}^n f(z) \equiv D^n$  [2], the generalized Salagean derivative operator  $\mathfrak{D}_{0,\beta}^n f(z) \equiv D_\beta^n$  [1] and the generalized Ruscheweyh derivative operator  $\mathfrak{D}_{\lambda,\beta}^1 f(z) \equiv D_{\lambda,\beta}$  [4].

For  $n, \lambda \in \mathbb{N}_0$  and  $\beta > 0$ , we obtain the following inclusion relations:

$$\mathfrak{D}_{\lambda,\beta}^{n+1} f(z) = (1 - \beta)\mathfrak{D}_{\lambda,\beta}^n f(z) + \beta z(\mathfrak{D}_{\lambda,\beta}^n f(z))', \quad (2)$$

$$z(\mathfrak{D}_{\lambda,\beta}^n f(z))' = (1 + \lambda)\mathfrak{D}_{\lambda+1,\beta}^n f(z) - \lambda\mathfrak{D}_{\lambda,\beta}^n f(z) \quad (3)$$

The main object of the present paper is to find sufficient condition for certain normalized analytic functions  $f$  to satisfy

$$q_1(z) \prec \frac{D_{\lambda,\beta}^{n+1} f(z)}{D_{\lambda,\beta}^n f(z)} \prec q_2(z),$$

where  $n, \lambda \in \mathbb{N}_0, \beta > 0$  and  $q_1, q_2$  are given univalent functions in  $\mathbb{U}$ . Also, we obtain the number of known results as their special cases.

In order to prove the original results we use shall need the following definition and lemmas. In this paper unless otherwise mentioned  $\alpha, \delta$  are complex numbers.

**Definition 1.2:** [5] Denote by  $Q_\alpha$  the set of all functions  $f$  that are analytic and injective on  $\mathbb{U} - E(f)$ , where

$$E(f) = \left\{ \zeta \in \partial\mathbb{U} : \lim_{z \rightarrow \zeta} f(z) = \infty \right\}$$

and are such that  $f'(\zeta) \neq 0$  for  $\zeta \in \partial\mathbb{U} - E(f)$ .

<sup>1,2</sup>School of Mathematical Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi 43600 Selangor D. Ehsan, Malaysia, email: <sup>1</sup>maslina@ukm.my email: <sup>2</sup>ommath@hotmail.com

**Lemma 1.3:** [7] Let  $q$  be univalent in the unit disk  $\mathbb{U}$  and  $\theta$  and  $\phi$  be analytic in a domain  $\mathcal{D}$  containing  $q(\mathbb{U})$  with  $\phi(w) \neq 0$  when  $w \in q(\mathbb{U})$ . Set

$$\psi(z) = zq'(z)\phi(q(z)) \quad \text{and} \quad h(z) = \theta(q(z)) + \psi(z).$$

Suppose that

- 1)  $\psi(z)$  is starlike univalent in  $\mathbb{U}$ , and
- 2)  $\operatorname{Re} \left\{ \frac{zh'(z)}{\psi(z)} \right\} > 0$  for  $z \in \mathbb{U}$ .

If  $p$  is analytic with  $p(0) = q(0)$ ,  $p(\mathbb{U}) \subseteq \mathcal{D}$ , and

$$\theta(p(z)) + zp'(z)\phi(p(z)) \prec \theta(q(z)) + zq'(z)\phi(q(z)), \quad (4)$$

then

$$p(z) \prec q(z)$$

and  $q$  is the best dominant.

**Lemma 1.4:** [8] Let  $q$  be convex univalent in the unit disk  $\mathbb{U}$  and  $\vartheta$  and  $\varphi$  be analytic in a domain  $\mathcal{D}$  containing  $q(\mathbb{U})$ .

Suppose that

- 1)  $\operatorname{Re} \left\{ \frac{\vartheta'(q(z))}{\varphi(q(z))} \right\} > 0$  for  $z \in \mathbb{U}$  and
- 2)  $\psi(z) = zq'(z)\varphi(q(z))$  is starlike univalent in  $\mathbb{U}$ .

If  $p(z) \in \mathcal{H}[q(0), 1] \cap \mathcal{Q}$  with  $p(\mathbb{U}) \subseteq \mathcal{D}$ , and

$\vartheta(p(z)) + zp'(z)\varphi(p(z))$  is univalent in  $\mathbb{U}$  and

$$\vartheta(q(z)) + zq'(z)\varphi(q(z)) \prec \vartheta(p(z)) + zp'(z)\varphi(p(z)), \quad (5)$$

then  $q(z) \prec p(z)$  and  $q$  is best subordinant.

## II. SUBORDINATION RESULTS

Using Lemma 1.3, we first prove the following theorem.

**Theorem 2.1:** Let  $n, \lambda \in \mathbb{N}_0$ ,  $\beta > 0$  and  $q(z)$  be convex univalent in  $\mathbb{U}$  with  $q(0) = 1$ . Further, assume that

$$\operatorname{Re} \left\{ \frac{2\delta q(z)}{\alpha} + 1 + \frac{zq''(z)}{q'(z)} \right\} > 0 \quad (z \in \mathbb{U}). \quad (6)$$

Let

$$\begin{aligned} \Psi(n, \lambda, \beta, \delta, \alpha; z) = & \frac{\delta[2 - \beta(2 + \lambda)]}{\beta} \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \\ & + \delta\beta(\lambda + 2)(\lambda + 1) \frac{\mathfrak{D}_{\lambda+2, \beta}^n f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \\ & - \delta\beta(\lambda + 1)^2 \frac{\mathfrak{D}_{\lambda+1, \beta}^n f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \\ & + [\alpha + \delta(1 - \frac{1}{\beta})] \left( \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \right)^2 \\ & - \frac{\delta(1 - \beta)[1 - \beta(\lambda + 1)]}{\beta} \end{aligned} \quad (7)$$

If  $f \in \mathcal{A}$  satisfies

$$\Psi(\lambda, \delta, \alpha; z) \prec \delta zq'(z) + (\delta + \alpha)(q(z))^2 \quad (8)$$

then

$$\frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \prec q(z)$$

and  $q$  is the best dominant.

**Proof.** Define the function  $p(z)$  by

$$p(z) = \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \quad (z \in \mathbb{U}). \quad (9)$$

Then the function  $p(z)$  is analytic in  $\mathbb{U}$  and  $p(0) = 1$ .

Therefore, by making use of (2), (3) and (4), we obtain

$$\begin{aligned} & \frac{\delta[2 - \beta(2 + \lambda)]}{\beta} \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} + \delta\beta(\lambda + 2)(\lambda + 1) \frac{\mathfrak{D}_{\lambda+2, \beta}^n f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \\ & - \delta\beta(\lambda + 1)^2 \frac{\mathfrak{D}_{\lambda+1, \beta}^n f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} + [\alpha + \delta(1 - \frac{1}{\beta})] \left( \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \right)^2 \\ & - \frac{\delta(1 - \beta)[1 - \beta(\lambda + 1)]}{\beta} \\ & = \delta zp'(z) + (\delta + \alpha)(p(z))^2 \end{aligned} \quad (10)$$

By using (10) in (8), we have

$$\delta zp'(z) + (\delta + \alpha)(p(z))^2 \prec \delta zq'(z) + (\delta + \alpha)(q(z))^2.$$

By setting  $\theta(w) = \delta w^2$  and  $\phi(w) = \alpha$ , it can be easily observed that  $\theta(w)$  and  $\phi(w)$  are analytic in  $\mathbb{C} - \{0\}$  and that  $\phi(w) \neq 0$ . Hence the result now follows by an application if Lemma 1.3.

**Corollary 2.2:** Let  $q(z) = \frac{1+Az}{1+Bz}$  ( $-1 \leq B < A \leq 1$ ) in Theorem 2.1, further assuming that (6) holds. If  $f \in \mathcal{A}$  then,

$$\Psi(n, \lambda, \beta, \delta, \alpha; z) \prec \delta \frac{(A - B)z}{(1 + Bz)^2} + (\delta + \alpha) \left( \frac{1 + Az}{1 + Bz} \right)^2,$$

$$\Rightarrow \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \prec \frac{1+Az}{1+Bz}, \text{ and } \frac{1+Az}{1+Bz} \text{ is the best dominant.}$$

Also, let  $q(z) = \frac{1+z}{1-z}$ , then for  $f \in \mathcal{A}$  we have,

$$\Psi(n, \lambda, \beta, \delta, \alpha; z) \prec \frac{2\delta z}{(1 - z)^2} + (\delta + \alpha) \left( \frac{1 + z}{1 - z} \right)^2,$$

$$\Rightarrow \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \prec \frac{1+z}{1-z}, \text{ and } \frac{1+z}{1-z} \text{ is the best dominant.}$$

By taking  $q(z) = \left( \frac{1+z}{1-z} \right)^\mu$ , ( $0 < \mu \leq 1$ ), for  $f \in \mathcal{A}$ , we have

$$\Psi(n, \lambda, \beta, \delta, \alpha; z) \prec \frac{2\delta\mu z}{(1 - z)^2} \left( \frac{1 + z}{1 - z} \right)^{\mu-1} + (\delta + \alpha) \left( \frac{1 + z}{1 - z} \right)^{2\mu},$$

$$\Rightarrow \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \prec \left( \frac{1+z}{1-z} \right)^\mu, \text{ and } \left( \frac{1+z}{1-z} \right)^\mu \text{ is the best dominant.}$$

## III. SUPERORDINATION AND SANDWICH RESULTS

Now, by applying Lemma 1.4, we prove the following theorem.

**Theorem 3.1:** Let  $q$  be convex univalent in  $\mathbb{U}$  with  $q(0) = 1$ .

Assume that

$$\operatorname{Re} \left\{ \frac{2(\delta + \alpha)q(z)q'(z)}{\delta} \right\} > 0. \quad (11)$$

Let  $f \in \mathcal{A}$ ,  $\frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \in \mathcal{H}[q(0), 1] \cap \mathcal{Q}$ .

Further, let  $\Psi(n, \lambda, \beta, \delta, \alpha; z)$  given by (7) be univalent in  $\mathbb{U}$  and

$$(\delta + \alpha)(q(z))^2 + \delta z q'(z) \prec \Psi(n, \lambda, \beta, \delta, \alpha; z)$$

then

$$q(z) \prec \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)},$$

and  $q$  is the best subordinant.

**Proof.** Theorem 3.1 follows by using the same technique to prove Theorem 2.1 and by an application of Lemma 1.4.

By using Theorem 3.1, we have the following corollary.

*Corollary 3.2:* Let  $q(z) = \frac{1+A_1z}{1+B_1z}$  ( $-1 \leq B_1 < A_1 \leq 1$ ),  $f \in \mathcal{A}$  and  $\frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \in \mathcal{H}[q(0), 1] \cap Q$ . Further assuming that (11) satisfies. If

$$(\delta + \alpha) \left( \frac{1 + Az}{1 + Bz} \right)^2 + \frac{\delta(A - B)z}{(1 + Bz)^2} \prec \Psi(n, \lambda, \beta, \delta, \alpha; z)$$

then

$$\frac{1 + Az}{1 + Bz} \prec \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)}$$

and  $\frac{1+Az}{1+Bz}$ , is the best subordinant.

Also, by let  $q(z) = \frac{1+z}{1-z}$ ,  $f \in \mathcal{A}$  and  $\frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \in \mathcal{H}[q(0), 1] \cap Q$ . Further assuming that (11) satisfies. If

$$\frac{2\delta z}{(1-z)^2} + (\delta + \alpha) \left( \frac{1+z}{1-z} \right)^2 \prec \Psi(n, \lambda, \beta, \delta, \alpha; z),$$

$\Rightarrow \frac{1+z}{1-z}$  and  $\frac{1+z}{1-z} \prec \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)}$ , is the best subordinant.

Finally, by taking  $q(z) = \left( \frac{1+z}{1-z} \right)^\mu$ , ( $0 < \mu \leq 1$ ),  $f \in \mathcal{A}$  and  $\frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \in \mathcal{H}[q(0), 1] \cap Q$ . Further assuming that (11) satisfies. If

$$\frac{2\delta\mu z}{(1-z)^2} \left( \frac{1+z}{1-z} \right)^{\mu-1} + (\alpha + \delta) \left( \frac{1+z}{1-z} \right)^{2\mu} \prec \Psi(n, \lambda, \beta, \delta, \alpha; z),$$

$\Rightarrow \left( \frac{1+z}{1-z} \right)^\mu$ , and  $\left( \frac{1+z}{1-z} \right)^\mu \prec \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)}$  is the best subordinant.

Combining the results of differential subordination and superordination, we state the following Sandwich Theorems .

*Theorem 3.3:* Let  $q_1$  and  $q_2$  be convex univalent in  $\mathbb{U}$  and satisfies (11) and (6), respectively. If  $f \in \mathcal{A}$ ,  $\frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \in \mathcal{H}[q(0), 1] \cap Q$  and  $\Psi(n, \lambda, \beta, \delta, \alpha; z)$  is univalent in  $\mathbb{U}$ , and

$$\delta z q_1'(z) + (\delta + \alpha)(q_1(z))^2 \prec \Psi(n, \lambda, \beta, \delta, \alpha; z) \delta z q_2'(z) + (\delta + \alpha)(q_2(z))^2, \tag{12}$$

then

$$q_1(z) \prec \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \prec q_2(z),$$

and  $q_1$  and  $q_2$  are respectively the best subordinant and best dominant.

For  $q_1(z) = \frac{1+A_1z}{1+B_1z}$ ,  $q_2(z) = \frac{1+A_2z}{1+B_2z}$ ,

where  $(-1 \leq B_2 \leq B_1 < A_1 \leq A_2 \leq 1)$ ,

we have the following corollary.

*Corollary 3.4:* If  $f \in \mathcal{A}$ ,  $\frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \in \mathcal{H}[q(0), 1] \cap Q$  and

$$\begin{aligned} &\Psi_1(A_1, B_1, n, \lambda, \beta, \delta, \alpha; z) \\ &\prec \Psi(n, \lambda, \beta, \delta, \alpha; z) \\ &\prec \Psi_2(A_2, B_2, n, \lambda, \beta, \delta, \alpha; z) \end{aligned}$$

then

$$\frac{1 + A_1z}{1 + B_1z} \prec \frac{\mathfrak{D}_{\lambda, \beta}^{n+1} f(z)}{\mathfrak{D}_{\lambda, \beta}^n f(z)} \prec \frac{1 + A_2z}{1 + B_2z}$$

where

$$\begin{aligned} \Psi_1(A_1, B_1, n, \lambda, \beta, \delta, \alpha; z) &:= (\delta + \alpha) \left( \frac{1 + A_1z}{1 + B_1z} \right)^2 \\ &+ \frac{\delta(A_1 - B_1)z}{(1 + B_1z)^2}, \end{aligned}$$

$$\begin{aligned} \Psi_2(A_2, B_2, n, \lambda, \beta, \delta, \alpha; z) &:= (\delta + \alpha) \left( \frac{1 + A_2z}{1 + B_2z} \right)^2 \\ &+ \frac{\delta(A_2 - B_2)z}{(1 + B_2z)^2}. \end{aligned}$$

Hence  $\frac{1+A_1z}{1+B_1z}$  and  $\frac{1+A_2z}{1+B_2z}$  are respectively the best subordinant and best dominant.

*Remark 3.5:* For special cases of the above results follows by choosing different values of  $n, \lambda$  and  $\beta$ .

ACKNOWLEDGMENT

The work presented here was fully supported by eScience-fund: 04-01-02-SF0425, MOSTI, Malaysia.

REFERENCES

- [1] F. M. Al-Oboudi, On univalent functions defined by a generalized Sălăgean operator, *Ind. J. Math. Math. Sci.* **27**, (2004), 1429-1436.
- [2] G. Ş. Sălăgean, Subclasses of univalent functions, *Lecture Note in Math.*(Springer-Verlag), **1013**, (1983), 362-372.
- [3] K. Al-Shaqsi and M. Darus, Differential subordination with generalized derivative operator. (Submitted)
- [4] K. Al Shaqsi and M. Darus, On univalent functions with respect to  $k$ -symmetric points defined by a generalized Ruscheweyh derivatives operator.(Submitted)
- [5] S. S. Miller and P. T. Mocanu, Subordinants of differential superordinations, *Complex Variables Theory Appl.* **48**(10), (2003), 815-826.
- [6] St. Ruscheweyh, New criteria for univalent functions, *Proc. Amer. Math. Soc.* **49**, (1975), 109-115.
- [7] S. S. Miller and P. T. Mocanu, Differential Subordinations: Theory and Applications, *Marcel Dekker Inc.*, New York, 2000.
- [8] T. Bulboacă, Classes of first order differential superordinations, *Demonstratio Math.* **35**(2), (2002), 287-292.

**Maslina Darus** received her Bachelor degree in Mathematics from Acadia University, Canada, in 1992 and PhD in Pure Mathematics (Complex Analysis) from University of Wales, Swansea, U. Kingdom in 1996. Currently, she is a Professor of Mathematics in School of Mathematical Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia. She is also holding a post as Head of Mathematics program. Her special interest is in the geometric function theory and its applications.

**Khalifa Al-Shaqsi** is currently a PhD student under supervision of Professor Maslina Darus. He obtained Master degree in Mathematics at School of Mathematical Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia in 2006.