

# Emergence of Fluoroquinolone Resistance in Pigs, Nigeria

Igbakura I. Luga, Alex A. Adikwu

**Abstract**—A comparison of resistance to quinolones was carried out on isolates of Shiga toxin-producing *Escherichia coli* O157:H7 from cattle and *mecA* and *nuc* genes harbouring *Staphylococcus aureus* from pigs. The isolates were separately tested in the first and current decades of the 21<sup>st</sup> century. The objective was to demonstrate the dissemination of resistance to this frontline class of antibiotic by bacteria from food animals and bring to the limelight the spread of antibiotic resistance in Nigeria. A total of 10 isolates of the *E. coli* O157:H7 and 9 of *mecA* and *nuc* genes harbouring *S. aureus* were obtained following isolation, biochemical testing, and serological identification using the Remel Wellcolex *E. coli* O157:H7 test. Shiga toxin-production screening in the *E. coli* O157:H7 using the verotoxin *E. coli* reverse passive latex agglutination (VTEC-RPLA) test; and molecular identification of the *mecA* and *nuc* genes in *S. aureus*. Detection of the *mecA* and *nuc* genes were carried out using the protocol by the Danish Technical University (DTU) using the following primers *mecA*-1: 5'-GGGATCATAGCGTCATTATTC-3', *mecA*-2: 5'-AACGATTGTGACACGATAGCC-3', *nuc*-1: 5'-TCAGCAAATGCATCACAAACAG-3', *nuc*-2: 5'-CGTAAATGCACTTGCTTCAGG-3' for the *mecA* and *nuc* genes, respectively. The *nuc* genes confirm the *S. aureus* isolates and the *mecA* genes as being methicillin-resistant and so pathogenic to man. The fluoroquinolones used in the antibiotic resistance testing were norfloxacin (10 µg) and ciprofloxacin (5 µg) in the *E. coli* O157:H7 isolates and ciprofloxacin (5 µg) in the *S. aureus* isolates. Susceptibility was tested using the disk diffusion method on Muller-Hinton agar. Fluoroquinolone resistance was not detected from isolates of *E. coli* O157:H7 from cattle. However, 44% (4/9) of the *S. aureus* were resistant to ciprofloxacin. Resistance of up to 44% in isolates of *mecA* and *nuc* genes harbouring *S. aureus* is a compelling evidence for the rapid spread of antibiotic resistance from bacteria in food animals from Nigeria. Ciprofloxacin is the drug of choice for the treatment of Typhoid fever, therefore widespread resistance to it in pathogenic bacteria is of great public health significance. The study concludes that antibiotic resistance in bacteria from food animals is on the increase in Nigeria. The National Food and Drug Administration and Control (NAFDAC) agency in Nigeria should implement the World Health Organization (WHO) global action plan on antimicrobial resistance. A good starting point can be coordinating the WHO, Office of International Epizootics (OIE), Food and Agricultural Organization (FAO) tripartite draft antimicrobial resistance monitoring and evaluation (M&E) framework in Nigeria.

**Keywords**—Fluoroquinolone, Nigeria, resistance, *Staphylococcus aureus*.

## I. INTRODUCTION

THE WHO has described fluoroquinolones as critically important medicines for the treatment of human infections and advised prudent use in both human and veterinary medicine [1]. Quinolones including their fluorinated

Igbakura Innocent Luga is with the Federal University of Agriculture, Nigeria (e-mail: lugaigbakura@gmail.com).

derivatives are widely used as the drug of choice in the treatment of urinary tract infection and typhoid fever. This extensive use along with non-therapeutic use in animal production has in part been responsible for the spread of resistance to fluoroquinolones. Several reports have described fluoroquinolone resistance in food animals [2]-[4] as well as commensal bacteria populations from humans, animals and environment [5]. On the other hand, the role of plasmid-mediated quinolone resistance (PMQR) gene, other resistant genes and mechanisms have produced widespread transfer of quinolone resistance between unrelated bacteria [5].

Pork is the most preferred source of animal protein in Makurdi, Nigeria, and virtually all residents consume it. However, pork has been reported in many parts of the world to be a vehicle for spread of pathogens that cause food borne illnesses due to its contamination during the slaughtering process [6]. For verification of slaughter hygiene conditions in the daily practice, the microbial status of carcasses is often determined by monitoring indicator organisms on carcasses at the end of slaughter [7]-[9], some of which include *Escherichia coli*, *Salmonella* spp., *Listeria monocytogenes*, *Staphylococcus aureus*, and *Campylobacter jejuni*. Methicillin-resistant *Staphylococcus aureus* (MRSA) is an increasing cause of health care-associated and livestock-associated infections worldwide [10]. The confirmation of the presence of the *mecA* gene has been the “golden standard” for detection of methicillin resistant *S. aureus* (MRSA) worldwide [11].

Antimicrobial resistance is a threat to humanity's way of life. Individual countries must recognize this threat and join in the global effort at developing strategies for antimicrobial resistance. This study has the objective of highlighting an important aspect of antibiotic resistance which is the resistance to a frontline antimicrobial agent, as well as increasing awareness of the menace in Nigeria.

## II. MATERIALS AND METHODS

### A. Ethical Considerations

Ethical approval for the studies was granted by the Research Ethics Committee of the Faculty of Veterinary Medicine, Ahmadu Bello University Zaria, Certificate no. 99/051 and 015/101.

### B. Sample Collection

Between January, 2000 and November, 2006 *E. coli* O157:H7 isolates were obtained from cattle on farms in North-western Nigeria (n=1,800). This was done using a cross-sectional design. Later, from January to November, 2015,

swab samples were obtained from pigs at slaughter in Makurdi, North-central, Nigeria (n=241).

#### C. Isolation of *E. coli* O157:H7 and *Staphylococcus aureus*

The samples from cattle were cultured [12]. Colourless colonies on CR-SMAC agar were selected and biochemically identified using standard procedures. Presumptive identification of *E. coli* O157:H7 was carried out using the Remel Wellcolex *E. coli* kit. (Remel Europe Ltd., UK). The performance of the test and control latex reagents was confirmed using fresh overnight cultures of a control strain EHEC EDL 933. The production of Shiga toxins 1 and 2 (Stx 1 and Stx 2) by isolates that were serologically detected using the Vero Toxin *E. coli* Reverse Passive Latex Agglutination (VTEC-RPLA) test kit (Oxoid Ltd., England). While the sample from pigs were enriched and plated on baird parker agar supplemented with egg yolk tellurite (Oxoid Ltd., England). Typical colonies were subcultured and tested for beta-haemolysis based on [13]. Beta-haemolytic isolates were subjected to DNase [14] and microgen Staph- ID identification [15].

#### D. Identification of *mecA* and *nuc* Genes in *S. aureus*

2.4.1 DNA extraction was carried out using the ZR Fungal/Bacterial DNA mini prepk kit (ZymoBIOMICS™ Epigenetics®, USA), based on the manufacturer's recommendation.

##### 1) Detection of *mecA* and *nuc* Genes

A monoplex PCR assay for detection of *mecA* and *nuc* genes (species specific) were carried out using the protocol described by [16]. Briefly, amplification of the target genes was performed in a total reaction volume of 25 µl containing 2.5 µl of 10 X PCR buffer, 1 µl MgCl<sub>2</sub> (25 mM), 0.25 µl of each dNTP (25 mM each), and 0.25 µl of 2U Taq DNA polymerase, 0.25 µl of each primers (130 µg/ml), 15.75 µl of PCR water and 5 µl template DNA. The PCR amplification was carried out with an initial denaturation of 94°C for 5 min, followed by 30 cycles of denaturation at 94°C for 30 sec, annealing at 55°C for 30 sec and extension at 72°C for 1 min and a final extension at 72°C for 10 min. Of the amplified products, 10 µl were electrophoresed in 1% agarose, stained with ethidium bromide using a 100 bp ladder (ThermoScientific®) and visualized with a UV transilluminator.

#### E. Fluoroquinolone Resistance Testing of *E. coli* O157:H7 and Methicillin-Resistant *S. aureus*

*In vitro* susceptibility tests were performed by the standardized single disc diffusion method [17] in Muller-Hinton Agar (MHA) (Oxoid Ltd., England) with antimicrobial disc using a recommended *E. coli* O157:H7 reference strain (EDL 933) as the control organism. The antimicrobial agents tested and their disc concentrations in micrograms were as follows: ciprofloxacin (CIP5) and norfloxacin (NOR10) with the *E. coli* O157:H7. The isolates of methicillin-resistant *S. aureus* (MRSA) were tested using ciprofloxacin (CIP5).

From a pure culture of the organisms on SMAC agar, five

colonies of the organisms to be tested were transferred with a wire loop to a test tube containing 4 ml of tryptone soya broth. All the five colonies were colourless, similar in morphology and were picked successively. The tubes were incubated at 37°C for 2 hours. The suspension was standardized by diluting with 0.85% saline to a density visually equivalent to the McFarland standard 0.5 (a turbidity standard prepared by add 0.5ml of 1% barium chloride solution to 99.5 of 1% sulphuric acid). This equates to approximately 10<sup>8</sup> enterobacteriaceae organisms per ml. MHA plates were prepared and dried in an incubator, cotton swabs were immersed into the culture and the entire surface of MHA was spread evenly with the suspension. The inoculum was allowed to dry with the plates covered for 10 minutes. Using aseptic techniques, each disc was placed and pressed gently to ensure full contact with the agar surface. The inverted plates were incubated immediately at 37°C for 24 hours.

The diameter of the zone of inhibition of growth including the diameter of the disc was measured to the nearest millimetre. The zone produced by each isolate was compared with that produced by the control strain (EDL 933). A strain was considered sensitive if the diameter of its zone of inhibition of growth by the antimicrobial agent was greater or equal to 4 mm less than that of the control culture. The average diameter of the inhibition zone produced by the control strain was 16 mm, so a zone was moderately sensitive if its zone of inhibition was 11-12 mm and resistant if it showed no zone if inhibition of growth or if the diameter of its zone was less or equal to 10 mm [18].

### III. RESULTS

#### A. Identification of *nuc* and *mecA* Genes in *S. aureus*

A total of 40 isolates from all the locations combined out of the 54 isolates positive for *Staphylococcus aureus* based on biochemical test were selected for polymerase chain reaction (PCR). A total of 30 (75%) isolates were positive for *Staphylococcus aureus* showing amplification of *nuc* gene at 255 bp (Fig. 1) and nine (22.5%) isolates were positive for the *mecA* gene (Fig. 2).




Fig. 1 Gel picture showing PCR for detection of *nuc* gene to confirm *Staphylococcus aureus* species from isolates. Lane M: 100 bp molecular ladder (ThermoScientific). Lanes 1-40 are presumptive *Staphylococcus aureus* isolates. Thirty lanes showed amplification of the *nuc* gene at 255 bp (DTU) specific for *Staphylococcus aureus*




Fig. 2 Gel picture showing PCR for detection of *mecA* gene to confirm methicillin-resistant *Staphylococcus aureus* strains from isolates. Lane M: 100 bp molecular ladder (ThermoScientific). Lanes 1-40 are representatives of isolates. Nine lanes showed amplification of the *mecA* gene at 527bp (DTU) specific for methicillin-resistant *Staphylococcus aureus*

#### B. Detection of Shiga Toxin-Producing *E. coli* O157:H7 and MRSA and Fluoroquinolone Resistance of the Isolates

Up to 0.6% (10/1800) isolates of *E. coli* O157:H7 were confirmed as Shiga toxin-producing based on the result of the VTEC- RPLA test (Table I) and 3.7% (9/241) of MRSA were obtained based on detection of *mecA* gene (Fig. 2, Table I). Fluoroquinolone resistance was not detected in the 10 isolates of *E. coli* O157:H7, as all the phenotypes tested were susceptible to both norfloxacin and ciprofloxacin. However, 44% (4/9) of the MRSA tested were resistant to ciprofloxacin (Table I).

TABLE I  
EMERGENCE OF FLUOROQUINOLONE RESISTANCE IN PIGS, NIGERIA

Time frame	No. of isolates tested		No. positive	% positive
	<i>E. coli</i> O157:H7	MRSA		
1 <sup>st</sup> decade (2000-2010)	10	-	0	0
2 <sup>nd</sup> decade (2011-date)	-	9	4	44

MRSA – Methicillin-resistant *Staphylococcus aureus*

#### IV. DISCUSSION

The study identified the two categories of *E. coli* Shiga toxins. The Shiga toxin 1 (Stx1) was in 10% (1/10) and Shiga toxin 2 (Stx2) in 90% (9/10) of the isolates confirming pathogenicity and membership of STEC strains [19]. A similar lop-sided ratio of production of Stx1: Stx2 has been reported [20]. Resistance to fluoroquinolones, norfloxacin and ciprofloxacin was not detected in a population of *E. coli* O157:H7 in the dawn of the 21<sup>st</sup> century in the cattle study. This could have been the case since fluoroquinolone preparations were not yet available for non-therapeutic use in animals in the country. The availability and indiscriminate use of the antibiotic explains the emergence of phenotypes of *S. aureus* that are resistance to fluoroquinolones in pigs. Extensive use and emergence of fluoroquinolone resistance in food animals has been reported by [5].

Methicillin resistance is the mechanism of *S. aureus* to overcome anti-staphylococcal agents. Methicillin resistant *S. aureus* (MRSA) is an increasing cause of health and livestock

associated infection worldwide [10]. The altered penicillin-binding protein (PBP2a) is associated with methicillin resistance and this protein has reduced affinity for beta-lactam antibiotics which is encoded by the *mecA* has been the 'golden standard' for detection of methicillin resistant *S. aureus* (MRSA) worldwide [11]. This study provided evidence of MRSA by the detection of *mecA* gene in 3.7% (9/241) of the isolates tested; similar findings have also demonstrated and reported in studies performed by [21]-[24]. The *nuc* gene encodes the thermonuclease enzyme of *S. aureus* its amplification is used for speciation of *S. aureus* [25], [26]. This gene was amplified in all the isolates that harboured the *mecA* gene in this study.

The study has demonstrated that resistance to fluoroquinolones is very high in isolates of MRSA from pigs in Nigeria. Similar high or extremely high resistance to fluoroquinolones in food animals have been reported only in *E. coli* populations [27], making this finding of public health significance.

#### V. CONCLUSION AND RECOMMENDATIONS

##### A. Conclusion

The study concludes that extensive and indiscriminate use of fluoroquinolones in humans and in food animals has led to the emergence of fluoroquinolones resistance in Nigeria.

##### B. Recommendation

It is important that Nigeria should be a part of the WHO, OIE, FAO initiative for addressing antimicrobial resistance. A first step could be implementation of M&E of antimicrobial resistance protocols that are designed specifically for the country.

#### REFERENCES

- [1] World Health Organization. "WHO Advisory Group on Integrated Surveillance of Antimicrobial Resistance (AGISAR). Critically important antimicrobials for human medicine 3rd Revision 2011. WHO Document Production Services, Geneva, Switzerland, "Clinical Infectious Disease, vol. 55, pp. 712–719, Oct. 2012
- [2] R. J. Gosling, C. S. Clouting, L. P. Randall, R. A. Horton and R. H. Davies, "Ciprofloxacin resistance in *E. coli* isolated from turkeys in Great Britain," *Avian Pathology*, vol. 41, pp. 83–89, Feb. 2012.
- [3] I. Literak, M. Micudova, D. Tausova, A. Cizek, M. Dolejska, I. Papousek, J. Prochazka, J. Vojtech, F. Borleis, L. Guardone, S. Guenther, J. Hordowski, C. Lejas, W. Meissner, B. F. Marcos, and M. Tucakov, "Plasmid-mediated quinolone resistance genes in fecal bacteria from rooks commonly wintering throughout Europe," *Microbial Drug Resistance*, vol. 18, pp. 567–573, Nov. 2012.
- [4] A. Agabou, N. Lezzar, Z. Ouchenane, S. Khemissi, D. Satta, A. Sotto, J. P. Lavigne and A. Pantel, "Clonal relationship between human and avian ciprofloxacin-resistant *Escherichia coli* isolates in North-Eastern Algeria," *European Journal of Clinical Microbiology and Infectious Diseases*, vol. 35, pp. 227–234, Feb. 2016.
- [5] M. Röderova, D. Halova, I. Papousek, M. Dolejska, M. Masarikova, V. Hanulik, V. Pudova, P. Broz, M. Htoutou-Sedlakova, P. Sauer, J. Bardon, A. Cizek, M. Kolar and I. Literak, "Characteristics of Quinolone Resistance in *Escherichia coli* Isolates from Humans, Animals, and the Environment in the Czech Republic," *Frontiers in Microbiology*, vol. 7, pp. 2147, Jan. 2017.
- [6] A. S. Fahrion, L. Jamir, K. Richa, B. Sonuwar, V. Rutsa, S. Ao, V. P. Padmakumar, R. P. Deka and D. Grace, "Food Safety Hazards in the Pork chain in Nagaland, North East India: Implication for human health," *International Journal of Environmental Research and Public*

- Health*, vol. 11, pp. 403-417, Jan. 2014.
- [7] M. H. Brown, C. O. Gill, J. Hollingsworth, I. R. Nickelson, S. Seward, J. J. Sheridan, T. Stevenson, J. L. Sumner, D. M. Theno, W. R. Osborne, and D. Zink, "The role of microbiological testing in systems for assuring the safety of beef," *International Journal of Food Microbiology*, vol. 62, pp.7-1, Dec. 2000.
- [8] J. R. Ruby, J. Zhu, and S. C. Ingham, "Using indicator bacteria and Salmonella spp. test results from three large-scale beef abattoirs over an 18-month period to evaluate intervention system efficacy and plan carcass testing for Salmonella spp.," *Journal of Food Protection*, vol. 70, pp. 2732-2740, Sep. 2007.
- [9] C. Zweifel, D. Baltzerand R. Stephan, "Microbiological contamination of cattle and pig carcasses at five abattoirs determined by swab sampling in accordance with EU decision 2001/471/EC," *Meat Science*, vol. 69, pp. 559-566, Mar. 2005.
- [10] G. R. Golding, L. Bryden, P. N. Levett, R. R. McDonald and A. Wong, "Livestock-associated methicillin-resistant Staphylococcus aureus sequence type 398 in humans, Canada," *Emerging Infectious Diseases*, vol. 16, pp. 587-594, Apr. 2010.
- [11] H. F. Chambers, and F. R. Deleo, "Waves of resistance: Staphylococcus aureus in the antibiotic era", *Nature Reviews Microbiology*, vol. 7, no. 9, pp. 629-641, Sep. 2009.
- [12] T. Zhao, M. P. Doyle, J. Shere, and L. Garber, "Prevalence of enterohemorrhagic Escherichia coli O157:H7 in a survey of dairy herds", *Applied and Environmental Microbiology*, vol. 61 no. 4, pp. 1290-1293. April 1995.
- [13] P. Vos, G. Garrity, D. Jones, N. R. Krieg, W. Ludwig, F. A. Rainey, and W. Whitman, *Bergey's Manual of Systematic Bacteriology*; New York: Springer-Verlag, 2009, ch 4.
- [14] B. G. Weckman, and B. W. Catlin, "Deoxyribonuclease activity of Mirococci from clinical sources" *Journal of Bacteriology*, vol. 73, no. 6, pp. 747-753, June 1957.
- [15] P. Mugg, S. Seymour, and S. Clark, "Efficiency of the New Microgen® Staph ID for the Identification of Medically Important Staphylococcus species" United Kingdom: Microgen biproducts limited, U. K. 2003.
- [16] Danish Technical University, Multiplex PCR for the detection of the mecA gene and identification of Staphylococcus aureus, From the community reference library of antimicrobial resistance, DTU, 2014.
- [17] A. W. Bauer, W. M. Kirby, J. C. Sherris, and M. Truch, "Antibiotic susceptibility testing by a standardized single disk method" *American Journal of Clinical Pathology*, vol. 45, no. 4, pp. 493-496, April 1966.
- [18] J. A. Coghlan, J. G. Collee, R. Cruickshank, J. P. Duguid, R. R. Gilles, J. C. Gould, D. M. Green, and N. J. Hayward, *Medical Microbiology: The Practice of Medical Microbiology*, 12th ed. Churchill Livingstone, Edinburgh: 1975 pp. 587-588.
- [19] K. Haugum, J. Johansen, C. Gabrielsen, L. T. Brandal, K. Bergh, D. W. Ussery, F. Drablos, J. E. Afset, "Comparative Genomics to Delineate Pathogenic Potential in Non-O157 Shiga Toxin-Producing Escherichia coli (STEC) from Patients with and without Haemolytic Uremic Syndrome (HUS) in Norway," *PLoS ONE*, vol. 9(10), e111788. <https://doi.org/10.1371/journal.pone.0111788>, Oct. 2014.
- [20] M. d. Zohorul Islam, A. Musekiwa, K. Islam, S. Ahmed, S. Chowdhury, A. Ahad, "Regional Variation in the Prevalence of E. coli O157 in Cattle: A Meta-Analysis and Meta-Regression," *PLoS ONE*, vol 9(4), pp. e93299, Apr. 2014.
- [21] H. C. Lewis, K. Mølbak, C. Reese, F. M. Aarestrup, M. Selchau, M. Sorum, and R. L. Skov, "Pigs as source of methicillin-resistant Staphylococcus aureus CC398 infections in humans, Denmark", *Emerging Infectious Diseases*, vol. 14 no. 9, pp. 1383-1389, Sep. 2008.
- [22] E. de Boer, J. T. Zwartkruis-Nahuis, B. Wit, X. W. Huijsdens, A. J. de Neeling, T. Bosch, R. A. van Oosterom, A. Vila, A. E. Heuvelink, "Prevalence of methicillin-resistant Staphylococcus aureus in meat", *International Journal of Food Microbiology*, vol. 134, no. 1-2, pp. 52-56, Aug. 2009.
- [23] A. U. Nnachi, F. E. Emele, C. O. Ukaegbu, M. V. Agah, O. E. Udu-Ibiam, O.S. Chukwu, and M. M. Agwu, "Prevalence of methicillin-resistant Staphylococcus aureus in raw meat and meat handlers in Onitsha, Nigeria", *European Journal of Preventive Medicine*, vol. 2, no. 1, pp. 9-15, 2014.
- [24] V. Velasco, J. S. Sherwood, P. P. Rojas-Garcia, and C. M. Logue, "Multiplex Real-Time PCR for detection of Staphylococcus aureus, mecA and Panton-Valentine Leukocidin (PVL) genes from selective enrichments from animals and retail meat", *PLoS ONE*, vol. 9, no. 5, pp. e97617. May 2014.
- [25] R. Ali, K. Al-Achkar, A. Al-Mariri, and M. Safi, "Role of polymerase chain reaction in detection of antibiotic-resistant Staphylococcus aureus", *The Egyptian Journal of Medical Human Genetics*, vol. 15, pp. 293-298, June 2014.
- [26] A. Costa, I. Kay, and S. Palladino, "Rapid detection of mecA and nuc genes in Staphylococci by Real-Time multiplex PCR", *Diagnostic Microbiology and Infectious Diseases*, vol. 51, no. 1, pp. 13-17, Jan. 2005.
- [27] ECDC (European Centre for Disease Prevention and Control), EFSA (European Food Safety Authority) and EMA (European Medicines Agency), "ECDC/EFSA/EMA first joint report on the integrated analysis of the consumption of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from humans and food-producing animals," *Stockholm/parma/London ECDC/EFSA/EMA. EFSA Journal*vol.13, pp. 4006, Jan. 2015.