

Mineral Chemistry and Petrography of Lava Successions From Kepsut-Dursunbey Volcanic Field, NW Turkey: Implications For Magmatic Processes and Crystallization Conditions

Kamaci O. and Altunkaynak S.

Abstract—Kepsut-Dursunbey volcanic field (KDVF) is located in NW Turkey and contains various products of the post-collisional Neogene magmatic activity. Two distinct volcanic suites have been recognized; the Kepsut volcanic suite (KVS) and the Dursunbey volcanic suite (DVS). The KVS includes basaltic trachyandesite-basaltic andesite-andesite lavas and associated pyroclastic rocks. The DVS consists of dacite-rhyodacite lavas and extensive pumice-ash fall and flow deposits. Petrographical features (i.e. existence of xenocrysts, glomerocrysts, and mixing-compatible textures) and mineral chemistry of phenocryst assemblages of both suites provide evidence for magma mixing/AFC. Calculated crystallization pressures and temperatures give values of 5.7–7.0 kbar and 927–982 °C for the KVS and 3.7–5.3 kbar and 783–787°C for the DVS, indicating separate magma reservoirs and crystallization in magma chambers at deep and mid crustal levels, respectively. These observations support the establishment and evolution of KDVF magma system promoted by episodic basaltic inputs which may generate and mix with crustal melts.

Keywords—mineral chemistry, mixing, basaltic inputs, NW Turkey

I. INTRODUCTION

TURKEY is situated in the Alpine-Himalayan mountain belt between the continents of Eurasia and Africa and has a complex geology as a result of the collision between these major continents. In western Turkey, the terminal continental collision following the closure of the Tethys Ocean was followed by extensive magmatism which initiated around the Eocene and lasted almost continuously to prehistoric times ([1], [2], [3], [4]). Post collisional magmatism produced a wide range of magmatic associations, including plutonic, hypabyssal and volcanic rocks, which are closely related in space and time to tectonic processes ([5], [6], [7], [8], [9] and references there in).

The nature, origin, and evolution of the Neogene volcanism in NW Turkey have been the subject of many studies focusing

on the whole-rock geochemistry and petrogenesis of the magmatic rocks (e.g. [2], [10], [7], [5], [11], [12], [9], [13] etc.). However, there are very few published mineralogical and petrographical data on any of the lava successions of W Turkey. (e.g. [14], [15]).

The Kepsut-Dursunbey volcanic field is located in NW Turkey and comprise lavas represented by various compositions ranging from basaltic andesite to rhyolite and associated pyroclastics (Fig. 1). The knowledge on the volcanic rocks of study area is limited. Previous works have provided some introductory geological and geochemical information about the study area (e.g. [16], [17], [18], [19]). Studies establishing the volcanic successions, descriptions of the volcanic episodes, constraining evolution of magma conduit system and storage conditions are lacking in this area. Their petrogenetic characteristics are also poorly known.

This paper seeks to fill that gap by focusing on the mineral chemistry and petrography of lavas from the KDVF. In this study, as an alternative to using whole-rock geochemistry, mineral chemistry and petrographical features are used to determine crystallization conditions and magmatic processes occurring during early stages of magma evolution.

II. ANALYTICAL METHODS

Mineral compositions were determined with a JEOL JXA-8900R electron probe micro analyzer in the Electron Microanalysis and Imaging Laboratory at the University of Nevada, Las Vegas. Amphibole and plagioclase analyses were conducted with a 15 kV accelerating voltage, 10 nA current, and 10 μ m beam diameter. Mica analyses were conducted with a 15 kV accelerating voltage, 5 nA current, and 10 μ m beam diameter. Natural minerals were used as standards and counting times were 30 seconds on peak and background for all elements.

Omer Kamaci, Geological Engineering Department, Faculty of Mines, Istanbul Technical University, Istanbul, Turkey. (e-mail: kamaciom@itu.edu.tr).

Safak Altunkaynak, Geological Engineering Department, Faculty of Mines, Istanbul Technical University, Istanbul, Turkey. (e-mail: safak@itu.edu.tr).




Fig. 1 Geological map of the area studied

III. PETROGRAPHY AND MINERAL CHEMISTRY

The andesite, basaltic andesite and basaltic trachyandesite lavas of KVS are porphyritic in texture and composed of 20-35 modal % phenocrysts. The most common phenocryst assemblage observed in this rock group is Plagioclase (60-85%), biotite (3-8%), hornblende (5-8%), sanidine (3-5%), clinopyroxene (8-10 %) and orthopyroxene (1-5%). The matrix of the basaltic trachyandesites and basaltic andesites is commonly microlithic or glassy and shows flow-related parallel alignment of microlites (Trachytic texture) in groundmass. The glomerophyric textures, in which different phenocrysts such as plagioclase, clinopyroxenes and orthopyroxenes are clustered in irregular groups, are also seen in this group.

The rhyodacite lavas of DVS consist of sanidine (20-30%), quartz (20-25%), plagioclase (20-40%), biotite (3-5%) and hornblende (3-5%). Dacite lavas of this group contain quartz

hornblende (3-5%). (10-20%), plagioclase (55-80%), biotite (5-10%) and The groundmass is either glassy or microcrystalline. Many of rhyodacite lavas are devitrified and display spherulitic texture. Phenocrysts range in size up to a maximum of 4 mm.

Plagioclase is the main felsic mineral of KVS and DVS. It is euhedral, partly zoned with typical polysynthetic twinning. Compositions of the plagioclase phenocrysts and microlites of KVS and DVS lavas range from An₇₀ to An₅₉ and An₂₂ to An₃₆, respectively. Many plagioclase phenocrysts display patchy zones and oscillatory zoning. Some KVS samples have both euhedral and rounded plagioclases indicating crystallization in different stages (Fig.2). Rounded plagioclase phenocrysts with pre-resorption zoning display post-resorption reaction rims. Synneusis and sieve textured crystals are also common in both KVS and DVS (Fig. 3, 4). In some KVS samples acidic plagioclases together with quartz




Fig. 2 Coexistence of euhedral and rounded plagioclases in KVS

represent xenocrysts from felsic phase which are surrounded by biotite and amphibole microcrystals from mafic phase.

Clinopyroxene is present in all intermediate to basic rocks of KVS. Some of them are mantled by hornblende. Its composition ranges from Wo44 En47 Fs9 to Wo32 En54 Fs14., generally represented by endiopside, augite and rarely diopside. Orthopyroxenes are characterized by enstatite and their composition range from Wo2 En78 Fs20 to Wo1.5 En71 Fs27.5 in basic rocks of KVS.




Fig. 3 Synneusis texture in plagioclase phenocrysts of KVS lavas

Biotite is common mafic mineral of DVS lavas and seen as phenocrysts and interstitial microlites in some KVS lavas. Biotite crystals of DVS are rich in Ti (3.7-4.3 wt % TiO₂) and also in Al (13.76-14.32 wt. % Al₂O₃).



Fig. 4 Sieve textured plagioclase phenocrysts in DVS lavas

Amphiboles are common mafic mineral in both DVS and KVS. They are dominantly idiomorphic and represented by magnesio-hornblende in DVS lavas and tschermakitic hornblende- tschermakite, pargasite-pargasitic hornblende in KVS lavas (Fig 5).




Fig. 5 Amphibole compositions of DVS and KVS lavas (after [20])

IV. AMPHIBOLE-PLAGIOCLASE THERMOMETER

Electronmicroprobe data for the amphibole and plagioclase phenocrysts from the KVS and DVS were used to calculate temperature using the equations suggested by [21]. The temperatures were calculated using rim compositions of plagioclase-amphibole pairs and uncertainty in the geothermometer is considered to be ± 40 °C as proposed by Holland and Blundy (1994). Pressures for the mineral assemblages used in the temperature calculations were calculated using Al-in-hornblende method and equations proposed by different authors ([22], [23], [24], Table 1). The results obtained from the amphibole-plagioclase thermometer and Al-in-hornblende geobarometer are given in Table 1. Calculated pressures and temperatures using geothermometers/barometers give values of 5.7–7.0 kbar and 927–982 °C for the KVS and 3.7–5.3 kbar and 783–787 °C for the DVS, indicating crystallization in magma chambers at deep and mid crustal levels (21km and 12km, respectively).

TABLE I
TABLE SHOWING CALCULATED CRYSTALLIZATION TEMPERATURES AND PRESSURES

Sample no	18-OS	17-OS	16-OS	20-OS	22-OS
Group	KVS	KVS	KVS	DVS	DVS
Rock name	bas. and.	bas. and.	bas. and.	dacite	rhyodacite
T (°C)	980	982	927	783	787
P (kbar)					
HZ86*	5,7	6,6	6,1	4,8	3,7
H87*	6,0	7,0	6,4	5,0	3,8
S92*	6,1	6,9	6,5	5,3	4,2
P ave.(kbar)	5,90	6,90	6,30	4,90	3,90

HZ86*= Hammarstrom and Zen 1986, H87*= Hollister et al. 1987, S92*=Schmidt 1992.

V. DISCUSSION AND CONCLUSIONS

Two distinct volcanic suites have been recognized in KDVF; the Kepsut volcanic suite (KVS) and the Dursunbey volcanic suite (DVS). Both KVS and DVS are Early Miocene in age and representative for the post-collisional magmatism in western Anatolia (Turkey).

Mainly andesites, latites, basaltic andesites and pyroclastic rocks of KVS were formed during the early stage of volcanism in the study area. Without a major interruption in the volcanism, partly contemporaneously, KVS followed by an extensive felsic volcanism. Products of this phase are named as the Dursunbey volcanic suite (DVS) in this study. Rhyolitic domes, lavas and extensive pyroclastic units were formed during this volcanic activity. Pyroclastic rocks of DVS are represented by pyroclastic fall deposits and pyroclastic flow deposits. Pyroclastic fall deposits formed from ash fall, ash-block fall and pumice- ash fall deposits. Flow deposits are represented by ash-block flow deposits and ignimbrites, probably produced by explosive sub-plinian and ignimbrite forming eruptions. This was followed by basaltic trachyandesitic and basaltic dykes and lava flows in surrounding areas.

The lavas of KDVF display textural and chemical evidence for interaction of mafic and silicic magmas. This is evidenced by disequilibrium textures such as existence of rounded plagioclase phenocrysts with reaction rims (regrowth), hornblende-mantled clinopyroxenes (corona texture), synneusis, sieve textured plagioclase, patchy zones and oscillatory zoning in plagioclase phenocrysts. Basic lavas of KVS also contains xenocrysts represented by biotite mantled quartz and asidic plagioclase ovoids and glomerocrysts entrained from melt zones near base of the crust. The observed disequilibrium textures in both suites are consistent with models involving magma mixing/AFC proposed for the origin and evolution of other continental volcanic fields ([25], [26], [27], [28] etc.). A number of recent studies indicate that such textures are present on a regional scale within the volcanic fields of Western Anatolia (e.g. [5], [15], [16]) and were probably caused by the influx of a hotter basaltic magma into the crustal magma chambers. Therefore, the mineral chemistry and petrographical features discussed in this paper, rules out closed system fractionation and suggests that the

processes of mixing and AFC have played an important role in the development of the KDVF; KVS and DVS.

Mainly based on the geothermometry and barometry studies on both KVS and DVS lavas, we present a model for the evolution of KDVF. Calculated pressures and temperatures using a range of geothermometers/barometers give values of 5.7–7.0 kbar and 927–982 °C for the KVS and 3.7–4.2 kbar and 787–790°C for the DVS. Two levels of magma storage may be distinguished: deep reservoirs between 17 and 21 km and shallow reservoirs between 12 and 9 km for KVS and DVS, respectively, indicating separate magma reservoirs and crystallization in magma chambers at deep and mid crustal levels. A main deep magma reservoir served as a source for the intermediate to basic lavas of KVS. However, the existence of similar mixing-compatible (disequilibrium) textures in both KVS and DVS support the establishment and evolution of KDVF magma system promoted by episodic basaltic inputs which may generate and mix/mingle with crustal melts at different depths.

REFERENCES

- [1] Sengor, A. M. C., and Yilmaz, Y., "Tethyan evolution of Turkey: a plate tectonic approach," *Tectonophysics*, v. 75, p. 181-241, 1981.
- [2] Yilmaz, Y., "An approach to the origin of young volcanic rocks of western Turkey", in Şengör A.M. C. Ed., "Tectonic evolution of the Tethyan region," Kluwer, the Hague, p. 159-189, 1989.
- [3] Yilmaz, Y., "Comparision of young volcanic associations of western and eastern Anatolia under compressional regime; a review," *Journal of Volcanology and Geothermal Research*, v. 44, p. 69-87, 1990.
- [4] Harris, N.B.W., Kelley, S., and Okay, A.I., "Post-collisional magmatism and tectonics in northwest Anatolia," *Contributions to Mineralogy and Petrology*, v. 117, p. 241-252, 1994.
- [5] Altunkaynak, S., and Yilmaz, Y., "The mount Kozak magmatic complex, Western Anatolia," *Journal of Volcanology and Geothermal Research*, v. 85, p. 211-131, 1998.
- [6] Altunkaynak, S., and Yilmaz, Y., "The Kozak Pluton and its emplacement," *Geological Journal*, v. 34, p. 257-274, 1999.
- [7] Genc, S.C., "Evolution of the Bayramic magmatic complex, northwestern Anatolia," *Journal of Volcanology and Geothermal Research*, v. 85 no. 1-4, p. 233-249, 1998.
- [8] Yilmaz, Y., Genc, S.C., Karacik, Z., and Altunkaynak, S., "Two contrasting magmatic associations of NW Anatolia and their tectonic significance," *Journal of Geodynamics*, v. 31, p. 243-271, 2001.
- [9] Altunkaynak, S., and Dilek, Y., "Timing and nature of post collisional volcanism in Western Anatolia and Geodynamic Implications," in Dilek, Y. and Pavlides, S., eds., "Post-Collisional Tectonics and Magmatism of the Eastern Mediterranean Region and Asia," *Geological Society of America Special Paper*, v. 409, p. 321-351, 2006.
- [10] Gulec, N., "Crust-mantle interaction in western Turkey: implications from Sr and Nd isotope geochemistry of Tertiary and Quaternary volcanics," *Geological Magazine*, v. 23, p. 417-435, 1991.
- [11] Yilmaz, Y., Genc, S.C., Gurer, O.F., Bozcu, M., Yilmaz, K., Karacik, Z., Altunkaynak, S., Elmas, A., "When did the western Anatolian grabens begin to develop?," in Bozkurt, E., Winchester, J.A., Piper, J. A. D. eds., "Tectonics and Magmatism in Turkey and the Surrounding Area," *Geological Society, London, Special Publication*, v.173, p. 353-384, 2000.
- [12] Aldanmaz, E., Pearce, J., Thirwall, M.F., and Mitchell, J., "Petrogenetic evolution of late Cenozoic, post-collision volcanism in western Anatolia, Turkey," *Journal of Volcanology and Geothermal Research*, v. 102, p. 67-95, 2000.
- [13] Altunkaynak, S., "Collision-driven slab breakoff magmatism in northwestern Anatolia, Turkey," *Journal of Geology*, v.115, p. 63-82, 2007.
- [14] Aldanmaz, E., "Mineral-Chemical Constraints on the Miocene Calc-alkaline and Shoshonitic Volcanic Rocks of Western Turkey:

- Disequilibrium Phenocryst Assemblages as Indicators of Magma Storage and Mixing Conditions,” Turkish Journal of Earth Sciences (Turkish J. Earth Sci.), Vol. 15, pp. 47-73, 2006.
- [15] Altunkaynak, S., Rogers, N.W., Kelley, S.P., “Causes and effects of geochemical variations in late Cenozoic volcanism of the Foça volcanic centre, NW Anatolia, Turkey,” International Geology Review, Vol. 52, No. 4-6, p.579-607, 2010.
- [16] Ercan, T., Ergul, E., Akcoren, F., Cetin A., Granit, S ve Asutay, J., “Balıkesir-Bandırma arasının jeolojisi, Tersiyer volkanizmasının petrolojisi,” MTA Dergi 110, 113-130, 1990.
- [17] Akyuz, S., Manyas-Susurluk-Kepsut (Balıkesir) civarının jeolojisi, Doktora tezi, İ.T.Ü., Fen Bilim. Ens., 163-171, 1995.
- [18] Delaloye, M., Bingöl, E., “Granitoids from western and northwestern Anatolia: Geochemistry and modelling of Geodynamic evolution,” Int. Geol. Rev. 42, 241-268, 2000.
- [19] Boztug, D., Harlavan, Y., Jonckheere, R., Can, I., Sari, R., “Geochemistry and K-Ar cooling ages of the Ilica, Çataldağ (Balıkesir) and Kozak (Izmir) granitoids, west Anatolia, Turkey,” Geol. J. 44: 79-103, 2009.
- [20] Leake, B.E., “Nomenclature of amphiboles,” American Mineralogist, v. 63, p. 1023 -1052, 1978.
- [21] Holland, T.J.B. and Blundy, J., “Non-ideal interactions in calcic amphiboles and their bearing on amphibole-plagioclase thermometry,” Contributions to Mineralogy and Petrology 116,433-447, 1994.
- [22] Hammarstrom, J.M., Zen, E.A., “Aluminium in hornblende; an empirical igneous geobarometer,” American Mineral., 71, 1297-1313, 1986.
- [23] Hollister, L.S., Grissom, G.C., Peters, E.K., Stowell, H.H., Sisson, V.B., “Confirmation of the empirical correlation of Al in hornblende with pressure of solidification of calc-alkaline plutons,” American Mineralogist, Volume 72, pages 231-239, 1987.
- [24] Schmidt, W.S. “Amphibole composition in tonalite as a function of pressure: an experimental calibration of the Al-in hornblende barometer,” Contributions to Mineralogy and Petrology 110, 304-310, 1992.
- [25] Taylor, T.R., Vogel, T.A., and Wilband, J.T., “The composite dikes at Mount Desert Island, Maine; An example of coexisting acidic and basic magmas,” Jour. of Geol., 88, 433-444, 1980.
- [26] Vernon, R.H., “Microgranitoid enclaves in granites-globules of hybrid magma quenched in a plutonic environment,” Nature, 309, 438-439, 1984.
- [27] Hibbard, M.J. “The magma mixing origin of mantled feldspars,” Contributions to Mineralogy and Petrology 76, 158-170, 1981.
- [28] Kerr, A.C. “Mineral chemistry of the Mull-Morvern Tertiary lava succession, western Scotland,” Mineralogical Magazine, Vol. 62(3), pp. 295-312, 1998.