

Robust Numerical Scheme for Pricing American Options under Jump Diffusion Models

Salah Alrabeei, Mohammad Yousuf

Abstract—The goal of option pricing theory is to help the investors to manage their money, enhance returns and control their financial future by theoretically valuing their options. However, most of the option pricing models have no analytical solution. Furthermore, not all the numerical methods are efficient to solve these models because they have nonsmoothing payoffs or discontinuous derivatives at the exercise price. In this paper, we solve the American option under jump diffusion models by using efficient time-dependent numerical methods. Several techniques are integrated to reduce the computational complexity. Fast Fourier Transform (FFT) algorithm is used as a matrix-vector multiplication solver, which reduces the complexity from $\mathcal{O}(M^2)$ into $\mathcal{O}(M \log M)$. Partial fraction decomposition technique is applied to rational approximation schemes to overcome the complexity of inverting polynomial of matrices. The proposed method is easy to implement on serial or parallel versions. Numerical results are presented to prove the accuracy and efficiency of the proposed method.

Keywords—Integral differential equations, American options, jump-diffusion model, rational approximation.

I. INTRODUCTION

OPTION pricing models have been developed rapidly after the classical Black-Scholes model [1]. Just years later, empirical studies revealed that the Black-Scholes model is inconsistent with the market movements. Many studies have revealed to overcome these shortcomings, such as Lévy models and jump-diffusion models (JDM) [2], [3]. Unlike in the European options [4], [5], American options do not have closed-form to calculate. Thus, extensive research in numerical methods has been conducted and applied in valuating American options [6], [7]. Different methods have been used to approximate the linear complementary problems (LCPs), such as the penalty method proposed by [8], [9], operator splitting method [10], IMEX [11]–[14]. Khaliq et al. [15] developed numerical schemes based on Padè approximations of matrix exponential functions using exponential time differencing combined with Runge-Kutta (ETDRK) applied to the Black-Scholes model, Burgers equation, etc. (see [16], [17]). In this paper, we extend the work of Khaliq et al. [15] to apply it for American options with jump diffusion model. We use several methods to increase the computation speed, such as the FFT algorithm, partial fraction decomposition technique. This paper is organized as follows: the continuous problem is described in Section I. Discretization in space as well as in time are given in Sections

III and IV respectively. The fast Fourier transform algorithm (FFA) is described in Section V. Several numerical examples are given in Section VI to show the efficiency and accuracy of our algorithms. We finalized the paper by a short conclusion in Section VII.

II. JUMP DIFFUSION MODEL

Consider that $v(x, \tau)$ is the value price of the asset x at time τ satisfying the integral differential equation

$$v_\tau - \frac{1}{2}\sigma v_{xx} - (r - \frac{1}{2}\sigma^2 - \kappa\lambda)v_x + (r + \lambda)v - \lambda \int_{-\infty}^{\infty} v(z, \tau)\phi(z - x)dz = \tilde{p}(v, \tilde{v}) \quad (1)$$

with the following are the initial and boundary conditions:

- Call option

$$\begin{cases} v(x, 0) = \max(Ee^x - E), \\ v(X_{min}, \tau) = 0, \\ v(X_{max}, \tau) = Ee^{x_{max}} - E, \end{cases}$$

- Put option

$$\begin{cases} v(x, 0) = \max(E - Ee^x), \\ v(X_{min}, \tau) = E, \\ v(X_{max}, \tau) = 0, \end{cases}$$

where $\tilde{v}(x, \tau)$ is the payoff received at the exercising time $\tau \leq T$ and $\tilde{p}(V, \tilde{V})$ is the penalty term [18] given by

$$p(v, \tilde{v}) = \frac{1}{\epsilon} \max \left\{ v(x, \tau) - \tilde{v}(x, \tau), 0 \right\}, \quad 0 \leq \epsilon \leq 1$$

where E is the exercise price, σ is the volatility and r is the rate of interest, λ is the Poisson intensity and κ is the expectation of the impulse function. For the boundary conditions, X_{min} and X_{max} are boundaries of the truncated domain $\Omega \in (-\infty, \infty)$ and ϕ is the density function of the normal distribution function with mean μ , variance δ^2 and $\kappa = e^{(\mu - \frac{\delta^2}{2})} - 1$.

Salah Alrabeei: Department of Computing, Mathematics, and Physics, Western Norway University of Applied Sciences, Bergen, Norway (e-mail: salah.alrabeei@hvl.no).

Mohammad Yousuf: Department of Mathematics and Statistics, King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia (e-mail: myousuf@kfupm.edu.sa).

III. SPATIAL DISCRETIZATION

The infinite space is truncated to $\Omega = [X_{min}, X_{max}]$ [19]. Then we discretize the differentiation term by the 2nd order central finite difference and fourth order Chebyshev Spectral Method [20] whereas the integral term is approximated the 2nd order composite Trapezoidal Rule and a fourth order quadrature rule at each subinterval in space. Therefore, the PIDE(1) can be written as a semi-linear system of ODEs given by

$$v'(\tau) + Av = \lambda F(v, \tau) \tag{2}$$

where A is given by

$$A = \frac{1}{2}\sigma^2 D_M^2 + \left(r - \frac{1}{2}\sigma^2 - \kappa\lambda \right) D_M - (r + \lambda)I$$

in case of Chebyshev spectral methods, where D_M and D_M^2 are the first and second Chebyshev spectral differentiation matrices respectively whereas, in case of the FDM, A is given by

$$A = \text{tridiag} \left(\frac{c_1}{2h^2} - \frac{c_2}{2h}, -\frac{c_1}{h^2} - r - \lambda, \frac{c_1}{2h^2} + \frac{c_2}{2h} \right)$$

where

$$c_1 = \sigma^2 \quad \text{and} \quad c_2 = r - \lambda\kappa - \frac{\sigma^2}{2}$$

Approximating the integral term by either composite Trapezoidal Rule, which is less accurate or Clenshaw-Curtis Quadrature, which is higher accurate, will end up with either tridiagonal Toeplitz matrix which is computationally cheaper or a dense matrix which is computationally exhaustive respectively. The semi-linear function F is the approximated integral term and the penalty term of the PIDE(1).

IV. FULL DISCRETIZATION

For the semi-discretized system of the ODEs given in (2), define $k \geq 0$ by the time step size, $\tau_n = nk, 0 \leq n \leq N$. Thus, using Duhamel principle [15], the semi-discretized system of the ODEs has exact solution given by

$$v(\tau) = e^{-kA}v(\tau) + \int_0^\tau e^{-kA(\tau-\nu)}F(v(\nu), \nu)d\nu, \tag{3}$$

by changing variable, $\nu - \tau = ks$ and putting $v_n = v(\tau_n)$, we get a simpler form of the exact solution

$$v_{n+1} = e^{-kA}v_n + k \int_0^1 e^{-kA(1-s)}F(v(\tau_n + ks), \tau_n + ks)ds, \tag{4}$$

Several time stepping schemes were developed (see [21], [22]) to approximate the recurrence formula (4). However, their schemes require to invert matrix higher order polynomials which is computationally expensive. Later, [23] used partial fraction form of rational approximations to approximate the matrix exponential functions arising in the exact solutions.

A. Padé Approximation

Padé approximation, type of rational approximation, is an important tool to approximate matrix exponential functions. It is named after the French mathematician Henri Padé (1863-1953). Padé schemes are the approximants derived by a ratio of two power series approximations. Due to the rational form of Padé approximations, it is better than Taylor expansions when approximating functions containing poles. Following [24], Padé approximation of order $(n + m + 1)$ of the exponential function e^{-x} is given by

$$R_m^n(x) = \frac{P_m^n(x)}{Q_m^n(x)} + O(x^{n+m+1})$$

where $P_m^n(x)$ and $Q_m^n(x)$ are polynomials of order n and m respectively given by

$$P_m^n(x) = \sum_{i=0}^n \frac{(m+n+i)n!}{(m+n)!i!(n-i)!} (-x)^i$$

$$Q_m^n(x) = \sum_{j=0}^m \frac{(m+n+j)n!}{(m+n)!j!(m-i)!} (x)^j$$

with the property $R_m^n(x) = e^{-x} + O(x^{n+m+1})$ when $x \rightarrow 0$.

Illustrative examples: The following functions are different lower and higher orders Padé approximations of e^{-x} :

$$\begin{aligned} R_1^0(x) &= (1+x)^{-1} && \text{(Backward Euler)} \\ R_1^1(x) &= (1-\frac{1}{2}x)(1+\frac{1}{2}x)^{-1} && \text{(Crank-Nicolson)} \\ R_2^1(x) &= (1-\frac{1}{3}x)(1+\frac{2}{3}x+\frac{1}{6}x^2)^{-1} \\ R_4^0(x) &= 24(24+24x+12x^2+4x^3+x^4)^{-1} \end{aligned}$$

B. Partial Fraction form Padé Approximation

The ETDRK schemes mentioned in the previous section contain lower and higher order polynomials of matrices which cause computational difficulties. In this regard, [25] and [26] have made important contributions to address this issue. They used the partial fraction technique to implement the Padé schemes. Not only this, but also they implemented efficient serial and parallel algorithms. Yousuf et al. [27] developed algorithms to implement diagonal and damping subdiagonal schemes and obtained the following version of schemes. Although we are interest only on two particular types of Padé approximations, for the sake of generalization, we shall give schemes of (n,m) - Padé in general for any positive integer numbers n, m .

Case1: $n < m$,

$$R_m^n(x) = \sum_{j=1}^{q_1} \frac{\omega_j}{x - c_j} + 2 \sum_{j=q_1+1}^{q_1+q_2} \Re \left(\frac{\omega_j}{x - c_j} \right)$$

Case2: $n = m$,

$$R_m^m(x) = (-1)^m + \sum_{j=1}^{q_1} \frac{\omega_j}{x - c_j} + 2 \sum_{j=q_1+1}^{q_1+q_2} \Re \left(\frac{\omega_j}{x - c_j} \right)$$

where q_1 and q_2 are the number of real and non-real poles \bar{c}_j of \bar{R}_m^n .

Therefore, our fully discretized scheme is given by

$$v_{n+1} = R_4^0(kA)v_n + \mathcal{P}_1(kA)F(v_n, t_n) + \mathcal{P}_2(kA)(F(\alpha_n, t_n + k/2) + F(\beta_n, t_n + k/2)) + \mathcal{P}_3(kA)F(\gamma_n, t_n), \quad (5)$$

where

$$\alpha_n = \bar{R}_4^0(kA)v_n + \bar{\mathcal{P}}(kA)F(v_n, t_n),$$

$$\beta_n = \bar{R}_4^0(kA)v_n + \bar{\mathcal{P}}(kA)F(\alpha_n, t_n + k/2),$$

$$\gamma_n = \bar{R}_4^0(kA)v_n + \bar{\mathcal{P}}(kA)\left(2F(\beta_n, t_n + k/2) - F(v_n, t_n)\right),$$

and

$$R_4^0(x) = 2\Re\left(\frac{\omega_1}{x - c_1}\right) + 2\Re\left(\frac{\omega_2}{x - c_2}\right),$$

$$\mathcal{P}_i = 2\Re\left(\frac{\omega_{i1}}{x - c_1}\right) + 2\Re\left(\frac{\omega_{i2}}{x - c_2}\right), \quad i = 1, 2, 3.$$

$$\bar{R}_4^0(x) = 2\Re\left(\frac{\bar{\omega}_1}{x - \bar{c}_1}\right) + 2\Re\left(\frac{\bar{\omega}_2}{x - \bar{c}_2}\right),$$

and

$$\mathcal{P}(x) = 2\Re\left(\frac{\Omega_1}{x - \bar{c}_1}\right) + 2\Re\left(\frac{\Omega_2}{x - \bar{c}_2}\right),$$

where the poles c_i and c^-_i and the corresponding weights ω_i , ω^-_i , ω_{ij} and Ω_i , $i = 1, 2$ and $j = 1, 2, 3$ are given by

- $c_1 = -1.72944423106769 + i0.888974376121862.$
- $c_2 = -0.2705557689322 - i2.50477590436244.$
- $\omega_1 = 0.541413348429182 - i1.58885918222330.$
- $\omega_2 = -0.541413348429154 - i0.248562520866115.$
- $\omega_{11} = 0.244153693956274 - i0.0497524711964030.$
- $\omega_{12} = -0.244153693956268 - i0.0750708534900480.$
- $\omega_{21} = -0.0240066687966667 - i0.210771761184790.$
- $\omega_{22} = 0.0240066687966698 + i0.110830774318527.$
- $\omega_{31} = 0.473042583717175 + i0.293424221840328.$
- $\omega_{32} = 0.0269574162828241 - i0.165188084403066.$
- $\bar{c}_1 = -3.45888846213543 - i1.77794875224371.$
- $\bar{c}_2 = -0.541111537864595 - i5.00955180872487.$
- $\bar{\omega}_1 = 1.08282669685827 + i3.17771836444659.$
- $\bar{\omega}_2 = -1.08282669685831 - i0.497125041732246.$
- $\Omega_1 = -0.621169602486758 - i0.599415294095229.$
- $\Omega_2 = 0.121169602486770 - i0.203064159380992.$

C. Stability Analysis

Consider the nonlinear ODE,

$$v_t = cv + F(v) \quad (6)$$

where $F(v)$ is the nonlinear term. We assume that there exist a fixed point $v_0 = v(t_0)$, such that $cv_0 + F(v_0) = 0$. We linearize about the fixed point to lead to

$$v_t = cv + \lambda v. \quad (7)$$

where v becomes the perturbation to v_0 , whereas, $\lambda = F'(v_0)$. If $\Re(c + \lambda) < 0$, then the fixed point v_0 is stable [21].

To obtain the stability region of the numerical methods, we first denote $\xi = \lambda k$ and $\eta = ck$, where k is the time step-size,




Fig. 1 Stability regions of (0,4)-Padé scheme in the complex ξ -plane

then we apply (5) to the ODE (6) leading to a recurrence relation involving v_n and v_{n+1} .

$$\frac{v_{n+1}}{v_n} = r(\xi, \eta) = \frac{c_0 + c_1\xi + c_2\xi^2 + c_3\xi^3 + c_4\xi^4}{D} \quad (8)$$

where

$$D = (24 - 24\eta + 12\eta^2 - 4\eta^3 + \eta^4)(384 - 192\eta + 48\eta^2 - 8\eta^3 + \eta^4)^3$$

where c , λ , ξ and η are complex numbers. Following [21], for a better and useful method, the stability regions grow as $|\eta|$ becomes larger. Therefore, we fix η with several negative real values, $\eta = 0$, $\eta = -5$, $\eta = -10$ and $\eta = -20$, in the complex ξ -plane.

In Fig. 1, we show the stability region in the complex ξ -plane for different negative real η , $\eta = 0$, $\eta = -5$, $\eta = -10$ and $\eta = -20$. When $\eta \rightarrow 0$ the stability region tends to the fourth-order Runge-Kutta scheme; and, as η decreases from -10 to -20 the region grows. This region gives an indication of the stability of the (0,4)-Padé.

V. THE FFT ALGORITHM

Thanks to the approximation of the integral part is a Toeplitz matrix, we can reduce the cost of vector-matrix multiplication using what is so-called Fast Fourier Transform (FFT) algorithm [28]

Definition 1. The Discrete Fourier Transform (DFT) is a Transform on \mathbf{C}^m given as

$$[D\{b\}]_j = \frac{1}{\sqrt{m}} \sum_{k=0}^{m-1} b_k e^{-i2\pi jk/m}, \quad j = 0, 1, \dots, m-1 \quad (9)$$

where b is a vector and D is a Fourier matrix $\in \mathbf{C}^{m \times m}$

Whereas the inverse DFT of a vector b is given by

$$[D^{-1}\{b\}]_j = \frac{1}{\sqrt{m}} \sum_{k=0}^{m-1} b_k e^{i2\pi jk/m}, \quad j = 0, 1, \dots, m-1 \quad (10)$$

The use FFT algorithm is to evaluate the The Discrete Fourier Transform (DFT) of a vector b of length M whose operations cost is $O(M^2)$ with a cost of $O(M \log M)$ operation [29].

Definition 2. A square matrix $[A]_{M \times M}$ is called a Toeplitz Matrix if it has the form

$$A = \begin{bmatrix} a_0 & a_{-1} & \dots & a_{2-M} & a_{1-M} \\ a_1 & a_0 & a_{-1} & \ddots & a_{1-M} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ a_{M-2} & \ddots & a_1 & a_0 & a_{-1} \\ a_{M-1} & a_{M-2} & \dots & a_1 & a_0 \end{bmatrix}$$

We may generate the toeplitz matrix using the mathematical software Matlab by calling $toeplitz(a)$, where $a = (a_{M-1}, a_{M-2}, \dots, a_0, a_{-1}, a_{-2}, \dots, a_{1-M})$

Definition 3. A Circulant matrix $[C]_{M \times M}$ is special case of a Toeplitz matrix when for each of its rows is a right cyclic shift of the row preceding it, (i.e)

$$C = \begin{bmatrix} a_0 & a_{M-1} & \dots & a_2 & a_1 \\ a_1 & a_0 & a_{M-1} & \ddots & a_2 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ a_{M-2} & \ddots & a_1 & a_0 & a_{M-1} \\ a_{M-1} & a_{M-2} & \dots & a_1 & a_0 \end{bmatrix}$$

To evaluate the vector-matrix multiplication $[A]_{M \times M} [b]_{M \times 1}$, we embed [28] the matrix A into a circulant matrix $[C]_{(2M-1) \times (2M-1)}$, let $\hat{b} = [b, 0, \dots, 0]$ where \hat{b} is of size $2M - 1$. Next step is to apply the DFT transform to the vector \hat{b} and vector c where c is the first row and column of the circulant matrix C , then apply the inverse DFT transform to the product of transformed two vectors. (i.e)

$$v = \text{ifft}(\text{fft}(c) \cdot \text{fft}(\hat{b})) \tag{11}$$

Finally we extract the first M elements of the vector v to be the desired result.

VI. NUMERICAL EXPERIMENTS

We test performance of our scheme by showing its efficiency and accuracy. Our concern is not only the convergence, but also on the computational cost.

A. Convergence Test

To compare convergence of finite difference method and the spectral method, we shall use the second order central finite difference method and the Chebychev spectral method to discretize the domain Ω in space. Regarding to the integral term (jump part), we use composite trapezoidal rule when using finite difference method. Whereas, the Clenshaw-Curtis Quadrature is used to approximate the integral term when using the spectral method. We consider the truncated domain $\Omega = [-2.2, 2.2]$, with $E = 100$, $\sigma = 0.15$, $\delta = 0.25$ $r = 0.04$, $\lambda = 1, \mu = 0, T = 1$ and $\epsilon = 0.01$.

TABLE I
ORDER OF CONVERGENCE IN SPACE OF FD AND SPECTRAL METHODS AT A FIXED NUMBER OF TIME-STEPS $N = 100$

M	CENTRAL FD METHOD			SPECTRAL METHOD		
	VALUE	ERROR	ORDER	VALUE	ERROR	ORDER
40	12.49678	-	-	12.71511	-	-
80	12.66399	1.67208E-01	-	12.71794	2.83265E-03	-
160	12.70438	4.03889E-02	2.04960	12.71811	1.74943E-04	4.01719
320	12.71462	1.02421E-02	1.97944	12.71813	1.09423E-05	3.99889
640	12.71723	2.60203E-03	1.97680	12.71813	6.84221E-07	3.99931

From Table I, it is noticed that the second and fourth order convergence are achieved when using the second order central finite difference method and spectral method respectively in the spatial discretizations.

B. Efficiency Test

In this experiment we compare the the efficiency our scheme when using two different spatial discretization methods as well as two different integral approximation methods.

TABLE II
COMPUTATIONAL COSTS (TIME IN SECONDS) REQUIRED WHEN USING FINITE DIFFERENCE METHOD VS CHEBYCHEV SPECTRAL METHOD

STEPS M=N	CENTRAL FD METHOD		SPECTRAL METHOD	
	VALUE	CPU	VALUE	CPU
40	14.84558872	0.040	15.03228280	0.041
80	14.98433811	0.100	15.03467073	0.121
160	15.02087991	0.120	15.03467073	0.184
320	15.03080592	0.418	15.03497510	1.241
640	15.03362563	1.659	15.03498484	9.471
1280	15.03449027	7.661	15.03498720	73.595

It can be observed from Table II that the required computational time is almost the same in the two different methods. However, when the discretisation becomes finer, which makes the matrix is larger, the first method (FDM) requires much less time compared with the spectral method. The reason behind that FFT method is used as a matrix-vector multiplication in the first method because we have a Toeplitz-tridiagonal matrix. However, when applying the spectral method, we will have to dense large matrix-multiplication in each time iteration.

VII. CONCLUSION

We have developed a robust and stable time-dependent method for pricing American options with Morton's jump diffusion model. We use different useful techniques to reduce the computational complexity. We have used the rational Padè approximation to approximate the exponential function of matrices, then used its partial fraction decomposition technique to overcome inverting polynomials of matrices. We also applied the FFT methods to as a matrix-vector multiplication solver. We finally gave numerical examples to test our algorithm.

ACKNOWLEDGMENT

This work was supported to be published by department of Computing, Mathematics and Physics, Western Norway

University of Applied Sciences whose support is gratefully acknowledged.

The work of the second author was partially supported by the Internal Project IN141026, King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia.

REFERENCES

- [1] F. Black and M. Scholes, "The pricing of options and corporate liabilities," *The journal of political economy*, pp. 637–654, 1973.
- [2] R. C. Merton, "Option pricing when underlying stock returns are discontinuous," *Journal of financial economics*, vol. 3, no. 1-2, pp. 125–144, 1976.
- [3] S. G. Kou, "A jump-diffusion model for option pricing," *Management science*, vol. 48, no. 8, pp. 1086–1101, 2002.
- [4] L. Andersen and J. Andreasen, "Jump-diffusion processes: Volatility smile fitting and numerical methods for option pricing," *Review of Derivatives Research*, vol. 4, no. 3, pp. 231–262, 2000.
- [5] K. I. Amin, "Jump diffusion option valuation in discrete time," *The journal of finance*, vol. 48, no. 5, pp. 1833–1863, 1993.
- [6] S. Hussain and N. Rehman, "Regularity of the american option value function in jump-diffusion model," *Journal of Computational Analysis and Applications*, vol. 22, no. 2, pp. 286–297, 2017.
- [7] Y. Chen, W. Wang, and A. Xiao, "An efficient algorithm for options under merton's jump-diffusion model on nonuniform grids," *Computational Economics*, vol. 53, no. 4, pp. 1565–1591, 2019.
- [8] X. Gan, Y. Yang, and K. Zhang, "A robust numerical method for pricing american options under kou's jump-diffusion models based on penalty method," *Journal of Applied Mathematics and Computing*, vol. 62, no. 1-2, pp. 1–21, 2020.
- [9] D. Tangman, A. Gopaul, and M. Bhuruth, "Exponential time integration and chebyshev discretisation schemes for fast pricing of options," *Applied Numerical Mathematics*, vol. 58, no. 9, pp. 1309–1319, 2008.
- [10] L. Boen and K. J. in't Hout, "Operator splitting schemes for american options under the two-asset merton jump-diffusion model," *Applied Numerical Mathematics*, 2020.
- [11] S. Salmi and J. Toivanen, "Imex schemes for pricing options under jump-diffusion models," *Applied Numerical Mathematics*, vol. 84, pp. 33–45, 2014.
- [12] M. K. Kadalbajoo, L. P. Tripathi, and A. Kumar, "Second order accurate imex methods for option pricing under merton and kou jump-diffusion models," *Journal of Scientific Computing*, vol. 65, no. 3, pp. 979–1024, 2015.
- [13] R. Mollapourasl, A. Fereshtian, H. Li, and X. Lu, "Rbf-pu method for pricing options under the jump-diffusion model with local volatility," *Journal of Computational and Applied Mathematics*, vol. 337, pp. 98–118, 2018.
- [14] K. Kazmi, "An imex predictor-corrector method for pricing options under regime-switching jump-diffusion models," *International Journal of Computer Mathematics*, vol. 96, no. 6, pp. 1137–1157, 2019.
- [15] A. Khaliq, B. Wade, M. Yousuf, and J. Vigo-Aguiar, "High order smoothing schemes for inhomogeneous parabolic problems with applications in option pricing," *Numerical Methods for Partial Differential Equations*, vol. 23, no. 5, pp. 1249–1276, 2007.
- [16] H. P. Bhatt and A.-Q. M. Khaliq, "Fourth-order compact schemes for the numerical simulation of coupled burgers' equation," *Computer Physics Communications*, vol. 200, pp. 117–138, 2016.
- [17] H. P. Bhatt and A. Khaliq, "A compact fourth-order l-stable scheme for reaction-diffusion systems with nonsmooth data," *Journal of Computational and Applied Mathematics*, vol. 299, pp. 176–193, 2016.
- [18] R. Zvan, P. Forsyth, and K. Vetzal, "Penalty methods for american options with stochastic volatility," *Journal of Computational and Applied Mathematics*, vol. 91, no. 2, pp. 199–218, 1998.
- [19] Y. dHalluin, P. A. Forsyth, and G. Labahn, "A penalty method for american options with jump diffusion processes," *Numerische Mathematik*, vol. 97, no. 2, pp. 321–352, 2004.
- [20] L. N. Trefethen, *Spectral methods in MATLAB*, vol. 10. Siam, 2000.
- [21] S. Cox and P. Matthews, "Exponential time differencing for stiff systems," *Journal of Computational Physics*, vol. 176, no. 2, pp. 430–455, 2002.
- [22] A.-K. Kassam and L. N. Trefethen, "Fourth-order time-stepping for stiff pdes," *SIAM Journal on Scientific Computing*, vol. 26, no. 4, pp. 1214–1233, 2005.
- [23] A. Khaliq, J. Martin-Vaquero, B. Wade, and M. Yousuf, "Smoothing schemes for reaction-diffusion systems with nonsmooth data," *Journal of Computational and Applied Mathematics*, vol. 223, no. 1, pp. 374–386, 2009.
- [24] V. Thomée, *Galerkin finite element methods for parabolic problems*, vol. 1054. Springer, 1984.
- [25] E. Gallopoulos and Y. Saad, "On the parallel solution of parabolic equations," in *Proceedings of the 3rd international conference on Supercomputing*, pp. 17–28, ACM, 1989.
- [26] A. Khaliq, E. Twizell, and D. Voss, "On parallel algorithms for semidiscretized parabolic partial differential equations based on subdiagonal padé approximations," *Numerical Methods for Partial Differential Equations*, vol. 9, no. 2, pp. 107–116, 1993.
- [27] M. Yousuf, A. Khaliq, and B. Kleefeld, "The numerical approximation of nonlinear black-scholes model for exotic path-dependent american options with transaction cost," *International Journal of Computer Mathematics*, vol. 89, no. 9, pp. 1239–1254, 2012.
- [28] C. R. Vogel, *Computational methods for inverse problems*, vol. 23. Siam, 2002.
- [29] A. Almendral and C. W. Oosterlee, "Numerical valuation of options with jumps in the underlying," *Applied Numerical Mathematics*, vol. 53, no. 1, pp. 1–18, 2005.



Salah A. Alrabeai: received the B.S. degree in applied Mathematics from Dhamar University, Dhamar, Yemen . He received his M.Sc in applied Mathematics from King Fahd University of Petroleum and Minerals, Dhahran, Saudi Arabia . He is currently pursuing the Ph.D degree in engineering computing in the western Norway University of Applied Sciences, Bergen, Norway. His research interests include numerical analysis and scientific computing, ecological modeling, modeling and simulations and dynamical systems.



Mohammad Yousuf: received his PhD degree in Applied Mathematics from University of Wisconsin-Milwaukee, Milwaukee, Wisconsin, USA. He has been working as an assistant professor in department of Mathematics and Statistics, King Fahd University of Petroleum and Minerals, Dhahran 31261, Saudi Arabia. His research of interest is in using Numerical schemes for homogeneous as well as inhomogeneous linear and semi-linear parabolic partial differential equations arising in actuarial science, financial mathematics.