

Strict Stability of Fuzzy Differential Equations by Lyapunov Functions

Mustafa Bayram Gücen, Coşkun Yakar

Abstract—In this study, we have investigated the strict stability of fuzzy differential systems and we compare the classical notion of strict stability criteria of ordinary differential equations and the notion of strict stability of fuzzy differential systems. In addition that, we present definitions of stability and strict stability of fuzzy differential equations and also we have some theorems and comparison results. Strict Stability is a different stability definition and this stability type can give us an information about the rate of decay of the solutions. Lyapunov's second method is a standard technique used in the study of the qualitative behavior of fuzzy differential systems along with a comparison result that allows the prediction of behavior of a fuzzy differential system when the behavior of the null solution of a fuzzy comparison system is known. This method is a usefull for investigating strict stability of fuzzy systems. First of all, we present definitions and necessary background material. Secondly, we discuss and compare the differences between the classical notion of stability and the recent notion of strict stability. And then, we have a comparison result in which the stability properties of the null solution of the comparison system imply the corresponding stability properties of the fuzzy differential system. Consequently, we give the strict stability results and a comparison theorem. We have used Lyapunov second method and we have proved a comparison result with scalar differential equations.

Keywords—Fuzzy systems, fuzzy differential equations, fuzzy stability, strict stability.

I. INTRODUCTION

STABILITY of differential equations is very important topic for math, engineering and related areas [1],[2], [6]-[15], [17]-[21], [24]-[40], [46]-[48]. Strict Stability is a different stability definition and this stability can give us some information about the rate of decay of the solutions [22]. Strict stability of differential equations, strict stability in impulsive functional differential equations and strict stability of differential equations with initial time difference studied and new results obtained [22], [38]-[45], [49].

In this paper, we studied the definitions and comparison theorems of strict stability of fuzzy differential equations.

II. PRELIMINARIES

First of all, we must define the fundamental properties of fuzzy sets, fuzzy derivative and fuzzy calculus for the fuzzy differential equations and the stability of fuzzy differential equations. For fuzzy sets, we take as the base space R^n . The fuzzy set $u \in E^n$ is a function $u : R^n \rightarrow [0,1]$. Where E^n is the space of all fuzzy subsets of R^n and $E^n = \{u : R^n \rightarrow [0,1] | u \text{ satisfies } (i) \text{ to } (iv)\}$ below

M. B. Gücen is with the Department of Mathematics, Yildiz Technical University, Esenler, stanbul, 34200 Turkey (e-mail: mgucen@yildiz.edu.tr).

C. Yakar is with the Department of Mathematics, Gebze Technical University, Gebze, Kocaeli, 41400 Turkey (e-mail: cyakar@gtu.edu.tr).

(i) u is normal if there exists an $x_0 \in R^n$ such that $u(x_0) = 1$;

(ii) u is fuzzy convex if for $x, y \in R^n$ and $0 \leq \lambda \leq 1$ such that

$$u(\lambda x + (1 - \lambda)y) \geq \min[u(x), u(y)];$$

(iii) u is upper semi-continuous;

(iv) $[u]^0 = \{x \in R^n : u(x) > 0\}$ is compact subset of R^n .

The function $u(x)$ for $x \in R^n$ taking values in $[0,1]$. Now we state the following membership grade of the $u(x)$, $u(x) = 0$ meaning to nonmembership, $0 < u(x) < 1$ to partial membership and $u(x) = 1$ to full membership.

Let K_c^n be the family of all nonempty compact, convex subsets of R^n .

For $0 < \alpha \leq 1$, we denote the α -level sets as $[u]^\alpha = \{x \in R^n : u(x) \geq \alpha\}$. Then from (i) to (iv), it follows that $[u]^\alpha \in K_c^n$ for $0 \leq \alpha \leq 1$.

Let $d_H(A, B)$ be the Hausdorff distance between the sets $A, B \in K_c^n$. Then we define

$$d[u, v] = \sup_{0 \leq \alpha \leq 1} d_H([u]^\alpha, [v]^\alpha)$$

which defines a metric in E^n and (E^n, d) is a complete metric space. We have the following properties of $d[u, v]$:

$$(i) \quad d[u + w, v + w] = d[u, v],$$

$$(ii) \quad d[u, v] = d[v, u],$$

$$(iii) \quad d[\lambda u, \lambda v] = |\lambda|d[u, v],$$

$$(iv) \quad d[u, v] \leq d[u, w] + d[w, v],$$

for all $u, v, w \in E^n$ and $\lambda \in R$.

For $x, y \in E^n$ if there exists a $z \in E^n$ such that $x = y + z$, then z is called the H -difference of x and y and is denoted by $x - y$.

Definition 1: Let $F : (a, b) \rightarrow E^n$, $t_0 \in (a, b)$ and F is differentiable at t_0 . If there exists $F'(t_0) \in E^n$ such that

(i) For all $h > 0$ sufficiently small there exist $F(t_0 + h) - F(t_0)$, $F(t_0) - F(t_0 - h)$ and the limits (in (E^n, d))

$$\begin{aligned} \lim_{h \rightarrow 0^+} (1/h)(F(t_0 + h) - F(t_0)) &= \lim_{h \rightarrow 0^+} (1/h)(F(t_0) - F(t_0 - h)) \\ &= F'(t_0) \end{aligned} \quad (1)$$

under the similar conditions, and other definition is,

(ii) For all $h < 0$ sufficiently small there exist $F(t_0 + h) - F(t_0)$, $F(t_0) - F(t_0 - h)$ and the limits (in (E^n, d))

$$\begin{aligned} \lim_{h \rightarrow 0^-} (1/h)(F(t_0 + h) - F(t_0)) &= \lim_{h \rightarrow 0^-} (1/h)(F(t_0) - F(t_0 - h)) \\ &= F'(t_0) \end{aligned} \quad (2)$$

For details please see [3]-[5].

III. FUNDAMENTAL DEFINITIONS AND THEOREMS OF STRICT STABILITY CRITERIA OF FUZZY DIFFERENTIAL SYSTEMS

Consider the initial value problem for the fuzzy differential equations

$$u' = f(t, u), u(t_0) = u_0 \text{ for } t \geq t_0 \geq 0 \quad (3)$$

$$u' = f(t, u), u(\tau_0) = v_0 \text{ for } t \geq t_0 \geq 0 \quad (4)$$

where $f \in C[R_+ \times S\rho, E^n]$, $S\rho = [u \in E^n : d[u, \tilde{0}] < \rho < \infty]$, $f(t, \tilde{0}) \equiv \tilde{0}$ for $t \geq 0$.

Before we can establish our comparison theorem and Lyapunov functions with strict stability criteria of fuzzy differential equations we need to introduce the following definitions.

Definition 2: The trivial solution of the system (3) is said to be strictly stable, if given $\epsilon_1 > 0$ and $t_0 \in R_+$, there exist a $\delta_1 = \delta_1(\epsilon_1, t_0) > 0$ such that $d[u_0, \tilde{0}] < \delta_1$ implies $d[u(t), \tilde{0}] < \epsilon_1$, $t \geq t_0$, and for every $0 < \delta_2 \leq \delta_1$, there exists an $0 < \epsilon_2 < \delta_2$ such that

$$\delta_2 < d[u_0, \tilde{0}] \text{ implies } \epsilon_2 < d[u(t), \tilde{0}] \text{ for } t \geq t_0$$

If δ_1, δ_2 and ϵ_2 in Definition 2 are independent of t_0 , then the trivial solution of the system (3) is said to be strictly uniformly stable for $t \geq t_0$.

Definition 3: The trivial solution of the system (3) is said to be strictly attractive, if given $\alpha_1 > 0$, $\epsilon_1 > 0$ and $t_0 \in R_+$, for every $\alpha_2 \leq \alpha_1$, there exists $\epsilon_2 < \epsilon_1$ and $T_1 = T_1(\epsilon_1, t_0)$ and $T_2 = T_2(\epsilon_2, t_0)$ such that

$$\alpha_2 < d[u_0, \tilde{0}] < \alpha_1$$

implies that

$$\epsilon_2 < d[u(t), \tilde{0}] < \epsilon_1 \text{ for } T_1 + t_0 \leq t \leq T_2 + t_0.$$

If T_1, T_2 in Definition 3 are independent of t_0 , then the trivial solution of the system (3) is said to be uniformly strictly attractive.

Definition 4: The trivial solution of the system (3) is said to be strictly asymptotically stable if Definition 3 holds and the trivial solution is stable.

The trivial solution of the system (3) is said to be strictly uniformly asymptotically stable if trivial solution of the system (3) is uniformly strictly attractive and the trivial solution of the system (3) is strictly uniformly stable.

Definition 5: A function $\phi(r)$ is said to belong to the class \mathcal{K} if $\phi \in C[(0, \rho), R_+]$, $\phi(0) = 0$, and $\phi(r)$ is strictly monotone increasing in r . It is said to belong to class \mathcal{K}_∞ if $\rho = \infty$ and $\phi(r) \rightarrow \infty$ as $r \rightarrow \infty$.

Definition 6: For a real-valued function $V(t, u) \in C[R_+ \times E^n, R_+]$ we define the Dini derivatives as follows

$$D^+V(t, u) \equiv \limsup_{h \rightarrow 0^+} \frac{1}{h} [V(t+h, u+hf(t, u)) - V(t, u)]$$

$$D_-V(t, u) \equiv \liminf_{h \rightarrow 0^-} \frac{1}{h} [V(t+h, u+hf(t, u)) - V(t, u)]$$

for $(t, x) \in R_+ \times E^n$. In our earlier some studies and in the study of some others [39], [41]-[43], the differences between

the classical notion of stability and initial time difference stability did not allow the use of the behavior of the null solution in our initial time difference stability analysis. The main result presented in this section resolves those difficulties with a new approach that allows the use of the stability of the null solution of the comparison system to predict the stability properties of $v(t, \tau_0, v_0)$ the solution of (3) with respect to $\tilde{u}(t) = u(t - \eta, t_0, u_0)$ where $u(t, t_0, u_0)$ is any solution of the system (3).

Theorem 1: Assume that $f, F \in C[J \times E^n, E^n]$ and

$$(i) \lim_{h \rightarrow 0^+} \sup \frac{1}{h} [d[v - \tilde{u} + h(F(t, v) - \tilde{f}(t, \tilde{u})), \tilde{0}] - d[v - \tilde{u}, \tilde{0}]] \leq G(t, d[v - \tilde{u}, \tilde{0}])$$

for $t \in J$ and $\tilde{u}, v \in E^n$ where $G \in C[J \times R_+, R]$;

(ii) $r(t) = r(t, \tau_0, w_{10})$ is the maximal solution of the scalar differential equation exists on J

$$u' = G(t, u), u(\tau_0) = u_0 \geq 0 \text{ for } t \geq \tau_0 \text{ in } J. \quad (5)$$

Then, if $u(t)$ and $v(t)$ are any solutions of (3) and (4) through (t_0, u_0) and (τ_0, v_0) respectively on J , then we have

$$d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}] \leq r(t, \tau_0, w_{10}), t \in J \quad (6)$$

provided that $d[v_0 - u_0, \tilde{0}] \leq w_{10}$. For details [16], [23]

Corollary 1: Assume that $f, F \in C[J \times E^n, E^n]$ and either

(i)

$$d[F(t, v) - \tilde{f}(t, \tilde{u}), \tilde{0}] \leq G(t, d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), v_0 - u_0])$$

or

$$(ii) \lim_{h \rightarrow 0^+} \sup \frac{1}{h} [d[v - \tilde{u} + h(F(t, v) - \tilde{f}(t, \tilde{u})), \tilde{0}] - d[v - \tilde{u}, \tilde{0}]] \leq G(t, d[v - \tilde{u}, \tilde{0}]) \text{ where } G \in C[J \times R_+, R].$$

Then, if $d[v_0 - u_0, \tilde{0}] \leq w_{10}$, we have

$$d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}] \leq r(t, \tau_0, w_{10}) \text{ for } t \in J \quad (7)$$

where $r(t) = r(t, \tau_0, w_{10})$ is the maximal solution of the Fuzzy scalar differential equation

$$w'_1 = G(t, w_1), w_1(\tau_0) = w_{10} \geq 0 \text{ for } t \geq \tau_0, t \in J \quad (8)$$

For details [16], [23]

IV. STRICT STABILITY CRITERIA OF THE FUZZY DIFFERENTIAL SYSTEMS AND A COMPARISON RESULT IN STRICT STABILITY OF FUZZY DIFFERENTIAL EQUATIONS

Now, we are in a position to prove main result for the strict stability criteria of the Fuzzy differential systems.

Before we prove the general result in terms of the comparison principle. Let us consider the uncoupled comparison differential systems:

$$\begin{cases} (i) w'_1 = g_1(t, w_1), w_1(\tau_0) = w_{10} \geq 0 \\ (ii) w'_2 = g_2(t, w_2), w_2(\tau_0) = w_{20} \geq 0 \end{cases} \text{ where } g_1, g_2 \in C[\mathbb{R}_+^2, \mathbb{R}] \quad (9)$$

The comparison system (9) is said to be strictly stable:

If given any $\epsilon_1 > 0$ and $t \geq \tau_0, \tau_0 \in \mathbb{R}_+$, there exist a $\delta_1 > 0$ such that

$$w_{10} \leq \delta_1 \text{ implies } w_1(t) < \epsilon_1 \text{ for } t \geq \tau_0$$

and for every $\delta_2 < \delta_1$ there exists an $\epsilon_2 > 0, 0 < \epsilon_2 < \delta_2$ such that

$$w_{20} \geq \delta_2 \text{ implies } w_2(t) > \epsilon_2 \text{ for } t \geq \tau_0.$$

Here, $w_1(t)$ and $w_2(t)$ are any solutions of (i) in (9) and (ii) in (9); respectively.

The comparison system (9) is said to be strictly attractive:

If given any $\alpha_1 > 0, \gamma_1 > 0, \epsilon_1 > 0$ and $\tau_0 \in \mathbb{R}_+$, for every $\alpha_2 < \alpha_1$, there exist $\epsilon_2 < \epsilon_1, T_1 = T_1(\epsilon_1, \tau_0) > 0$ and $T_2 = T_2(\epsilon_1, \tau_0) > 0$ such that

$$w_1(t, \tau_0, w_{10}) < \epsilon_1 \text{ for } T_1 + \tau_0 \leq t \leq T_2 + \tau_0 \text{ whenever } w_{10} \leq \alpha_1$$

and

$$w_2(t, \tau_0, w_{20}) > \epsilon_2 \text{ for } T_2 + \tau_0 \geq t \geq T_1 + \tau_0 \text{ whenever } w_{20} \geq \alpha_2.$$

If T_1 and T_2 are independent of τ_0 , then the comparison system (9) is initial time difference uniformly strictly attractive for $t \geq \tau_0$.

Following main result based on this definition that result is formulated in terms of comparison principle.

In this section, we have a comparison theorem in strict stability of fuzzy differential systems via scalar differential equation and we give the proof of this theorem.

Theorem 2 Assume that

(S₁) Let $V_\mu \in C[\mathbb{R}_+ \times S_\rho, \mathbb{R}_+]$, $|V(t, u) - V(t, v)| \leq Ld[u, v], L > 0$ and for $(t, u) \in \mathbb{R}_+ \times S_\rho$, where $S_\rho = \{u \in E^n : d[u(t), \hat{0}] < \rho\}$, for each $\mu, 0 < \mu < \rho$, and V_μ is locally Lipschitzian in z and for $(t, z) \in \mathbb{R}_+ \times S_\rho$ and $d[z, \hat{0}] \geq \mu$, such that

$$b_1(d[z, \hat{0}]) \leq V_\mu(t, z) \leq a_1(d[z, \hat{0}]), a_1, b_1 \in \mathcal{K}$$

$$D^+V_\mu(t, v - \tilde{u})$$

$$= \limsup_{h \rightarrow 0^+} \frac{1}{h} [V_\mu(t+h, v - \tilde{u} + h(F(t, v) - \tilde{f}(t, \tilde{u}))) - V_\mu(t, v - \tilde{u})]$$

$$\leq g_1(t, V_\mu(t, z));$$

(10)

(S₂) for each $\theta, 0 < \theta < \rho, V_\theta \in C[\mathbb{R}_+ \times S_\rho, \mathbb{R}_+]$ and V_θ is locally Lipschitzian in z and for $(t, z) \in \mathbb{R}_+ \times S_\rho$ and $d[z, \hat{0}] \leq \theta$,

$$b_2(d[z, \hat{0}]) \leq V_\theta(t, z) \leq a_2(d[z, \hat{0}]), a_2, b_2 \in \mathcal{K},$$

$$D^+V_\theta(t, v - \tilde{u})$$

$$= \limsup_{h \rightarrow 0^+} \frac{1}{h} [V_\theta(t+h, v - \tilde{u} + h(F(t, v) - \tilde{f}(t, \tilde{u}))) - V_\theta(t, v - \tilde{u})]$$

$$\geq g_2(t, V_\theta(t, z));$$

(11)

where $g_2(t, w) \leq g_1(t, w), g_1, g_2 \in C[\mathbb{R}_+^2, \mathbb{R}], g_1(t, 0) = g_2(t, 0) = 0$ and $z(t) = v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0)$ for $t \geq \tau_0, v(t, \tau_0, v_0)$ of the system (4) through (τ_0, v_0) and

$u(t - \eta, t_0, u_0)$, where $u(t, t_0, u_0)$ is any solution of the system (3) for $t \geq \tau_0 \geq 0, t_0 \in \mathbb{R}_+$ and $\eta = \tau_0 - t_0$.

We have the strict stability properties of the comparison systems give us the corresponding strict stability concept of the fuzzy differential equation solution $v(t, \tau_0, v_0)$ of the system (4) through (τ_0, v_0) with respect to the solution $u(t - \eta, t_0, u_0)$ of the system (3) with initial time difference where $u(t, t_0, u_0)$ is any solution of the system (3) for $t \geq \tau_0 \geq 0, t_0 \in \mathbb{R}_+$.

Proof of Theorem 2: We will only prove the case of strict uniformly asymptotically stability. Suppose that the comparison differential systems in (9) is strictly uniformly asymptotically stable, then for any given $\epsilon_1, 0 < \epsilon_1 < \delta$, there exist a $\delta^* > 0$ such that $w_{10} \leq \delta^*$ means that $w_1(t, \tau_0, w_{10}) < b_1(\epsilon_1)$ for $t \geq \tau_0$.

For this ϵ_1 we choose δ_1 and δ_0 , such that $a_1(\delta_1^*) \leq \delta^*$ and $\delta_1^* < \epsilon_1$ where $\delta_1^* = \max\{\delta_1, \delta_0\}$, then we claim that

$$d[v_0 - u_0, \tilde{0}] < \delta_1, d[\tau_0 - t_0, \tilde{0}] < \delta_0 \text{ means that}$$

$$d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}] < \epsilon_1 \text{ for } t \geq \tau_0.$$

If it is not true, then there exist t_1 and $t_2, t_2 > t_1 > \tau_0$ and a solution $q(t)$ of

$$q' = \tilde{f}(t, q), q(\tau_0) = v_0 - u_0 \text{ for } t \geq \tau_0$$

$$\text{with } d[\tau_0 - t_0, \tilde{0}] < \delta_{11} \text{ and } d[v_0 - u_0, \tilde{0}] < \delta_1$$

$$d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}] < \delta_1^*,$$

$$d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}] = \epsilon_1 \text{ and}$$

$$\delta_1^* \leq d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}] < \epsilon_1 \text{ for } [t_1, t_2].$$

If we choose $\mu = \delta_1^*$ and using the theory of differential inequalities, we get

$$b_1(\epsilon_1) = b_1(d[v(t_2, \tau_0, v_0) - u(t_2 - \eta, t_0, u_0), \tilde{0}])$$

$$\leq V_\mu(t_2, v(t_2, \tau_0, v_0) - u(t_2 - \eta, t_0, u_0))$$

$$\leq r(t_2, t_1, V_\mu(t_1, v(t_1, \tau_0, v_0) - u(t_1 - \eta, t_0, u_0)))$$

$$\leq r(t_2, t_1, a_1(\delta_1^*))$$

$$\leq r(t_2, t_1, \delta^*)$$

$$< b_1(\epsilon_1).$$

and then we have a contradiction here that this inequality is impossible $b_1(\epsilon_1) < b_1(\epsilon_1)$, and this inequality that we obtain is a contradiction. Here the function $r(t, \tau_0, w_{10})$ is the maximal solution of the comparison system. We have that the strict stability of the fuzzy differential equation is true and we have uniformly stability with initial time difference. Then, we have proved that the fuzzy differential equation is strictly uniformly attractive with initial time difference.

For any given $\delta_2, \epsilon_2 > 0, \delta_2 < \delta^*$, we choose $\bar{\delta}_2$ and $\bar{\epsilon}_2$ such that $a_1(\delta_2) < \bar{\delta}_2$ and $b_1(\epsilon_2) \geq \bar{\epsilon}_2$. For these $\bar{\delta}_2$ and $\bar{\epsilon}_2$, since (9) is strictly uniformly attractive, for any $\bar{\delta}_3 < \bar{\delta}_2$ there exist $\bar{\epsilon}_3$ and T_1 and T_2 and we assume that the inequality $T_2 < T_1$ such that $\bar{\delta}_3 < w_{10} = w_{20} < \bar{\delta}_2$ implies

$$\begin{aligned} r(t, \tau_0, w_{10}) &\leq r(t, \tau_0, \bar{\delta}_2) < \bar{\epsilon}_2 \\ \rho(t, \tau_0, w_{20}) &\geq \rho(t, \tau_0, \bar{\delta}_3) > \bar{\epsilon}_2 \end{aligned}$$

where $r(t, \tau_0, w_{10})$ and $\rho(t, \tau_0, w_{20})$ is the maximal solution and minimal solution of the comparison system (9) part (i) and the comparison system (9) part (ii); respectively.

Now, for any δ_3 , let $b_2(\delta_3) \geq \bar{\delta}_3$. We choose ϵ_3 such that $a_2(\epsilon_3) < \bar{\epsilon}_3$. Then by using comparison system (9), (i) and (A₁), we have

$$\begin{aligned} b_1(d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}]) \\ \leq V_\mu(t, v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0)) \\ \leq r(t, \tau_0, V_\mu(\tau_0, v_0 - u_0)) \\ \leq r(t, \tau_0, a_1(d[v_0 - u_0, \tilde{0}])) \\ \leq r(t, \tau_0, \bar{\delta}_2) \\ < \bar{\epsilon}_2 \leq b_1(\epsilon_2) \end{aligned}$$

So that we obtain that we have the inequality

$$b_1(d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}]) < b_1(\epsilon_2)$$

if we have b_1^{-1} and it exists which implies that $d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}] < \epsilon_2$ for $t \in [\tau_0 + T_2, \tau_0 + T_1]$.

We have used the comparison principle and (ii) and (S₂) and then we have

$$\begin{aligned} a_2(d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}]) \\ \geq V_\theta(t, v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0)) \\ \geq \rho(t, \tau_0, V_\theta(\tau_0, v_0 - u_0)) \\ \geq \rho(t, \tau_0, b_2(\delta_3)) \\ \geq \rho(t, \tau_0, \bar{\delta}_3) \\ > \bar{\epsilon}_3 \geq a_2(\epsilon_3) \end{aligned}$$

Then we have an important inequality that we obtain the inequality

$$a_2(d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}]) > a_2(\epsilon_3)$$

if we have that a_2^{-1} exists which implies that for $d[v(t, \tau_0, v_0) - u(t - \eta, t_0, u_0), \tilde{0}] > \epsilon_3$ for $t \in [\tau_0 + T_2, \tau_0 + T_1]$, and we have $v(t, \tau_0, v_0)$ solution of (4) through (τ_0, v_0) is strictly uniformly attractive with respect to the

solution $u(t - \eta, t_0, u_0)$ is solution of the system (3) for $t \geq \tau_0 \geq 0, t_0 \in \mathbb{R}_+$. So that we have completed the proof of the comparison theorem of fuzzy differential equations with strict stability of fuzzy differential equations with Lyapunov functions.

This completes the proof of the Theorem 2.

V. CONCLUSION

In this study, we present some results for the study of stability of fuzzy differential equations and the strict stability of fuzzy systems. First of all, we present the fundamental concept of fuzzy calculus and differential equations in fuzzy calculus. Firstly, we consider and define the strict stability of fuzzy differential equations by Lyapunov functions. Secondly, we develop a new comparison principle for the strict stability of fuzzy differential systems, then we prove the strict stability criteria with Lyapunov functions. It may provide a greater prospect of solving problems which exhibit strict stability of fuzzy differential equations by Lyapunov functions.

ACKNOWLEDGMENT

This work has been supported by The Yıldız Technical University, Gebze Technical University and Yeditepe University Department of Mathematics and The Scientific and Technological Research Council of Turkey. The authors also would like to thank the referees for their insightful comments and detailed suggestions which improved the quality of the paper.

REFERENCES

- [1] Aumann, R.J. Integrals of set-valued functions, *J. Math. Anal. Appl.* 12 (1965) 1-12.
- [2] Bernfeld, S. and Lakshmikantham, V. *An Introduction to Nonlinear Boundary Value Problems*. Academic Press, New York, 1974.
- [3] Bede, B. and Gal, S. G. *Generalizations of the differentiability of fuzzy-number valued functions with applications to fuzzy differential equations*, *Fuzzy Sets and Systems*, Vol. 151, No. 3, (2005) 581-589.
- [4] Bede, B., Rudas, I. J. and Bencsik, A. L. *First order linear fuzzy differential equations under generalized differentiability*, *Information Sciences*, vol. 177, no. 7, (2007) pp. 1648-1662.
- [5] Bede, B., Stefanini, L. *Generalized differentiability of fuzzy-valued functions*, *Fuzzy Sets and Systems*, Vol. 230, November, (2013) pp. 119-141.
- [6] Buckley, J.J. and Feuring, T.H. *Fuzzy differential equations*, *Fuzzy Sets and Systems*, Vol. 110, (2000) pp. 43-54.
- [7] Gomes, L. T., Barros, L. C., Bede, B. *Fuzzy Differential Equations in Various Approaches*, Springer International Publishing (2015).
- [8] Buckley, J. J. and Feuring, T. H. *Fuzzy differential equations*, *Fuzzy Sets and Systems*, 110 (2000), 43 -54.
- [9] Chen, Z., Fu, X., The variational Lyapunov function and strict stability theory for differential systems, *Nonlinear Analysis* 64, 1931 - 1938, (2006).
- [10] Ding, Z., Ming, M. and Kandel, A. Existence of solutions of Fuzzy differential equations, *Inform. Sci.*, 99 (1997), 205 - 217.
- [11] Dubois, D. and Prade, H. Towards fuzzy differential calculus, Part I, Part II, Part III, *Fuzzy Sets and Systems*, 8 (1982), 1 - 17, 105 - 116, 225 - 234.
- [12] Kaleva, O. Fuzzy differential equations. *Fuzzy Sets and Systems* 24 (1987) 301-317.
- [13] Kaleva, O. On the calculus of fuzzy valued mappings, *Appl. Math. Lett.*, 3 (1990), 55 - 59.
- [14] Kaleva, O. The Cauchy problem for fuzzy differential equations. *Fuzzy Sets and Systems* 35 (1990) 389-396.
- [15] Lakshmikantham, V. and Leela, S. *Differential and Integral Inequalities*, Vol. I. Academic Press, New York, 1969.

- [16] Lakshmikantham, V. and Leela, S. Fuzzy differential systems and the new concept of stability. *Nonlinear Dynamics and Systems Theory*, 1 (2) (2001), 111-119
- [17] Lakshmikantham, V. and Leela, S. A new concept unifying Lyapunov and orbital stabilities. *Communications in Applied Analysis*, (2002), 6 (2).
- [18] Lakshmikantham, V. and Leela, S. Stability theory of fuzzy differential equations via differential inequalities. *Math. Inequalities and Appl.* 2 (1999) 551-559.
- [19] Lakshmikantham, V., Leela, S. and Martynyuk, A.A. *Practical Stability of Nonlinear System*. World Scientific Publishing, NJ, 1990.
- [20] Lakshmikantham, V., Leela, S. and Martynyuk, A. A. *Stability Analysis of Nonlinear System*. Marcel Dekker, New York, 1989.
- [21] Lakshmikantham, V. and Mohapatra, R. Basic properties of solutions of fuzzy differential equations. *Nonlinear Studies* 8 (2001) 113-124.
- [22] Lakshmikantham, V. and Mohapatra, R. N. : Strict Stability of Differential Equations, *Nonlinear Analysis*, Volume 46, Issue 7, Pages 915-921 (2001)
- [23] Lakshmikantham, V. and Mohapatra, R. N. *Theory of Fuzzy Differential Equations*. Taylor and Francis Inc. New York, 2003.
- [24] Lakshmikantham, V. and Vatsala, A.S., Differential inequalities with time difference and application, *Journal of Inequalities and Applications* 3, (1999) 233-244.
- [25] Li, A., Feng, E. and Li, S., Stability and boundedness criteria for nonlinear differential systems relative to initial time difference and applications. *Nonlinear Analysis: Real World Applications* 10(2009) 1073-1080.
- [26] Liu, K., Yang, G., Strict Stability Criteria for Impulsive Functional Differential Systems, *Journal of Inequalities and Applications* , 2008:243863 doi:10.1155/2008/243863, (2008).
- [27] Lyapunov, A. Sur les fonctions-vecteurs complement additives. *Bull. Acad. Sci. URSS, Ser. Math* 4 (1940) 465-478.
- [28] Massera, J. L. The meaning of stability. *Bol. Fac. Ing. Montevideo* 8 (1964) 405-429.
- [29] Nieto, J. J. The Cauchy problem for fuzzy differential equations. *Fuzzy Sets and Systems*, (102 (1999), 259 - 262.
- [30] Park, J. Y. and Hyo, K. H. Existence and uniqueness theorem for a solution of Fuzzy differential equations, *Inter. J. Math. and Math. Sci.* 22 (1999), 271-279.
- [31] Puri, M. L. D. and Ralescu, A. Differential of Fuzzy functions, *J. Math. Anal. Appl.* 91 (1983), 552 - 558.
- [32] Rojas K., Gomez D., Monteroa J., Rodriguez J. Tinguaro, Strictly stable families of aggregation operators, *Fuzzy Sets and Systems*, 228, 44-63 (2013).
- [33] Rojas K., Gomez D., Monteroa J., Rodriguez J. Tinguaro, Valdivia A., Paiva F., Development of child's home environment indexes based on consistent families of aggregation operators with prioritized hierarchical information, *Fuzzy Sets and Systems* 241, 41-60, (2014).
- [34] Shaw, M. D. and Yakar, C., Generalized variation of parameters with initial time difference and a comparison result in term Lyapunov-like functions, *International Journal of Non-linear Differential Equations-Theory-Methods and Applications* 5, (1999). 86-108.
- [35] Shaw, M. D. and Yakar, C., Stability criteria and slowly growing motions with initial time difference, *Problems of Nonlinear Analysis in Engineering Systems* 1, (2000) 50-66.
- [36] Song, S. J., Guo, L. and Feng, C. H. Global existence of solutions of Fuzzy differential equations, *Fuzzy Sets and Systems*, 115 (2000), 371 - 376.
- [37] Song, S.J. and Wu, C. Existence and Uniqueness of solutions to Cauchy problem of Fuzzy differential equations, *Fuzzy Sets and Systems*, 110 (2000), 55 - 67.
- [38] Yakar, C. Boundedness criteria with initial time difference in terms of two measures, *Dynamics of Continuous, Discrete & Impulsive Systems*. Series A, vol. 14, supplement 2, (2007) 270-274, .
- [39] Yakar C. and Çiçek M. and Gücen, M.B. "Boundedness and Lagrange stability of fractional order perturbed system related to unperturbed systems with initial time difference in Caputo's sense." *Advances in difference Equations* (2011):54. doi:10.1186/1687-1847-2011-54. ISSN: 1687-1847.
- [40] Yakar C., Çiçek M. and Gücen M. B. "Practical Stability in Terms of Two Measures for Fractional Order Dynamic Systems in Caputo's Sense with Initial Time Difference" *Journal of the Franklin Institute*. PII: S0016-0032(13)00377-3 DOI: <http://dx.doi.org/10.1016/j.jfranklin.2013.10.009>. Ref.: FI1903. (2013). (SCI).
- [41] Yakar C., Çiçek M. and Gücen M. B. "Practical Stability, Boundedness Criteria and Lagrange Stability of Fuzzy Differential Systems" *Journal of Computers and Mathematics with Applications*. 64 (2012) 2118-2127. Doi: 10.1016/j.camwa.2012.04.008. (2012).
- [42] Yakar, C. Strict stability criteria of perturbed systems with respect to unperturbed systems in terms of initial time difference. *Complex Analysis and Potential Theory*, World Scientific, Hackensack, NJ, USA (2007) 239-248.
- [43] Yakar, C. and Shaw, M. D. A comparison result and Lyapunov stability criteria with initial time difference. *Dynamics of Continuous, Discrete & Impulsive Systems*. Series A, vol. 12, no. 6, (2005) 731-737.
- [44] Yakar, C. and Shaw, M. D. Initial time difference stability in terms of two measures and a variational comparison result. *Dynamics of Continuous, Discrete & Impulsive Systems*. Series A, vol. 15, no. 3, (2008) 417-425, .
- [45] Yakar, C. and Shaw, M. D. Practical stability in terms of two measures with initial time difference. *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 12, (2009) e781-e785.
- [46] Yoshizawa, T. *Stability Theory by Lyapunov's second Method*, *The Mathematical Society of Japan*, Tokyo, 1966.
- [47] Zadeh, L. A. Fuzzy Sets, *Inform. Control.*, 8 (1965), 338 - 353.
- [48] Zhang, Y. Criteria for boundedness of Fuzzy differential equations, *Math. Ineq. Appl.*, 3 (2000), 399 -410.
- [49] Zhang, Y., Sun J., Strict stability of impulsive functional differential equations, *J. Math. Anal. Appl.*, 301, 237-248 (2005).