Synthesis and Reactions of Sulphone Hydrazides

Mohamed E. Khalifa

Abstract—The chemistry of sulphone hydrazide has gained increase interest in both synthetic organic chemistry and biological fields and has considerable value. The therapeutic importance of these compounds is the attractive force to continue research in such a point. The present review covers the literature up to date for the synthesis, reactions and applications of such compounds.

Keywords—Sulphone hydrazide compounds, Reactions, Synthesis, Biological activities.

I. INTRODUCTION

CULPHONE hydrazides exhibit a great interest due to their Dimportance in synthesis a variety of valuable heterocyclic compounds which have pharmaceutical activities and chemotherapeutic value [1]-[4] against cognitive disorder, Alzheimer's disease [5], [6], hepatitis C virus [7]–[9], βlactamase inhibitory properties against class A and C enzymes, antitumor [10], tuberculosis [11]-[16], antimycobacterial agents [14] and bacteriostatic activity with respect to the Koch bacillus [17]. They also produce ringclosed hydrazide-imide copolymers with novel properties used in manufacturing separating membranes [18], cross-linked tough foam copolymer which have good resistance to heat and chemicals [19]-[21] besides their industrial uses in various fields such as rubber [22], epoxy resins [23], leather [24], flame-resistant fibers [25]-[27], color developer [28], [29], stain bleaching and/or anti dye-transfer [30], decomposable blowing agents [31]-[36], flexible self-expandable and selfcontained unit with pyrotechnic sheet [37] and their utility analytical field as a sensitive fluorescent labeling reagent for determination of aromatic and aliphatic aldehydes e.g. 4-(5,6dimethoxy-2-phthalimidinyl)phenyl sulfonyl hydrazide (DPSH) [38].

II. SYNTHESIS

The main objective of this section is to provide a comprehensive account for the synthesis of various sulphone hydrazide compounds. Usually, they are prepared by the reaction of sulphonyl chloride and hydrazines [39] or reduction of azo compounds [40]. However, these reactions were carried out in organic solvent such as pyridine and DMF [41]–[43].

A. Diels-Alder Reactions with Cyclic Sulfones Reactions of spiro[1-benzothiophene-4,5'-[1,3]dioxane]-

Mohamed E. Khalifa is with the Faculty of Science, department of Chemistry, Taif University, Taif 21974, Saudi Arabia and with Higher Institute of Engineering and Technology, department of Chemical Engineering, New Damietta, Egypt (phone: +966569846966, e-mail: mohamedezzat200@hotmail.com)

4',6'-dione **1** with various amines led to the formation of 4carbamoylhexahydrobenzo[b]thiophene-4-carboxylic acid 1,1dioxides **2** or their decarboxylation products, depending on the conditions. Hydrazinolysis of the spiro adducts in DMF gave the corresponding monohydrazides **3** [44].



Fig. 1 Diels Alder reaction with cyclic sulfones

B. From Hydrazines

Reaction of triflouromethanesulfonic anhydride and triflouromethanesulfonyl chloride with hydrazine, phenyl hydrazine and 1,1-dimethylhydrazine at low temperature gave the corresponding triflouromethanesulfonic hydrazides **4** [45].

$$2CF_3SO_2Cl + 3N_2H_4 \longrightarrow [CF_3SO_2NH.NH]_2$$

Fig. 2 Synthesis of triflouromethanesulfonic hydrazides 4

Treating of p-MeC₆H₄SO₂Na with alkyl bromide in EtOH to yield p-MeC₆H₄SO₂R **5** [R, Me, Et, Pr, Bu, iso-Amyl, benzyl, which on heating with KMnO₄, and water to yield p-HO₂C.C₆H₄SO₂R **6**. The mixture of **6** in 20 cc. EtOH and 0.012 mol 60% N₂H₄.H₂O was heated on a water bath for 1 h., and the residue recrystalized from EtOH to give a quantitative yield of 4-(alkylsulfonyl)benzohydrazide **7** having the same antituberculous activity as isonicotinyl hydrazide [46].



Fig. 3 Synthesis of 4-(alkylsulfonyl)benzohydrazide 7

Treatment of hydrazine derivatives with sulfonyl chlorides in the presence of basic alumina, in a mortar with grinding by a pestle, afforded the corresponding sulphone hydrazide derivatives 8.

$$R-SO_2-Cl + H_2NR' \xrightarrow{\text{Basic Alumina}} R-SO_2NHR'$$

Fig. 4 Synthesis of sulphone hydrazide derivatives 8

On the other hand, t-butyl carbazate can be used instead of hydrazine that reacted with sulphonyl chloride in the presence of basic alumina to produce intermediate 9. Hydrolysis of the intermediate carbazate 9 gave directly in situ the sulphone hydrazide derivatives 10 under mild conditions [47].



Fig. 5 Synthesis of sulphone hydrazide derivatives 10

C. By Microwave Irradiation

The chemical application of microwave irradiation has now become an area of interest for the synthesis of a wide variety of compounds and efficient functional group transformations under solvent-free conditions [48]–[53], where this method was initially introduced in 1986 [54]. The advantages of microwave-expedited chemical transformations are cleaner reactions, higher efficiency and selectivity, shorter reaction times, and the ease of manipulation.

A synthesis of 1-aryl-2-*p*-toluenesulphonyl hydrazides **12** under solvent-free conditions involved mixing the *p*-toluenesulphonyl chloride **11** and aryl hydrazine (1:2 ratio) followed by exposure to microwave irradiation (600W) under solvent-free conditions, where the reaction carefully monitored (by TLC) to regulate the ratio of the substrates, irradiation time and power level of the microwave oven to achieve the maximum yield [55].



Fig. 6 Synthesis of 1-aryl-2-p-toluenesulphonyl hydrazides 12

III. REACTIONS

A. Oxidation

Oxidation of p-RC₆H₄SO₂NHNH₂ (**13**; R = H, Me, MeO, Cl, Br, AcNH) in AcOH with KMnO4 and in several drops of 3% aq. H₂O₂ gave the symmetric diaryl disulphide S,S,S',S'-tetraoxides **14** [56].



Fig. 7 Oxidation using acidic potassium permanganate

The oxidation of sulphonhydrazides with benzeneselenic acid (PhSeO₂H) produce the corresponding selenosulphonate derivatives, where phenyl areneselenosulphonates (**15**, R = aryl) were obtained. The method is also applicable to alkaneselenosulphonates, as illustrated by the methane derivative (**15**, R = Me) [57].

$$\frac{\text{Oxid.}}{\text{R'SO}_2\text{NHNH}_2 + \text{PhSeO}_2\text{H}} \xrightarrow{\text{Oxid.}} \frac{\text{R'SO}_2 \text{ SePh} + \text{N}_2 + \text{H}_2\text{O}}{15}$$

Fig. 8 Oxidation of sulphonhydrazides with benzeneselenic acid

Phenyl areneselenosulphonates were reacted to the acetylinic derivatives to yield the adducts **16** [58]



Fig. 9 Reaction with acetylinic derivatives

B. Nucleophilic Reaction

Nucleophilic displacement reactions of p-MeC₆H₄SO₂NHNH₂ on RX (X = Br, Cl; R = aryl, alkyl) gave sulphones p-MeC₆H₄SO₂R **17** in 80-95% yields [59].



Fig. 10 Reaction with alkyl halides

C. Reaction with Thiophenecarboxaldehydes

4,4'-Sulphonyl bis(benzohydrazide) was treated with thiophenecarboxaldehydes to yield condensation products **18** (R = H, Cl, Br) and **19**, which showed bactericidal activity [60]. Also, methionine sulphone hydrazide was coupled to 6-aldehydosugars and the reaction was catalytically enhanced by Mn^{2+} under physiological condition. The reaction was applied to label surface glycoproteins of erythrocytes with I-35S after treating cells with galactose oxidase [61].



Fig. 11 Reaction with thiophenecarboxaldehydes

D. With Cyanogen Bromide and/or Acetyl Acetone

The synthetic potential of sulphanyl/-sulphonyl acetic acid hydrazide derivatives **20** has been utilized in two ways. In one case cyanogen bromide has been used as a source of single carbon and carbohydrazide acts as a source of four atoms reacting through its enol form generating the amino oxadiazole derivatives **21**. In the second case, double electrophilic character of acetyl acetone and the enhanced nucleophilicity of ¹N-nitrogen of the carbohydrazide have been used to synthesize 2,5-dimethyl pyrroles **22**. The synthesized compounds exhibited antimicrobial activity against gram -ve bacterium *Escherichia coli*, gram +ve bacterium *Bacillus staphylococci* and fungi Penicillium and Aspergillus [62].



Fig. 12 Reaction with cyanogen bromide and/or acetyl acetone

E. With Acetylenic Ester

The reaction of the aryl sulphonyl hydrazide with the acetylenic ester in the presence of triphenyl phosphine led to the corresponding derivatives **23** in good yields [63].



Fig. 13 Reaction with acetylinic esters

F. With 5-Aryl-2,3-Dihydrofuran-2,3-Diones

Decyclization of 5-aryl-2,3-dihydrofuran-2,3-diones under the action of *p*-toluenesulphonyl hydrazides in anhydrous dioxane afforded β -N-(4-methylphenylsulphonyl) hydrazides of aroyl pyruvic acid **24** which exhibit antiinflammatory, and antimicrobial activities [64].



Fig. 14 Decyclization action of sulphone hydrazides

REFERENCES

- A. Mavrodin, V. Zotta, M. Vorel-Stoenescu and D. Oteleanu, "Experiments on the sulfone group. V. New derivatives of 4phenylsulphone-benzoylhydrazones," *Pharmazeutische Zentralhalle fuer Deutschland*, 1957, vol. 96, pp. 509-16.
- [2] A. Mavrodin, V. Zotta, M. Vorel-Stoenescu and D. Oteleanu, "Studies on the sulfone group. Iv. New sulfone-hydrazide derivatives," *Pharmazeutische Zentralhalle fuer Deutschland*, 1956, vol. 95, pp. 353-61.
- [3] A. Mavrodin, C. Demetrescu and C. Chirita, "Analyse du cytosquelette par des approches bioinformatiques à haut débit de génomique comparative et de transcriptomique," *Revue Roumaine de Chimie*, 1965, vol.10, no.10, pp. 1025-33.
- [4] I. Matei and M.Grigoras, "Vinyl polymer analogs with biological activity," *Rom. Buletinul Institutului Politehnic din Iasi*, 1967, vol. 13, no. 3-4, pp. 261-8.
- [6] P. Castro, L. Jose, I. Churcher, K. Dinnell, T. Harrison, S. Kerrad, A. J. Nadin, P. J. Oakley, A. P. Owens, D. E. Shaw, M. R. Teall, B. J. Williams and S. Williams, "Preparation of aryl sulfones which modulate the action of gamma secretase," PCT Int. Appl. 2002, 159 pp.
- [7] S. Venkatraman, K. X. Chen, A. Arasappan,; F. G. Njoroge, V. M. Girijavallabhan, T. Chan, B. A. McKittrick, A. J. Prongay and V. S. Madison, "Preparation of macrocyclic NS3-serine protease inhibitors of hepatitis C virus comprising alkyl and aryl alanine p2 moieties," PCT Int. Appl. 2001, 218 pp.
- [8] K. X. Chen, A. Arasappan, S. Venkatraman, T. N. Parekh, H. Gu, F. G. Njoroge, V. M. Girijavallabhan, A. Ganguly, A. Saksena, E. Jao, N. H. Yao, A. J. Prongay, V. S. Madison and B. Vibulbhan, "Preparation of macrocyclic NS3-serine protease inhibitors of hepatitis C virus comprising n-cyclic p2 moieties," PCT Int. Appl. 2001, 402 pp.
- Z. Zhu, Z. Sun, S. Venkatraman, F. G. Njoroge, A. Arasappan, B. A. Malcolm, V. M. Girijavallabhan, R. G. Lovey and K. X. Chen, "Preparation of diaryl peptides as NS3-serine protease inhibitors of hepatitis C virus," PCT Int. Appl. 2002, 149 pp.
 O. A. Fathalia and M. Rizk, "Biological activity of some new uracil
- [10] O. A. Fathalia and M. Rizk, "Biological activity of some new uracil derivatives on Ehrlich ascites cells," *Bulletin of the National Research Centre* (Egypt), 1993, vol. 18, no.1, pp. 55-64.
 [11] S. K. Gupta, I. S. Mathur and M. C. Khosla, "The therapeutic activity of
- [11] S. K. Gupta, I. S. Mathur and M. C. Khosla, "The therapeutic activity of p-aminop'-(hydrazido methylamino) diphenyl sulphone (S.N.322) and its copper chelate in experimental tuberculosis of guinea-pigs and mice," *Archives Internationales de Pharmacodynamie et de Therapie*, 1958, vol. 114, pp. 373-80.
- [12] Y. Aoki, "On the antituberculous activity of chemotherapeutic agents bearing hydrazino radical," *Nippon Saikingaku Zasshi*, 1955, vol. 10, no. 11, pp. 919-23.
- [13] Y. Aoki, "On the antituberculous activity of chemotherapeutic agents bearing hydrazino radical," *Nippon Saikingaku Zasshi*, 1955, vol. 10, no. 10, pp. 867-72.
- [14] M. N. Shchukina and T. P. Sycheva,"Antituberculous chemotherapeutic preparations," *Khim. Nauka I Prom.* 1956, vol.1, pp. 419-32.
- [15] O. A. Phillips, A. V. N. Reddy, E. L. Setti, P. Spevak, D. P. Czajkowski, H. Atwal, S. Salama, R. G. Micetich and S. N. Maiti, "Synthesis and biological evaluation of penam sulfones as inhibitors of β-lactamases," *Bioorganic & Medicinal Chemistry*, 2005, vol. 13, no.8, pp. 2847-58.
- [16] D. Birca-Galateanu, C. Chirita, C. Demetrescu and A. Mavrodin, "Synthesis and infrared spectroscopic study of the structure of some new

International Journal of Chemical, Materials and Biomolecular Sciences

ISSN: 2415-6620

Vol:7, No:6, 2013

diarylsulfonhydrazidones with antituberculous action," *Studuii Cercetari Chim.*, 1965, vol.14, no.1, pp. 85-94.

- [17] M. C. Khosla, N. Anand and M. L. Dhar, "Studies in potential antimycobacterial agents. XI. Synthesis of p-amino-p'-(carboxyalkylamino)diphenyl sulfones, their esters, hydrazides, and amides," *Journal of Scientific & Industrial Research*, 1955, vol. 14C, pp. 222-7.
- [18] V. Zota, M. Stoenescu, D. Oteleanu, E. Chilibon and A. Mavrodin, "Diphenyl sulfone hydrazides with bacteriostatic activity with respect to the Koch bacillus," *Chemia*, 1963, vol. 8, no.1, pp. 303-9.
- [19] H. Nagata and A. Takada, "Separation membranes with excellent gas separation capacity, permeability, processability, and tensile strength," *Jpn. Kokai Tokkyo Koho*, 2003, 15 pp.
- [20] (Dynamit-Nobel A.-G.),"Cross-linked polyolefin foams," 1962, 8 pp.
- [21] H. Sheng-Huei and C. Jiun-Hsiang, Journal of Polymer Science, Part A: Polymer Chemistry, 2001, vol. 39, no.13, pp. 2271-2286.
- [22] Y. A. Noura and A. M. Nadia, "Thermal degradation behaviour of novel wholly para-oriented aromatic polyamide-hydrazides containing sulfoneether linkages," *Polymer Degradation and Stability*, 2009, vol. 94, no.10, pp. 1656-1665.
- [23] S. Y. Kim, "Sidewall rubber composition of radial pneumatic tire for truck/bus," KR Patent 2002089891, 2002.
- [24] T. Kousaka, M. Iwata and T. Ito, "Epoxy resin compositions with mechanical strength and tackiness stability for fiber-reinforced composites for prepregs for honeycomb panels," PCT Int. Appl., 2007, 24 pp.
- [25] A. Jakus, S. Sulc and P. Beno, "Leather substitutes for footwear linings," *Czech.*, 1982, 6 pp.
- [26] I. Tanaka, K. Kato, K. Watanabe, K. Kurita and M. Sugino, "Aromatic polyamide-hydrazide copolymers," *Jpn. Kokai Tokkyo Koho*, 1976, 5 pp.
- [27] I. Tanaka, K. Kato, K. Watanabe, K. Kurita and M. Sugino, "Fireresistant polyester-hydrazide copolymers," *Jpn. Kokai Tokkyo Koho*, 1976, 10 pp
- [28] I. Tanaka, K. Kato, K. Watanabe, "Studies on flame-resistant fibers. IV. Light-resistance and flammability of various aromatic polymers," *Kobunshi Ronbunshu*, 1979, vol. 36, no.11, pp. 713-19.
- [29] K. Okauchi, "Silver halide photographic material and its processing," Jpn. Kokai Tokkyo Koho, 1995, 26 pp.
- [30] D. Clarke, J. D. Goddard, P. L. R. Stanley, N. E. Milner, "Silver halide color photographic material," Eur. Pat. Appl., 1993, 18 pp.
- [31] D. Convents, M. Doornink, R. G. Smith and D. W. Thornthwaite, "Oxidation process and composition using hydrazides," U.S. Pat. Appl. Publ., 2002, 10 pp.
- [32] T. Fujiwara, S. Nagashitani, H. Motoki, T. Ouchi and K. Miyamoto, " Compositions for weather-resistant fireproof composite coatings and their formation," *Jpn. Kokai Tokkyo Koho*, 1997, 10 pp.
- [33] J. Gagliani, U. A. K. Sorathia and R. Lee, "Preparing polyimide foams with blowing agents," Eur. Pat. Appl., 1984, 17 pp.
- [34] J. Ahnemiller, "Process and performance optimization of chemically foamed polyesters," *Plastics Engineering*, 1983, vol. 39, no. 2, pp. 21-3.
- [35] Ontario Research Foundation. Brit. "Storage-stable, solid, foamable epoxy-amine composition," U.S. Pat. Appl. Publ., 1968, 4 pp.
- [36] B. N. Dinzburg and B. A. Safrai, "Sole rubbers with different blowing agents," *Plenochnykh Materialov I Iskusstv. Kozhi*, 1964, vol. 15, pp. 30-8.
- [37] T. Francis, J. F. Furrer, M. H. Jones, G. E. Murray and M. P. Thorne, "Flexible self-expandable and self-contained unit with pyrotechnic sheet," U.S. Pat. Appl. Publ., 1966, 6 pp.
- [38] H. Inouej, J. Yamazaki, E. Sugino, H. Fujioka, k. Kohashi, Y. Date, Y. Ishii and Y. Tsuruta, "4-(5,6-Dimethoxy-2-phthalimidinyl)phenyl hydrazide as a fluorescent labeling reagent for determination of aldehydes in hplc," *Analytical sciences*, 1997, vol. 13, no. 4, pp. 669-671.
- [39] C.S. Rooney, E.J. Cragoe, C.C. Porter and J.M. Sprague, "Synthesis of sulfonylhydrazine derivatives with monoamine oxidase inhibitory activity," *J. Med. Pharm. Chem*, 1962, vol. 52, pp. 155-67.
- [40] Reinhard, W.H. "Fragmentierung von Azoverbindungen, III. Protonierung und Halogenid-Abspaltung der 2-Halogen-phenyl-Anionen in Äthanol," *Chem. Ber.*, 1965, vol. 98, no. 1, pp. 222-34.

- [41] R.T. Hrubiec, K. L. Shyam, A. Cosby, R. Furubayashi and A.C. Sartorelli, "Synthesis and evaluation of 2-substituted 1-methyl-1-(4tolylsulfonyl)hydrazines as antineoplastic agents," *J. Med. Chem.*, 1986, vol. 29, no.7, pp. 1299-1301.
- [42] K. L. Shyam, R. Furubayashi, R.T. Hrubiec, A. Cosby and A.C. Sartorelli, "1,2-Bis(arylsulfonyl)hydrazines. 2. The influence of arylsulfonyl and aralkylsulfonyl substituents on antitumor and alkylating activity," *J. Med. Chem.*, 1986, vol. 29, no. 7, pp. 1323-5.
- [43] K. L. Shyam, R. Furubayashi, R.T. Hrubiec, A. Cosby and A.C. Sartorelli, "1,2-Bis(sulfonyl)hydrazines. Effects of structural modification on antineoplastic activity," *J. Med. Chem.*, 1987, vol. 30, no.11, pp. 2157-61.
- [44] E. E. Shul'ts, G. N. Andreev, M. M. Shakirov, I. Yu. Bagryanskaya and G. A. Tolstikov, "Diels-alder reactions with cyclic sulfones: X. Synthesis of 4-carbamoyl- and 4-hydrazido-substituted hexahydro-1benzothiophene S,S-dioxides," *Russian Journal of Organic Chemistry*, 2004, vol. 45, no.10, pp. 1546- 54.
- [45] B. A. Shainyan, L.L. Tolstikova, V. I. Meshcheryakov and Y. S. Danilevich, "Trifluoromethanesulfonic Hydrazides," *Russian Journal of Organic Chemistry*, 2009, vol. 40, no.8, pp. 1071-1075.
- [46] H. Shirai, M. Yoneda and N. Oda, "Syntheses of hydrazide derivatives containing sulfone group," *Nagoya-shiritsu Daigaku Yakugakubu Kiyo*, 1954, vol. 2, pp. 45-8.
- [47] J. Safaei-Ghomi, "An Efficient Synthesis of Sulfonylhydrazides and Sulfonylsemicarbazides by Utilizing Alumina as a Catalyst," *Journal of* the Chinese Chemical Society, 2007, vol. 54, no. 6, pp. 1561-3.
- [48] R.S.Varma, "Solvent-free organic syntheses . using supported reagents and microwave irradiation," *Green Chem.*, 1999, vol.1, pp. 43-55.
- [49] A.K. Bose, B.K. Banik, N. Lavlinskaia, M. Jayaraman and M.S. Manhas, "More chemistry in a microwave," *Chemtech.*, vol. 1997, no. 27, pp. 18-24.
- [50] R.S. Varma and R. Dahiya, "An expeditious and solvent free synthesis of 2- amino substituted isoflav-3-enes using microwave irradiation," J. Org.Chem., 1998, vol. 63, pp. 8038-41.
- [51] L. Perreux, A. Loupy and F. Volatron, "Solvent-free preparation of amides from acids and primary amines under microwave irradiation," *Tetrahedron*, 2002, vol. 58, no. 11, pp. 2155-62.
- [52] E. Gelens, L. Smeets, L.A.J.M. Sliedregt, B.J. Steen, C.G. Kruse, R. Leurs and V.A. Orru, "An atom efficient and solvent-free synthesis of structurally diverse amides using microwaves," *Tetrahedron Lett*, 2005, vol. 46, no. 21, pp. 3751-4.
- [53] A. Loupy, Microwave in Organic Synthesis, Wiley-VCH, Weinheim, 2004.
- [54] R. Gedye, F. Smith, K. Westaway, H. Ali, L. Baldisera, L. Laberge and J. Rousell, "The use of microwave ovens for rapid organic synthesis," *Tetrahedron Lett.*, 1986, vol. 27, no. 3, pp. 279- 82.
- [55] Y. Li, Y. Wang, J. Li and J. Wang, "Regioselective synthesis of 1-aryl-2-p-toluenesulfonyl hydrazides under microwave irradiation," *Journal of Sulfur Chemistry*, 2007, vol. 2, no. 28, pp. 177–180.
- [56] O. P. Smirnov, V. Y. Rodionov and B. I. Stepanov, "Synthesis of symmetric diaryl disulfide S,S,S',S'-tetraoxides," *VINITI.*, 1976, pp. 2651-76.
- [57] T. Backs, C. Ollins and M. V. Rishna, "Reactions of sulfonhydrazides with benzeneseleninic acid, selenium halides, and sulfur halides. A convenient preparation of selenosulfonates and thiosulfonates," *Can. J. Chem.*, 1987, vol. 65, no. 1, pp. 38-42.
- [58] T. G. Back and H. Zhai, "Cyclizations and cycloadditions of acetylenic sulfones on solid supports," *Chem. Commun.*, 2006, pp. 326–28.
- [59] R. Ballini, E. Marcantoni and M. Petrini, "A new general synthesis of sulfones from alkyl or aryl halides and p-toluenesulfonhydrazide," *Tetrahedron*, 1989, vol. 45, no. 21, pp. 6791-8.
- [60] A. Morait and A. Mavrodin, "Studies on the thenylidene hydrazide series (II). Synthesis, physicochemical and pharmacodynamic study of some thenylidene hydrazides of p,p'-diphenyl sulfone dicarboxylic acid," *Rom. Farmacia*, 1980, vol. 28, no. 4, pp. 229-34.
- [61] K. Itaya, C. G. Gahmberg and S. Hakomori, "Cell surface labeling of erythrocyte glycoproteins by galactose oxidase and Mn⁺⁺-catalyzed coupling reaction with methionine sulfone hydrazide," *Biochemical and Biophysical Research Communications*, 1975, vol. 64, no. 3, pp. 1028-35.
- [62] G. N. Alawandi and M. V. P. G. Kulkarni, Department "Chemoselective Knoevenagel reaction as a novel route for the synthesis of biscoumarinyl methylsulphones and 4+1, 1+4 approach synthesis of aminooxadiazolyl and N-(2,5-dimethyl)pyrrolylcoumarins," *Indian Journal of Chemistry*,

International Journal of Chemical, Materials and Biomolecular Sciences ISSN: 2415-6620 Vol:7, No:6, 2013

Section B: Organic Chemistry Including Medicinal Chemistry, 2006,

- Section B: Organic Chemistry Including Medicinal Chemistry, 2006, vol. 45B, no. 1, pp. 258-66.
 [63] H. Anaraki-Ardakani, S. Sadeghian, F. Rastegari, A. Hassanabadi and M. Anary-Abbasinejad, "Three-component reaction of triphenylphosphine, acetylenic esters, and arylsulfonyl hydrazides or aryl hydrazines: an efficient one-pot synthesis of stable β-nitrogen-substituted phosphorus ylides," Synthetic Communications, 2008, vol. 28, pp. 120, pp. 1000, 0. 38, no. 12, pp. 1990– 9.
 [64] O. V. Zvereva, A. V. Milyutin, O. V. Bobrovskaya and T. F. Odegova,
- "Synthesis and Antiinflammatory and Antibacterial Activity of β -N. (Halogenobenzoyl)- and β -N-(4-Methylphenylsulfonyl)hydrazides of 4-Aryl-2-hydroxy-4-oxo-butenoic (Aroylpyruvic) Acids," *Pharmaceutical* Chemistry Journal, 2004, vol. 38, no. 2, pp. 90-2.