# Visualized Characterization of Molecular Mobility for Water Species in Foods

Yasuyuki Konishi and Masayoshi Kobayashi

**Abstract**—Six parameters, the effective diffusivity (*De*), activation energy of *De*, pre-exponential factor of *De*, amount (ASOW) of self-organized water species, and amplitude ( $\alpha$ ) of the forced oscillation of the molecular mobility ( $1/\tau_{\rm C}$ ) derived from the forced cyclic temperature change operation, were characterized by using six typical foods, squid, sardines, scallops, salmon, beef, and pork, as a function of the correlation time ( $\tau_{\rm C}$ ) of the water molecule's proton retained in the foods. Each of the six parameters was clearly divided into the water species A<sub>1</sub> and A<sub>2</sub> at a specified value of  $\tau_{\rm C} = 10^{-8} \rm s$  (=C  $\tau_{\rm C}$ ), indicating an anomalous change in the physicochemical nature of the water species at the C  $\tau_{\rm C}$ . The forced oscillation of  $1/\tau_{\rm C}$  clearly demonstrated a characteristic mode depending on the food shown as a three dimensional map associated with  $1/\tau_{\rm C}$ , the amount of self-organized water, and  $\tau_{\rm C}$ .

*Keywords*—molecular mobility, self-organization, hysteresis, water species  $A_1$  and  $A_2$ , forced cyclic temperature change operation (FCTCO)

### I. INTRODUCTION

N the field of food engineering, as has been discussed by many researchers, the water activity of foods has been conveniently employed to create a stability map [1] for various reactions in foods illustrating the appearance of a specified water activity. Water activity (aw) has commonly been used as a parameter to evaluate the deterioration of foods [2]. Water species retained in foods, as is well known, expose a multifunctional nature dynamically responding to environmental conditions, such as temperature, pressure, dehydration, and water content. For the scientific analysis of the multifunctional water species retained in foods, a recognized difficulty is related to its nonlinear dynamic change, which influences food quality.

To demonstrate the nonlinearity of the water species, the effective water diffusivity (*De*) in foods has frequently been used to evaluate the dehydration rate of food [3]. In previous studies [4]–[6], the water species retained in fish-past-sausage and squid were categorized into two species as a function of the water content. The two were water species A<sub>1</sub>, recognized as the higher water content at  $W_0 > 120\%$ -d.b., with higher water diffusivity, *De*, and water species A<sub>2</sub>,  $W_0 < 120\%$ -d.b., with lower *De*.

For the quantitative evaluation of the water species in foods, the proton NMR technique was recently applied because of the need to use physicochemical data rather than the water activity [7], [8]. Focusing on a more direct identification of the water species on the molecular level, a proton nuclear magnetic resonance technique was applied to evaluate the food quality as demonstrated in the International Conferences on Applications of Magnetic Resonance in Food Science [9]. From the viewpoint of the molecular mobility of water species, the progress of research was recently accelerated to try to demonstrate the theoretical meaning [9], [10] and applications in various researches [11]–[14].

In the present study, a dynamic proton NMR technique is proposed as a new technique. The aims of this study are (1) to identify the two water species,  $A_1$  and  $A_2$ , by using the effective diffusivity (*De*) as a three-dimensional mobility, the activation energy ( $E_D$ ) of *De*, and the pre-exponential factor of *De*; (2) to visualize the self-organization of the two water species by using the dynamic NMR technique; (3) to visualize the forced oscillation of the molecular mobility ( $1/\tau_C$ ); and (4) to demonstrate the dynamism of the amount of the self-organized water species.

### **II. EXPERIMENTAL METHODS**

### A. Materials and Methods

Pork meat ( $P_{H}$ : produced in Hokkaido), beef meat ( $B_{A}$ : produced in Australia and  $B_{H}$ : produced in Hokkaido), squid, salmon, sardine, and scallop, were chosen as food samples. The pork and beef had an initial water content of 230~320%-d.b. and 230~280%-d.b., respectively; the initial water content of the squid and the salmon commonly ranged from 300~360%-d.b., and that of sardine and scallop, from 210~270%-d.b. and 180~230%-d.b. of the initial water content, respectively. To evaluate the effective diffusivity (*De*) of water species, each sample was placed in a stainless steel net tray (4 meshes) that was mechanically hung from a strain gage transducer in a dryer. To evaluate the dehydration rate, the sample weight was continuously recorded by the output of a strain-gage transducer using a data-logger.

To evaluate the correlation time ( $\tau_c$ ) of the water species for the samples, a nuclear magnetic resonance (NMR) technique was applied to measure the <sup>1</sup>H-NMR spectra and spin-spin relaxation time ( $T_2$ ) of water protons. All the samples cut into  $2 \times 2 \times 10$  mm pieces were inserted into an NMR sample tube (4mm in inner diameter and 180mm in length). <sup>1</sup>H-NMR spectra

Yasuyuki Konishi is with the Hokkaido Industrial Technology Centre, 379 Kikyo-cho, Hakodate, Hakodate, 041-0801, Japan (phone: +81-138-34-2600; fax: +81-138-34-2602; e-mail: konishi@techakodate.or.jp).

Masayoshi Kobayashi is with the Advanced Technology Institute of Northern Resources, 8-6, Tonden 2Jo 2Chome, Sapporo, Hokkaido, 002-0852, Japan (e-mail: kmasa@f2.dion.ne.jp).

were obtained using a JEOL A-500 FT-NMR spectrometer operating at 500MHz for protons. The observed frequency width was 20 kHz. The 90° pulse width was 12.5  $\mu s$ , and the number of pulse repetitions was 8. The proton chemical shifts were measured by using a small amount of water containing deuterium oxide as an external reference. Except the forced cyclic temperature change operation (FCTCO), all the NMR measurements were performed at 23.5±0.5°C. The spin-spin relaxation times,  $T_2$ , were obtained by the spin-locking method.

### B. Evaluation of the correlation time $(\tau_C)$ of water proton

The spin-locking pulse technique used was effective to detect a very fast relaxation signal at a low water content. For the evaluation of the relaxation time,  $T_2$ , the equation of  $M_t = M_0$  $exp(-ts/T_2)$  was used, where  $M_t$  is the magnitude of the magnetization vector after the spin locking pulse,  $M_0$  is the magnitude of the macroscopic equilibrium magnetization vector, and ts is the spin locking pulse length. In the present study, the plot of  $ln [M_t / M_0]$  vs. ts indicated a good linearity (which demonstrates a correlation coefficient higher than 0.99) through all water contents of the foods, suggesting that the evaluated  $T_2$  value was reasonable. From  $T_2$ , the correlation time of a water proton,  $\tau_C$ , was evaluated using Equation (3) as described by Abragam [15]:

$$\frac{1}{T_2} = \frac{\gamma^4 \cdot \hbar^2 \cdot I(I+1)}{5r^6} \left( 3\tau_c + \frac{5\tau_c}{1+\omega_0^2 \cdot \tau_c^2} + \frac{2\tau_c}{1+4\omega_0^2 \cdot \tau_c^2} \right) , \qquad (1)$$

where  $\gamma$  is the gyromagnetic ratio of a proton (= 2.675×10<sup>8</sup> rad·T<sup>-1</sup>·s<sup>-1</sup>),  $\hbar$  is the modified Plank's constant (6.63×10<sup>-34</sup> J·s), *I* is the nuclear-spin quantum number of a water proton (= 0.5), *r* is the proton-proton distance of a water molecule (0.16 nm),  $a_0$  is the resonance frequency of NMR (= 3.14×10<sup>9</sup> s<sup>-1</sup>), and  $\tau_C$  is the correlation time of a water proton (s).

### C. Forced cyclic temperature change operation (FCTCO)

To clearly distinguish the mode for the self-organization of water species among the foods used, a forced cyclic temperature change operation (FCTCO) was effectively conducted. The time schedule for the FCTCO was as follows: when the temperature was gradually decreased, a steep reduction of  $1/\tau_{\rm C}$ at a specified temperature  $(t_1)$  appeared, and, at the peak bottom of the  $1/\tau_{\rm C}$ , when the temperature was gradually increased, the locus of the  $1/\tau_{\rm C}$  showed a typical hysteresis reaching the temperature  $(t_2)$  previously used. The FCTCO was carefully repeated between  $t_1$  and  $t_2$ , and a forced oscillation of the  $1/\tau_{\rm C}$ was then clearly visualized. In the course of the FCTCO, as an expedient way, 2 min to increase or decrease the temperature and 13 min to evaluate the  $T_2$  as the time for the operation of the NMR equipment were required; the total time of 15 min was thus needed as the measuring time at each temperature in the hysteresis period. The oscillating curves obtained were demonstrated as a function of the elapsed time including the 15 min.

### III. RESULTS AND DISCUSSION

## A. Characterization of the water species due to De (the three-dimensional mobility: 3D mobility, $m^2/s$ )

As has been demonstrated by a large number of researchers [3, 10], the three-dimensional mobility (3D mobility) of water species retained in the foods was evaluated by using an effective diffusion coefficient (De, m<sup>2</sup>/h). The value of De at a given water content could be evaluated by using Equation (2).

$$\frac{W - We}{W_D - We} = \left(\frac{8}{\pi^2}\right)^3 \cdot exp\left(\frac{\pi^2 \cdot De \cdot t}{4} \cdot \left(L_a^2 + L_b^2 + L_c^2\right)\right) (2)$$

where *W* is the water content at the drying time t (%-d.b.),  $W_e$  is the equilibrium water content (%-d.b.),  $W_D$  is the initial water content of the rectangular sample of drying flesh (%-d.b.), t is the drying time (s), and  $L_a$ ,  $L_b$ , and  $L_c$  are the half distances of the rectangular sample (m).

Since *De* is strongly influenced by the water content of food, it should be evaluated as a function of the water content ( $W_0$ ). The water content of food consequently contributes to the correlation time ( $\tau_C$ ) evaluated from the proton NMR technique and could reasonably be replaced by  $\tau_C$  instead of the water content [16].



Fig. 1 De, ED, and  $\delta$  D0 as a function of  $\tau$  C .Key:  $\bigcirc$ , BA;  $\triangle$ , BH;  $\Box$ , PH

The value of  $\tau_{\rm C}$  means a restriction in the strength of water species due to the surrounding macro-molecule as protein. The

*De* described as a function of  $\tau_{\rm C}$  provides information as to how it will be influenced by  $\tau_{\rm C}$ . Fig. 1(A) illustrates *De* for B<sub>A</sub>, B<sub>H</sub>, and P<sub>H</sub> as a function of  $\tau_{\rm C}$ . As is evident from the results, two water species A<sub>1</sub> and A<sub>2</sub> regions were clearly distinguished at  $\tau_{\rm C}$ =10<sup>-8</sup>s (designated as a critical  $\tau_{\rm C}$ ,  $C\tau_{\rm C}$ ) indicating a stepwise reduction of *De* and an identical dependency as a function of  $\tau_{\rm C}$ within each of the two regions. This anomalous change of *De* for the three foods at the C $\tau_{\rm C}$  strongly supports the idea of a physical change of the food structure derived from the progress of dehydration, such as porosity ( $\varepsilon$ ) and labyrinth factor ( $\chi$ ) of food solids. This presumption would also be supported by the following discussion.

As is well known, De can be expressed as a function of the diffusibility ( $\delta$ ) and activation energy ( $E_{\rm D}$ ) of diffusion by Equation (3).

$$De = \left(\frac{\varepsilon}{\chi}\right) \cdot D = \delta \cdot D_0 \cdot exp\left[\frac{-E_{\rm D}}{R \cdot (T_{\rm D} + 273)}\right] , \qquad (3)$$

where  $\varepsilon$  is the porosity of solid food,  $\chi$  is the labyrinth factor of food,  $\delta$  is the diffusibility (= $\varepsilon/\chi$ ), and  $D_0$  is the pre-exponential factor of the diffusion coefficient (*D*).

Although the Arrhenius plot of *De* evaluated from the data obtained at 30, 40, and 50°C was widely scattered, the activation energy ( $E_D$ ) of *De* can roughly be evaluated as a function of  $\tau_C$ , as shown in Fig. 1(B). The obtained values of  $E_D$  again demonstrate a drastic reduction at the  $C\tau_C$  as a function of  $\tau_C$ , indicating an identical value of 16 (±1.5) kJ/mol within the water species A<sub>1</sub> region independently of the kind of foods; on the other hand, in the water species A<sub>2</sub> region, the evaluation was different depending on the kind of foods, i.e., 6.1 (±0.6) kJ/mol for B<sub>A</sub>, 8.1 (±0.4) kJ/mol for B<sub>H</sub>, and 2.0 (±0.8) kJ/mol for P<sub>H</sub>, indicating no change depending on  $\tau_C$ .

Focusing on Eq. (3), the pre-exponential factor  $(\delta D_0)$  can easily be evaluated by using the extrapolation of the Arrhenius plot line against the perpendicular axis, and the values obtained are plotted as a function of  $\tau_{\rm C}$ , as shown in Fig. 1(C). The obtained value of  $\delta D_0$  clearly demonstrated a remarkable reduction at  $C\tau_{\rm C}$ . From this anomalous change of  $\delta D_0$ , the values of  $\varepsilon$  and  $D_0$  can be presumed to decrease and those of  $\chi$  to steeply increase because of the progress of dehydration. The drastic change of the three parameters ( $\varepsilon$ ,  $D_0$ , and  $\chi$ ) indicates that there is a steep change in the physical structure of the solid food influencing the reduction of De in the water species  $A_2$ region, as seen in Fig. 1(A). In this region, regarding the significance of the reduction of  $E_{\rm D}$ , it may be presumed that the energy barrier for water species A2's diffusion is reduced due to the reduction of the value of  $D_0$  in the hydration water. On the basis of this presumption, the following discussion may be offered.

In a heterogeneous catalysis [17], as an empirical principle, the compensation effect between the pre-exponential factor  $(A_0)$ and activation energy (E) for the rate constant in a heterogeneously catalyzed reaction has been widely accepted. The linear relation between the log  $A_0$  and E gives another straight line when the reaction occurs on a different active center of the catalyst surfaces associated with some difference in the reaction mechanism. This empirical principle is possibly applied to the water diffusion process in the foods. Fig. 2 demonstrates the ln ( $\delta D_0$ ) as a function of  $E_D$  for P<sub>H</sub>, B<sub>A</sub>, squid, salmon, and sardine. Although the data obtained includes an appreciable number of experimental errors, except for the case of P<sub>H</sub> and B<sub>A</sub>, each straight line among the three foods clearly gives a different tendency. This evidence strongly demonstrates that the diffusion mechanism of water is different among the three foods. Since  $D_0$  is possibly associated with the nature of the adsorption sites for the water species, the protein structure, as an adsorption site, involved in the foods strongly influenced the value of  $D_0$ . The difference among the straight lines of the  $\ln(\delta D_0) \sim E_D$  obtained for the five foods in Fig. 2, therefore, strongly suggests that  $D_0$  changes depending on the foods. Although it is difficult to describe the physical meaning of the variation of  $D_0$ , it may be presumed that the number of adsorbed water molecules and the structure of the hydration water on the adsorption sites derived from the mechanical structure change of the food solid will change depending on the foods, as discussed in the above section using Fig. 1(C).



Fig. 2 Compensation effect derived from the ln (  $\delta$  D0 ) as a function of ED. Key:  $\bigcirc$ , PH ;  $\triangle$ , BA ;  $\bigcirc$ , squid ;  $\blacktriangle$ , salmon ;  $\blacksquare$ , sardine.

B. Characterization of the water species due to the molecular mobility  $(1/\tau_{c}, s^{-1})$ 

As has been described in the above section, the molecular mobility  $(1/\tau_{\rm C}, \rm s^{-1})$  can be easily evaluated by using the proton NMR technique. The value of  $1/\tau_{\rm C}$  obtained for all the foods demonstrates the self-organization of the water species retained in the food, which was caused by a programmed reduction in temperature. Fig. 3 demonstrates the  $\ln(1/\tau_{\rm C})$  as a function of 1/T for the scallop as a typical example. The self-organization of the water species is clearly evident at a specified temperature, which influences the water content of the sample (this can be replaced by the initial correlation time,  $\tau_{\rm C}$ , of the food sample). Since the molecular mobility  $(1/\tau_{\rm C}, \rm s^{-1})$  is the rotation rate of the water proton, the  $1/\tau_{\rm C}$  obtained at the steeply reduced value can be plotted as a function of 1/T. The Arrhenius plots obtained for various initial  $\tau_{\rm C}$  of the food samples evaluate Eso(I) = 29.5

(±3.0) for the water species  $A_1$  and  $Eso(II) = 57.6 (\pm 2.6) \text{ kJ/mol}$  for the water species  $A_2$ . All the values of Eso (I) and Eso (II) for the six foods were evaluated as presented in Table I. Each of the Eso obtained gave a characteristic value depending on the kind of foods, indicating the self-organization of the water species to be influenced by different environmental conditions due to the macromolecules of each type of food, such as proteins.



Fig. 3 Evaluation of the activation energy for the self-organization of scallop derived from the  $\ln(1/\tau_{\rm C})$  as a function of 1/T.  $W_0$  and the initial  $\tau_{\rm C}$  are respectively evaluated as the Key:  $\circ$ , 197%-d.b.,  $6.53 \times 10^{-9}$ s;  $\Delta$ , 162%-d.b., 7.58×10<sup>-9</sup>s;  $\bullet$ , 126%-d.b., 1.01×10<sup>-8</sup>s;  $\blacktriangle$ , 104%-d.b., 1.23×10<sup>-8</sup>s;  $\blacksquare$ , 64%-d.b., 4.37×10<sup>-8</sup>s.

 $\begin{tabular}{l} TABLE I \\ Activation Energies Of Self-Organization For Water Species A_1 \\ And A_2 Retained In B_A, B_H, P_H, Squid, Salmon, Sardine, And Scallop \\ \end{tabular}$ 

	$E_{\rm SO}$ (kJ/mol)	
	Species A <sub>1</sub>	Species A <sub>2</sub>
$\mathbf{B}_{\mathbf{A}}$	31	42
$P_{H}$	49	98
Squid	65	67
Salmon	160	63
Sardine	183	116
Scallop	30	58

Focusing on the character difference of the self-organization between water species  $A_1$  and  $A_2$ , each of which produces a different activation energy, it is noteworthy that the two water species can be identified from their molecular states associated with the pre-exponential factor,  $As_0$ , in the Arrhenius equation,  $\ln (1/\tau_C) = As_0 \exp (-Eso/RT)$ . To respond to a question how the adsorption sites for the two water species are discriminated, the compensation effect would be considered as discussed in Fig.1(C). Fig. 4 demonstrates ln ( $As_0$ ) as a function of *E*so for squid, salmon, sardine, P<sub>H</sub>, and B<sub>H</sub>. All the data for each food agreed, indicating that an identical compensation effect was in effect between the ln ( $As_0$ ) and *E*so. This result strongly demonstrates that there is no difference in the water species as hydration water among the foods because of the similarity of the hydration water even though water species  $A_1$  and  $A_2$  can be distinguished by the difference in their self-organization temperature. This evidence differs from the compensation effect between the  $\delta D_0$  and  $E_D$  for the 3D mobility (*De*), indicating a clear difference among the foods evaluated at 30~70°C, as described in Fig. 2. From these results, it is evident that the self-organization of the water species occurred as a result of an identical physical mechanism that was independent of the difference between water species  $A_1$  and  $A_2$  and among the foods.



Fig. 4 Compensation effect derived from the  $\ln(A_0)$  as a function of  $E_{SO}$ . Key:  $\circ$ ,  $P_H$ ;  $\Delta$ ,  $B_A$ ;  $\bullet$ , squid;  $\blacktriangle$ , salmon;  $\blacksquare$ , sardine;  $\blacklozenge$ , scallop.

## *C.* Visualization of the forced oscillation of the molecular mobility $(1/\tau_{c}, s^{-1})$

Focusing on the hysteresis behavior of the scallop in Fig. 3, the forced oscillating behavior of the molecular mobility  $(1/\tau_{\rm C})$  can be easily visualized by using a forced cyclic temperature change operation (FCTCO) between two specified temperatures (designated as  $t_1$  and  $t_2$ ) at which the hysteresis behavior of  $1/\tau_{\rm C}$  appears. Fig. 5 demonstrates a schematic explanation for the FCTCO. The temperature was cyclically scheduled by decreasing and increasing  $t_1$  and  $t_2$ , respectively. To evaluate the value of  $\tau_{\rm C}$  using the proton NMR spectra obtained at the given temperature, about 15min was needed for the temperature to



Fig. 5 Schematic visualization of the forced cyclic temperature change operation (FCTCO), the NMR-spectra dynamism, and the  $1/\tau_{c}$ -oscillation obtained

decrease and increase and for the evaluation of  $T_2$ . The values of  $1/\tau_C$  obtained at each temperature were visualized while taking the 15min into the elapsed time as an oscillating curve. The oscillating curve obtained was visualized as a function of time, as demonstrated in the figure.

The forced oscillations of the  $1/\tau_{\rm C}$  obtained were visualized for all the foods taken in this study. Fig. 6 demonstrates the forced oscillating curves for sardine, scallop, and P<sub>H</sub> as typical examples. As is evident in the oscillating response curves obtained, the amplitude, period, and waveform were changed depending on the kind of foods and the water content, which can be replaced by the initial  $\tau_{\rm C}$  of the sample. The amplitude (*a*) is drastically reduced in the water species A<sub>2</sub> region, indicating the reduction of the molecular mobility ( $1/\tau_{\rm C}$ ). This reduction of *a*, in the case of scallop, can be easily recognized from the diagram in Fig. 7 by demonstrating a width of  $\ln(1/\tau_{\rm C})$  given as a distance



Fig. 6 Oscillating dynamism characterized by the kind of foods



Fig. 7 Width of  $ln(1/\tau_C)$  for the forced oscillation as a function of the initial  $\tau_C$  for scallop

between the peak top and peak bottom of the oscillating curves in Fig. 6. Since the  $\alpha$  is demonstrated as a function of the initial  $\tau_{\rm C}$  of the sample, as shown in Fig. 8, the forced oscillations for squid, sardine, and scallop disappeared at the initial  $\tau_{\rm C}$ =3.1×10<sup>-8</sup>, 6.9×10<sup>-8</sup>, and 4.4×10<sup>-8</sup> s, respectively. In addition, since the slopes of the  $\alpha \sim \tau_{\rm C}$  straight lines in the species A<sub>2</sub> region gave a specified value depending on the food, designated as  $\Delta \alpha \sim \tau_{\rm C}$ , the  $\Delta \alpha \sim \tau_{\rm C}$  obtained for each of the foods can be used to visualize the difference among the characteristic modes of the foods, as is presented in Fig. 11.



Fig. 8  $\alpha$  as a function of the initial  $\tau_C$  . Key:  $\circ$ , squid ;  $\Box$ , scallop ;  $\Box$ , sardine

D.Visualization for the hysteresis behavior of the self-organized water amount (ASOW)

The amount of the self-organized water species (ASOW) can be evaluated by an examination of the graphical integration of the proton NMR spectrum presented in Fig. 5 and expressed as a modified water content (MWC, %-d.b.). The MWC was evaluated by using the initial water content of the sample food corresponding to the peak area of the proton NMR spectra. Fig. 9 demonstrates the MWC as a function of 1/T and the initial  $\tau_{\rm C}$ . It is clearly evident that the hysteresis of the MCW is similar to the  $1/\tau_{\rm C}$  hysteresis presented in Fig. 3, indicating the steep reduction of MWC at the same temperature as the  $1/\tau_{\rm C}$ .

Since the reduced amount of the MWC clearly indicates the ASOW, the amount is linearly reduced by increasing the initial  $\tau_{\rm C}$  of the sample. Fig. 10 demonstrates, as an example, the ASOW as a function of the initial  $\tau_{\rm C}$  for scallop. All the other foods showed similar results. Based on the results obtained, the linear dependency of the ASOW~ $\tau_{\rm C}$  straight line indicates a inflection point at  $\tau_{\rm C} = 10^{-8}$ s, pointing again to the existence of two water species as water species A<sub>1</sub> and A<sub>2</sub>, as shown in Fig. 1.



Fig. 9 The converted water content for scallop as a function of 1/T and the initial  $\tau_{\rm C}$ . Key:  $\circ$ ,  $6.5 \times 10^{-9} {\rm s}$ ;  $\Box$ ,  $7.9 \times 10^{-9} {\rm s}$ ;  $\Box$ ,  $1.0 \times 10^{-9} {\rm s}$ ;  $\bullet$ ,  $1.2 \times 10^{-8} {\rm s}$ ;  $\blacktriangle$ ,  $2.1 \times 10^{-9} {\rm s}$ ;  $\blacksquare$ ,  $4.4 \times 10^{-9} {\rm s}$ .



Fig. 10 ASOW as a function of the initial  $\tau_C$  for scallop.

In the water species  $A_2$  region in Fig. 10, the slope of the ASOW~ $\tau_{\rm C}$  straight line clearly changes depending on the food used. The slope may be designated as  $\Delta ASOW \sim \tau_C$ . The values of the  $\Delta ASOW{\sim}\tau_C$  for  $P_H,~B_A,$  squid, salmon, sardine, and scallop are evaluated to be  $-1.15 \times 10^9$ ,  $-8.19 \times 10^8$ ,  $-1.41 \times 10^9$ , -5.49×10<sup>9</sup>, -4.61×10<sup>9</sup>, and -1.18×10<sup>9</sup> %-d.b./s, respectively. The difference of the  $\triangle ASOW \sim \tau_{C}$  among the six foods would be recognized as the difference in the acceleration of the self-organization of water species in the foods, strongly associated with the  $\alpha$ 's in Fig. 6. It may be presumed that the  $\Delta$ ASOW~ $\tau_{\rm C}$  is larger with a higher  $\alpha$ . This assumption can be visualized by Fig. 11. Fig. 11 demonstrates a three-dimensional plot of  $\ln(1/\tau_{\rm C})$  as a function of the initial  $\tau_{\rm C}$  and the  $\Delta$ ASOW~ $\tau_{\rm C}$ for  $P_{\rm H}$ , squid, and salmon as typical foods. The  $\alpha$  can be expected to sensitively change depending on the kind of food, indicating a characteristic behavior. These results show that the ASOW strongly contributes to the mode of the oscillation wave and  $\alpha$ . From these results, the mode of the oscillating width can be used as the difference of the water species in the foods

indicating a characteristic dynamism of the hydration water depending on the value of the initial  $\tau_{\rm C}$ .



Fig. 11 Three-dimensional visualization for the forced oscillation of the molecular mobility as a function of the  $\Delta ASOW \sim \tau_C$  and the initial  $\tau_C$ . Key in the A<sub>1</sub> region:  $\bigcirc$ , squid;  $\triangle$ , salmon;  $\square$ , P<sub>H</sub>; and Key in the A<sub>2</sub> region:  $\bigcirc$ , squid;  $\blacktriangle$ , salmon;  $\blacksquare$ , P<sub>H</sub>.

### IV. CONCLUSIONS

The water species retained in the six foods was clearly separated into two categories as species  $A_1$  and  $A_2$ . The two species were characterized by two points of view, the effective water diffusivity (*De*, three-dimensional mobility, m<sup>2</sup>/s) and the molecular mobility ( $1/\tau_c$ , reciprocal correlation time, s<sup>-1</sup>). Based on the experimental evidence obtained, the following conclusions were demonstrated.

(1) *De* and the activation energy  $(E_D)$  of *De* were drastically changed at  $C\tau_C$ , at which the water species was divided into A<sub>1</sub> and A<sub>2</sub> in all the foods. This anomalous change was understood as a remarkable change in the physical structure of the food solids derived from the progress of the dehydration. This change brought about a change in a diffusion mechanism.

(2) The molecular mobility  $(1/\tau_c)$  in all the foods clearly demonstrated a characteristic hysteresis at a specified temperature lower than 0°C, indicating a self-organization of the water species (SOW). The hysteresis of the SOW visualized a characteristic forced oscillation derived from the forced cyclic temperature change operation (FCTCO) indicating a reproducibility of the SOW.

(3) The obtained forced oscillation of the  $1/\tau_{\rm C}$  clearly demonstrated a characteristic amplitude ( $\alpha$ ), period, frequency, and oscillation wave depending on the kind of foods and the two water species. The  $\alpha$  demonstrated a drastic reduction at C $\tau_{\rm C}$ .

(4) The ASOW was quantitatively evaluated by using the peak area of the proton NMR spectrum and determined as a function of the initial  $\tau_{\rm C}$  of the sample foods. The amount was changed depending on the kind of foods and understood as the difference of the hydration water amount retained in the foods.

(5) The oscillating behavior of  $1/\tau_C$  clearly demonstrated as a three-dimensional visualization using two parameters, the initial  $\tau_C$  and the  $\Delta ASOW \sim \tau_C$ . The results interestingly visualized the

water species to be clearly divided into two water species and to have a variety of oscillations depending on the foods.

#### NOMENCLATURES

### pre-exponential factor of k (kJ/mol)

- $A_0$  $As_0$ pre-exponential factor of the molecular mobility  $1/\tau_{\rm C}$ (kJ/mol)
- ASOW the amount of the self-organized water (%-d.b.)  $\Delta ASOW \sim \tau_C$  the slope of the ASOW  $\sim \tau_C$  straight line (%-d.b./s)
- $B_{\rm A}$ beef meat produced in Australia (-)
- $B_{\rm H}$ beef meat produced in Hokkaido, Japan (-)
- D water diffusion coefficient (m<sup>2</sup>/h)
- $D_0$ frequency factor of D (m<sup>2</sup>/h)
- effective water diffusion coefficient (m<sup>2</sup>/h) De
- $De^0$ pre-exponential factor of  $De (=\delta \cdot D_0, m^2/h)$
- Ε activation energy of the heterogeneous catalysis (kJ/mol)
- $E_{\rm D}$ activation energy of water diffusivity (kJ/mol)
- activation energy of the self-organization (kJ/mol) Eso
- FCTCO forced cyclic temperature change operation (-)
- I nuclear spin quantum number of water proton (=0.5) (-)
- half distance of a-axis of the rectangular sample (m)  $L_{a}$
- $L_{\rm b}$ half distance of b-axis of the rectangular sample (m)
- half distance of c-axis of the rectangular sample (m)  $L_{\rm c}$
- $M_{\rm t}$ magnitude of magnetization vector (-)
- $M_0$ magnitude of macroscopic equilibrium magnetisation vector (-)
- MWC *modified water content* (%-d.b.)
- pork meat produced in Hokkaido, Japan (-)  $P_{\rm H}$
- R gas constant (=8.314J/K·mol)
- *r* proton-proton distance of water molecule (= 0.16 nm)
- SOW self-organization of the water species (-)
- $\Delta S$ activation entropy (kJ/K·mol)
- Т temperature at the time to evaluate  $T_2$  (K)
- $T_2$ spin-spin relaxation time of water proton (s)
- $T_{\rm D}$ *drying temperature* (°C)
- *drying time* (s)
- the initiation temperature of the self-organization of water ti species in the course of the FCTCO (°C)
- the returning temperature of the self-organization of water t2 species in the course of the FCTCO (°C)
- ts spin locking pulse length (s)
- W water content at the drying time t (%-d.b.)
- $W_0$ initial water content for the given sample (%-d.b.)
- $W_{\rm D}$ initial water content of drying flesh sample (%-d.b.)
- We equilibrium water content (%-d.b.)
- Greek letters
  - amplitude of the forced oscillation response (-) α
  - porosity of the food tissue (-) З
  - the ratio of the circumference of a circle to its diameter π (=3.14)
  - gyromagnetic ratio of proton (= $2.675 \times 10^8 \text{ rad} \cdot \text{T}^{-1} \cdot \text{s}^{-1}$ ) γ
  - ħ modified Plank's constant (= $6.63 \times 10^{-34} \text{ J} \cdot \text{s}$ )
  - resonance frequency  $(=3.14 \times 10^9 \text{ s}^{-1})$  $\omega_{\rm b}$
  - correlation time of water proton (s)  $\tau_{\rm C}$
  - critical correlation time of water proton (s)  $C\tau_{\rm C}$
  - labyrinth factor of the food tissue (-) χ
  - *diffusibility* (= $\varepsilon / \chi$ ) (-)

### **ACKNOWLEDGMENTS**

This work was financially supported by the Cooperation of Innovation Technology and Advanced Research in Evolution Area (City Area) from the Japanese Ministry of Education, Culture, Sports, Science, and Technology. The authors wish to thank Associate Professor Koichi Miura, Kitami Institute of Technology, for his effective assistance with the proton NMR analysis.

#### REFERENCES

- S. J. Schmidt, Water and Solids Mobility in Foods, Advances in food and [1] nutrition research, volume 48, Edited by S.L. Taylor, Elsevier, 2004.
- [2] O. Fennema, In principles of food science, Part 1, Marcel Dekker, New York 1976
- [3] A. C. Jason, A Study of evaporation and diffusion processes in the drying of fish muscle in fundamental aspects of dehydration of food stuffs, ed. Society of Chemical Industry. McMillan, London, pp.103-134, 1958.
- [4] Y. Konishi, J. Horiuchi, and M. Kobayashi, Dynamic evaluation of the dehydration response curves of foods characterized by a poultice-up process using a fish-paste sausage-I. Determination of the mechanisms for moisture transfer, Drying Technology, 19(7), pp1253-1269, 2001.
- Y. Konishi, J. Horiuchi, and M. Kobayashi, Dynamic evaluation of the [5] dehydration response curves of foods characterized by a poultice-up process using a fish-paste sausage-II, A new tank model for a computer simulation, Drying Technology, 19(7), pp1271-1285, 2001.
- Y. Konishi, K. Miura, and M. Kobayashi, Drying efficiency design using [6] multifunctional dynamics of water molecules in foods-H-NMR analysis of a fish paste sausage and squid, AIDIC Conference series, vol.6, pp183-190, 2003.
- L. B. Rockland, and G. F. Stewart, Eds., International symposium on [7] properties of water, water activity: Influences on food quality, Academic Press Inc., London, 1981.
- [8] P. S. Belton, A. M. Gil, G. A. Webb, and D. Rutledge, Eds., Magnetic resonance in food science-latest developments, The Royal Society of Chemistry, Cambridge., 2003.
- B. P. Hills, NMR studies of water mobility in foods. In"Water [9] Management in the Design and Distribution of Quality Foods" ISOPOW 7(Y.H. Roos, R.B. Leslie, and P.J.Lillford, eds) Technomic Publishing, Lancaster, PA., 1999.
- [10] B. P. Hills, C. E. Manning, and J. A. Godward, A multistate theory of water relations in biopolymer systems. In: Advances in Magnetic Resonance in Food Science (edited by P. S. Belton, B. P. Hill and G. A. Webb). pp.45-62. Cambridge, UK: Royal Society of Chemistry, 1999.
- [11] E. Vittadini, and P. Chinachoti, Effect of physico-chemical and molecular mobility parameters on Staphylococcus aureus growth. International Journal of Food Science and Technology, 38, pp841-847, 2003.
- [12] Y. Konishi, and M. Kobayashi, Quantitative evaluation of the design parameters requested in beef and pork drying operation, AIDIC Conference Series, 9, pp 177-186, 2009.
- [13] Y. Konishi, M. Kobayashi, and K. Miura, Characeterization of water species revealed in the drying operation of Todarodes pacificus Steenstrup using water proton NMR analysis, International Journal of Food Science and Technology., 45. pp1889-1894, 2010.
- [14] Y. Konishi, M. Kobayashi, and Y. Kawai, Bacterial growth trend of a dried Japanese common squid (Todarodes pacificus Steenstrup) characterised by dehydration, International J.Food Science and Technology, 46, pp 2035-2041, 2011.
- [15] A. Abragam, The Principles of Nuclear Magnetism. Oxford at the Clarend Press, p347-349, 1963.
- [16] Y. Konishi, and M. Kobayashi, Dynamism of the water species as a probe molecule in food, Chemical Engineering Transactions, Ed. Sauro Pierucci, 24, pp 475-480, 2011.
- [17] H. M. C. Sosnovsky, The catalytic activity of silver crystals of various orientations after bombardment with positive ions,. J.Phys.Chem. Solids, Pergan press, 10, pp 304-310, 1959.