

A new robust stability criterion for dynamical neural networks with mixed time delays

Guang Zhou and Shouming Zhong

Abstract—In this paper, we investigate the problem of the existence, uniqueness and global asymptotic stability of the equilibrium point for a class of neural networks, the neural system has mixed time delays and parameter uncertainties. Under the assumption that the activation functions are globally Lipschitz continuous, we drive a new criterion for the robust stability of a class of neural networks with time delays by utilizing the Lyapunov stability theorems and the Homomorphic mapping theorem. Numerical examples are given to illustrate the effectiveness and the advantage of the proposed main results.

Keywords—Neural networks, Delayed systems, Lyapunov function, Stability analysis.

I. INTRODUCTION

IN recent years, neural networks have been widely used in solving various classes of engineering problems such as control systems, optimization, image processing, associative memory design and signal processing. The key feature of the designed neural network, in such applications, is to be convergent. When a neural network is designed to function as an associative memory, it is desired that the neural network has multiple equilibrium points. Therefore, there has been a great deal of interest to the stability properties of neural networks in the past literature. When a neural network is employed to solve optimization problems, then the neural network must have unique equilibrium point which is globally asymptotically stable. But, in hardware implementation of neural networks, some parameters associated with the dynamical behavior of neural network may be subjected to some changes. Therefore, in order to be able to completely characterize equilibrium and stability properties of the neural network, we must take into account the delay parameters and uncertainties in the mathematical model of the neural network. In the recent literature, many papers have studied the existence, uniqueness and global robust asymptotic stability of the equilibrium point for different classes of delayed neural networks and presented various robust stability conditions [1]-[15]. In [1], the author have applied many methods to study the existence, uniqueness and global asymptotic stability of the equilibrium point for the class of neural networks with multiple time delays and

parameter uncertainties, and got a new robust stability criterion. But the paper have conservation. In the current paper, we will present a new alternative sufficient condition for global robust stability of delayed neural networks with multiple and distributed time delays. At the end of this paper we will give two numerical examples to clarify the problem which we study.

II. PROBLEM STATEMENT

The dynamical behavior of the neural network we consider is assumed to be governed by the following system of ordinary differential equations:

$$\begin{aligned} \frac{dx_i(t)}{dt} = & -c_i x_i(t) + \sum_{j=1}^n a_{ij} f_j(x_j(t)) \\ & + \sum_{j=1}^n b_{ij} f_j(x_j(t - \tau_{ij})) \\ & + \sum_{j=1}^n d_{ij} \int_{t-\sigma}^t f_j(x_j(s)) ds + u_i, i = 1, 2, \dots, n. \end{aligned} \quad (1)$$

Where n is the number of the neurons, $x_i(t)$ denotes the state of the neuron i at time t , $f_i(\cdot)$ denotes activation functions, a_{ij} , b_{ij} and d_{ij} denote the strengths of connectivity between neurons j and i ; τ_{ij} and σ represent the time delay required in transmitting a signal from the neuron i , u_i is the constant input to the neuron i , c_i is the charging rate for the neuron i .

In order to complete characterize the equilibrium and stability properties of the neural network defined by (1), it will be assume that the parameters a_{ij} and b_{ij} and c_i and d_{ij} of neural system (1) are uncertain and have bounded norms with being defined in the following intervals:

$$\begin{aligned} C_I & := \{C = \text{diag}(c_i) : 0 < \underline{C} \leq C \leq \overline{C}, \\ & \quad \text{i.e., } 0 < \underline{c}_i \leq c_i \leq \overline{c}_i, i = 1, 2, \dots, n\} \\ A_I & := \{A = (a_{ij})_{n \times n} : 0 < \underline{A} \leq A \leq \overline{A}, \\ & \quad \text{i.e., } 0 < \underline{a}_{ij} \leq a_{ij} \leq \overline{a}_{ij}, i, j = 1, 2, \dots, n\} \\ B_I & := \{B = (b_{ij})_{n \times n} : 0 < \underline{B} \leq B \leq \overline{B}, \\ & \quad \text{i.e., } 0 < \underline{b}_{ij} \leq b_{ij} \leq \overline{b}_{ij}, i, j = 1, 2, \dots, n\} \\ D_I & := \{D = (d_{ij})_{n \times n} : 0 < \underline{D} \leq D \leq \overline{D}, \\ & \quad \text{i.e., } 0 < \underline{d}_{ij} \leq d_{ij} \leq \overline{d}_{ij}, i, j = 1, 2, \dots, n\} \end{aligned} \quad (2)$$

In order to achieve the task of finding conditions that ensure robust stability of neural network (1), we need to prove that the conditions to be obtained must guarantee that the unique equilibrium point of system (1) is globally asymptotically stable for all $C \in C_I, A \in A_I, B \in B_I, D \in D_I$. Therefore, our main goal will be studying the dynamical analysis of neural network (1) under the parameter uncertainties defined by (2).

This work was supported by the National Basic Research Program of China(2010CB732501) and the National Nature Science Foundation of China(61273015).

Guang Zhou and Shouming Zhong are with the School of Mathematics Science, University Electronic Science and Technology of China, Chengdu 611731, PR China.

Shouming Zhong is with Key Laboratory for NeuroInformation of Ministry of Education, University of Electronic Science and Technology of China, Chengdu 611731, PR China. Email address: zhoguang@163.com.

We will assume that the functions f_i are Lipschitz conditions satisfying

$$|f_i(x) - f_i(y)| \leq \ell_i |x - y|, i = 1, 2, \dots, n, \forall x, y \in R, x \neq y.$$

Where $\ell_i > 0$ denotes a Lipschitz constant. This class of function is denote by $f \in \mathcal{L}$.

In order to obtain our robust stability result, we will need to make use of some commonly used vector and matrix norms. Let $v = (v_1, v_2, \dots, v_n)^T$ be a vector of dimension n and $Q = (q_{ij})_{n \times n}$ be a real $n \times n$ matrix. For $v = (v_1, v_2, \dots, v_n)^T$ and $Q = (q_{ij})_{n \times n}$, $|v|$ will be denote $(|v_1|, |v_2|, \dots, |v_n|)^T$ and $|S|$ will be denote $|Q| = (|q_{ij}|)_{n \times n}$. Then, we consider the following norms:

$$\begin{aligned} \|v\|_1 &= \sum_{i=1}^n |v_i|, \\ \|v\|_2 &= \{\sum_{i=1}^n |v_i|^2\}^{\frac{1}{2}}, \\ \|v\|_\infty &= \max_{1 \leq i \leq n} |v_i|, \\ \|Q\|_1 &= \max_{1 \leq i \leq n} \sum_{j=1}^n |q_{ji}|, \\ \|Q\|_2 &= [\lambda_{\max}(Q^T Q)]^{\frac{1}{2}}, \\ \|Q\|_\infty &= \max_{1 \leq i \leq n} \sum_{j=1}^n |q_{ij}|. \end{aligned}$$

Lemma 1 [2]. If the map $H(x) \in C^0$ satisfies the following conditions:

- (i) $H(x) \neq H(y)$ for all $x \neq y$,
- (ii) $\|H(x)\| \rightarrow \infty$ as $\|x\| \rightarrow \infty$,

then, $H(x)$ is homeomorphism of R^n .

Lemma 2 [3]. Let A be any real matrix such that $A \in A_I$ with A_I being defined as

$$A_I := \{A = (a_{ij})_{n \times n} : 0 < \underline{A} \leq A \leq \bar{A}, \text{ i.e., } 0 < \underline{a}_{ij} \leq a_{ij} \leq \bar{a}_{ij}, i, j = 1, 2, \dots, n\}$$

Then, the following inequality holds:

$$\|A\|_2 \leq \|\hat{A}\|_2$$

where $\hat{A} = (\hat{a}_{ij})_{n \times n}$ with $\hat{a}_{ij} = \max\{|a_{ij}|, |\bar{a}_{ij}|\}$.

Lemma 3 [4]. Let A be any real matrix such that $A \in A_I$ with A_I being defined as

$$A_I := \{A = (a_{ij})_{n \times n} : 0 < \underline{A} \leq A \leq \bar{A}, \text{ i.e., } 0 < \underline{a}_{ij} \leq a_{ij} \leq \bar{a}_{ij}, i, j = 1, 2, \dots, n\}$$

Then, the following inequality holds:

$$\|A\|_2 \leq \sqrt{\|A^*\|_2^2 + \|A_*\|_2^2 + 2\|A_*^T | A^* \|_2}$$

where $A^* = \frac{1}{2}(\bar{A} + \underline{A})$, $A_* = \frac{1}{2}(\bar{A} - \underline{A})$.

Lemma 4 [4]. Let A be any real matrix such that $A \in A_i$ with A_i being defined as

$$A_i := \{A = (a_{ij})_{n \times n} : 0 < \underline{A} \leq A \leq \bar{A}, \text{ i.e., } 0 < \underline{a}_{ij} \leq a_{ij} \leq \bar{a}_{ij}, i, j = 1, 2, \dots, n\}.$$

Then, the following inequality holds:

$$\|A\|_2 \leq \|A^*\|_2 + \|A_*\|_2$$

where $A^* = \frac{1}{2}(\bar{A} + \underline{A})$, $A_* = \frac{1}{2}(\bar{A} - \underline{A})$.

Lemma 1 is very important for the prove of the existence and uniqueness of equilibrium point of the neural network defined by (1). And others define different upper bound norms for the interval matrices, which will play key roles in the following proofs.

III. MAIN RESULT

Now, we can studying the robust stability of the equilibrium point defined by (1).

1. Existence and uniqueness of equilibrium point.

Theorem 1. For the neural network defined by (1), assume that the network parameters satisfy (2) and $f \in \mathcal{L}$. Then, the neural network model (1) has a unique equilibrium point for every input vector $u = [u_1, u_2, \dots, u_n]^T$, if the following condition holds:

$$\varepsilon = c_m - \ell_M \|Q\|_2 - \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} - \lambda_{\max}(D^*) > 0$$

where $c_m = \min(c_i)$, $\ell_M = \max(\ell_i)$, $\|Q\|_2 = \min\{\|A^*\|_2 + \|A_*\|_2, \sqrt{\|A^*\|_2^2 + \|A_*\|_2^2} + 2\|A_*^T | A^* \|_2, \|\hat{A}\|_2\}$, with $A^* = \frac{1}{2}(\bar{A} + \underline{A})$, $A_* = \frac{1}{2}(\bar{A} - \underline{A})$, $\hat{A} = (\hat{a}_{ij})_{n \times n}$, $\hat{a}_{ij} = \max\{|a_{ij}|, |\bar{a}_{ij}|\}$, $\hat{B} = (\hat{b}_{ij})_{n \times n}$, with $\hat{b}_{ij} = \max(|b_{ij}|, |\bar{b}_{ij}|)$, $D^* = (d_{ij}^*)_{n \times n}$, $d_{ij}^* = \frac{|d_{ij}| \sigma \ell_j + |d_{ji}| \sigma \ell_i}{2}$, λ_i is eigenvalue of D^* , and $\lambda_{\max} = \max_{1 \leq i \leq n} \{\lambda_i\}$, $i = 1, 2, \dots, n$.

Proof: In order to proceed with proof of the existence and uniqueness, we consider the following mapping associated with system (1).

$$H(x) = -Cx + Af(x) + Bf(x) + D \int_{t-\sigma}^t f(x(s)) ds + u \quad (3)$$

Let x^* be an equilibrium point of Eq.(1), then we have

$$H(x^*) = -Cx^* + Af(x^*) + Bf(x^*) + D \sigma f(x^*) + u = 0$$

Note that $H(x) = 0$ is an equilibrium point of (1). Therefore, it follows from Lemma 1 that, for the system defined by (1), there exist a unique equilibrium point for every input vector u if $H(x)$ is homeomorphism of R^n . We now prove that $H(x)$ is homeomorphism of R^n . we choose two vectors $x, y \in R^n$ such that $x \neq y$. For $H(x)$ defined by (3), we can write

$$H(x) - H(y) = -C(x - y) + A(f(x) - f(y)) + B(f(x) - f(y)) + D \sigma (f(x) - f(y))$$

Next, we multiply both sides by $(x - y)^T$, we get

$$\begin{aligned} (x - y)^T (H(x) - H(y)) &= -(x - y)^T C (x - y) \\ &+ (x - y)^T A (f(x) - f(y)) + (x - y)^T B (f(x) - f(y)) \\ &+ (x - y)^T D \sigma (f(x) - f(y)) \\ &= -\sum_{i=1}^n c_i (x_i - y_i)^2 + (x - y)^T A (f(x) - f(y)) \\ &+ \sum_{i=1}^n \sum_{j=1}^n b_{ij} (x_i - y_i) (f_j(x_j) - f_j(y_j)) \\ &+ (x - y)^T D \sigma (f(x) - f(y)). \end{aligned} \quad (4)$$

We can write the following inequalities

$$-\sum_{i=1}^n c_i (x_i - y_i)^2 \leq -\sum_{i=1}^n c_i (x_i - y_i)^2 \leq -c_m \sum_{i=1}^n (x_i - y_i)^2 = -c_m \|x - y\|_2^2 \quad (5)$$

We also get that

$$\begin{aligned} (x - y)^T A (f(x) - f(y)) &\leq \|A\|_2 \|x - y\|_2 \|f(x) - f(y)\|_2 \\ &\leq \ell_M \|A\|_2 \|x - y\|_2^2 \\ &\leq \ell_M \|Q\|_2 \|x - y\|_2^2 \end{aligned} \quad (6)$$

and

$$\begin{aligned} & \sum_{i=1}^n \sum_{j=1}^n b_{ij} (x_i - y_i) (f_j(x_j) - f_j(y_j)) \\ & \leq \sum_{i=1}^n \sum_{j=1}^n |b_{ij}| |x_i - y_i| |f_j(x_j) - f_j(y_j)| \\ & \leq \sum_{i=1}^n \sum_{j=1}^n |b_{ij}| \ell_j |x_i - y_i| |x_j - y_j| \\ & \leq \ell_M \sum_{i=1}^n \sum_{j=1}^n |b_{ij}| \frac{1}{2} (\sqrt{\frac{\|B\|_1}{\|B\|_\infty}} (x_i - y_i)^2 \\ & + \sqrt{\frac{\|B\|_\infty}{\|B\|_1}} (x_j - y_j)^2) \\ & = \frac{1}{2} \ell_M \sum_{i=1}^n \sum_{j=1}^n (|b_{ij}| \sqrt{\frac{\|B\|_1}{\|B\|_\infty}} \\ & + |b_{ji}| \sqrt{\frac{\|B\|_\infty}{\|B\|_1}}) (x_i - y_i)^2 \\ & \leq \frac{1}{2} \ell_M (\|B\|_\infty \sqrt{\frac{\|B\|_1}{\|B\|_\infty}} + \|B\|_1 \sqrt{\frac{\|B\|_\infty}{\|B\|_1}}) \|x - y\|_2^2 \\ & = \ell_M \sqrt{\|B\|_1 \|B\|_\infty} \|x - y\|_2^2 \end{aligned}$$

From (2), it is easy to get $\|B\|_1 \leq \|\hat{B}\|_1$ and $\|B\|_\infty \leq \|\hat{B}\|_\infty$. So, we obtain the following inequalities

$$\begin{aligned} & \sum_{i=1}^n \sum_{j=1}^n b_{ij} (x_i - y_i) (f_j(x_j) - f_j(y_j)) \\ & \leq \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} \|x - y\|_2^2 \end{aligned} \tag{7}$$

$$\begin{aligned} & (x - y)^T D \sigma(f(x) - f(y)) \\ & = \sum_{i=1}^n \sum_{j=1}^n (x_i - y_i) d_{ij} \sigma(f_j(x_j) - f_j(y_j)) \\ & \leq \sum_{i=1}^n \sum_{j=1}^n |x_i - y_i| |d_{ij}| |\sigma| |f_j(x_j) - f_j(y_j)| \\ & \leq \sum_{i=1}^n \sum_{j=1}^n |x_i - y_i| |d_{ij}| |\sigma \ell_j| |x_i - y_i| \\ & = \begin{pmatrix} |x_1 - y_1| \\ |x_2 - y_2| \\ \vdots \\ |x_n - y_n| \end{pmatrix}^T D^* \begin{pmatrix} |x_1 - y_1| \\ |x_2 - y_2| \\ \vdots \\ |x_n - y_n| \end{pmatrix} \\ & \leq \lambda_{max}(D^*) \|x - y\|_2^2 \end{aligned} \tag{8}$$

Then, using (5)-(8) in (4) yields

$$\begin{aligned} & (x - y)^T (H(x) - H(y)) \leq -c_m \|x - y\|_2^2 \\ & + \ell_M \|Q\|_2 \|x - y\|_2^2 + \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} \|x - y\|_2^2 \\ & + \lambda_{max}(D^*) \|x - y\|_2^2 \\ & = -(c_m - \ell_M \|Q\|_2 - \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} \\ & - \lambda_{max}(D^*)) \|x - y\|_2^2 \\ & = -\varepsilon \|x - y\|_2^2. \end{aligned}$$

For $x \neq y$ and $\varepsilon > 0$, we get

$$(x - y)^T (H(x) - H(y)) < 0 \tag{9}$$

Thus, from which we can conclude that if $x \neq y$, then $H(x) \neq H(y)$.

Let $y = 0$ in (9), we have

$$x^T (H(x) - H(0)) \leq -\varepsilon \|x\|_2^2.$$

Taking the absolute value of both sides, we have

$$|x^T (H(x) - H(0))| \geq \varepsilon \|x\|_2^2.$$

Since,

$$|x^T (H(x) - H(0))| \leq \|x\|_\infty \|H(x) - H(0)\|_1.$$

$$\|x\|_\infty \|H(x) - H(0)\|_1 \geq \varepsilon \|x\|_2^2.$$

And, $\|x\|_\infty \leq \|x\|_2$, so

$$\|H(x) - H(0)\|_1 \geq \varepsilon \|x\|_2$$

Then, we have

$$\|H(x)\|_1 + \|H(0)\|_1 \geq \varepsilon \|x\|_2$$

$$\|H(x)\|_1 \geq \varepsilon \|x\|_2 - \|H(0)\|_1.$$

Since, $\|H(0)\|_1$ is finite, we can get the conclusion that if $\|x\| \rightarrow \infty$, that $\|H(x)\| \rightarrow \infty$. By Lemma 1, it is easy to know that $H(x)$ is homeomorphism of R^n . So, we have completed the proof of the existence and uniqueness of the equilibrium point for the neural networks defined by (1). ■

2. Stability of equilibrium point.

In the section 2, we have completed the proof of the existence and uniqueness of the equilibrium point for the networks defined by (1). Next, we begin to proof stability of equilibrium point for the system (1). We will first simplify system (1) as follows: we let $z_i(\cdot) = x_i(\cdot) - x_i^*$, $i = 1, 2, \dots, n$, and note that the $z_i(\cdot)$ are governed by:

$$\begin{aligned} \dot{z}_i(t) & = -c_i z_i(t) + \sum_{j=1}^n a_{ij} g_j(z_j(t)) \\ & + \sum_{j=1}^n b_{ij} g_j(z_j(t - \tau_{ij})) \\ & + \sum_{j=1}^n d_{ij} \int_{t-\sigma}^t g_j(z_j(s)) ds \\ & i = 1, 2, \dots, n. \end{aligned} \tag{10}$$

where $g_i(z_i(\cdot)) = f_i(z_i(\cdot) + x_i^*) - f_i(x_i^*)$, $i = 1, 2, \dots, n$, and $g_i \in \mathcal{L}$. We also note that

$$|g_i(z)| \leq \ell_i |z|, g_i(0) = 0, i = 1, 2, \dots, n.$$

Here, note that $z \rightarrow 0$ implies that $x \rightarrow x^*$. It is sufficient to prove the stability of the transformed system (10) instead of considering the stability of x^* of system (1).

Theorem 2. For the neural network defined by (1), assume that the network parameters satisfy (2) and $f \in \mathcal{L}$. Then, the neural network model (10) is globally asymptotically stably if the following condition holds:

$$\begin{aligned} \varepsilon & = c_m - \ell_M \|Q\|_2 - \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} \\ & - \lambda_{max}(D^*) > 0 \end{aligned}$$

where $c_m = \min(c_i)$, $\ell_M = \max(\ell_i)$, $\|Q\|_2 = \min\{\|A^*\|_2 + \|A_*\|_2, \sqrt{\|A^*\|_2^2 + \|A_*\|_2^2} + 2\|A_*^T\|_2\|A^*\|_2, \|\hat{A}\|_2\}$, with $A^* = \frac{1}{2}(\bar{A} + \underline{A})$, $A_* = \frac{1}{2}(\bar{A} - \underline{A})$, $\hat{A} = (\hat{a}_{ij})_{n \times n}$, $\hat{a}_{ij} = \max\{|a_{ij}|, |\bar{a}_{ij}|\}$, $\hat{B} = (\hat{b}_{ij})_{n \times n}$, with $\hat{b}_{ij} = \max(|b_{ij}|, |\bar{b}_{ij}|)$, $D^* = (d_{ij}^*)_{n \times n}$, $d_{ij}^* = \frac{|d_{ij}| \sigma \ell_j + |d_{ji}| \sigma \ell_i}{2}$, λ_i is eigenvalue of D^* , and $\lambda_{max} = \max_{1 \leq i \leq n} \{\lambda_i\}$, $i = 1, 2, \dots, n$.

Proof: We construct the follow positive definite Lyapunov functional:

$$\begin{aligned} V_1(z(t)) & = \frac{1}{2} \sum_{i=1}^n z_i^2(t) \\ & + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \ell_M |b_{ij}| \beta \int_{t-\tau_{ij}}^t z_j^2(s) ds \\ V_2(z(t)) & = \frac{1}{2\sigma} \int_{t-\sigma}^t \int_s^t |z(l)|^T D^* |z(l)| dlds \\ V(z(t)) & = V_1(z(t)) + V_2(z(t)) \end{aligned}$$

where β are positive constants to be determined later. The time derivative of the functional along the trajectories of

system (10) is obtained as follows:

$$\begin{aligned} \dot{V}_1(z(t)) = & -\sum_{i=1}^n c_i z_i^2(t) \\ & + \sum_{i=1}^n \sum_{j=1}^n a_{ij} z_i(t) g_j(z_j(t)) \\ & + \sum_{i=1}^n \sum_{j=1}^n b_{ij} z_i(t) g_j(z_j(t - \tau_{ij})) \\ & + \sum_{i=1}^n \sum_{j=1}^n d_{ij} z_i(t) \int_{t-\sigma}^t g_j(z_j(s)) ds \\ & + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \ell_M |b_{ij}| \beta (z_j^2(t) - z_j^2(t - \tau_{ij})) \end{aligned} \quad (11)$$

$$+ \left(\begin{array}{c} |z_1(s)| \\ |z_2(s)| \\ \vdots \\ |z_n(s)| \end{array} \right)^T D^* \left(\begin{array}{c} |z_1(s)| \\ |z_2(s)| \\ \vdots \\ |z_n(s)| \end{array} \right) ds$$

$$= \frac{1}{2} |z(t)|^T D^* |z(t)| + \frac{1}{2\sigma} \int_{t-\sigma}^t |z(s)|^T D^* |z(s)| ds$$

Then, we can write the following inequalities:

$$-\sum_{i=1}^n c_i z_i^2(t) \leq -\sum_{i=1}^n c_i z_i^2(t) \leq -\sum_{i=1}^n c_m z_i^2(t) = -c_m \|z(t)\|_2^2 \quad (12)$$

$$\dot{V}_2(z(t)) = \frac{1}{2} |z(t)|^T D^* |z(t)| - \frac{1}{2\sigma} \int_{t-\sigma}^t |z(s)|^T D^* |z(s)| ds \quad (17)$$

and

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^n a_{ij} z_i(t) g_j(z_j(t)) & = z^T(t) A g(z(t)) \\ & \leq \|z(t)\|_2 \|A\|_2 \|g(z(t))\|_2 \\ & \leq \ell_M \|A\|_2 \|z(t)\|_2^2 \\ & \leq \ell_M \|Q\|_2 \|z(t)\|_2^2 \end{aligned} \quad (13)$$

and

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^n b_{ij} z_i(t) g_j(z_j(t - \tau_{ij})) & \leq \sum_{i=1}^n \sum_{j=1}^n |b_{ij} z_i(t) g_j(z_j(t - \tau_{ij}))| \\ & \leq \sum_{i=1}^n \sum_{j=1}^n \ell_M |b_{ij}| |z_i(t)| |z_j(t - \tau_{ij})| \\ & \leq \frac{1}{2} \ell_M \sum_{i=1}^n \sum_{j=1}^n |b_{ij}| (\beta z_j^2(t - \tau_{ij}) + \frac{1}{\beta} z_i^2(t)) \end{aligned}$$

Thus, we can get

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^n b_{ij} z_i(t) g_j(z_j(t - \tau_{ij})) & + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \ell_M |b_{ij}| \beta (z_j^2(t) - z_j^2(t - \tau_{ij})) \\ & = \frac{1}{2} \ell_M \sum_{i=1}^n \sum_{j=1}^n (|b_{ij}| \beta z_j^2(t) + |b_{ij}| \frac{1}{\beta} z_i^2(t)) \\ & = \frac{1}{2} \ell_M \sum_{i=1}^n \sum_{j=1}^n (\frac{1}{\beta} |b_{ij}| + \beta |b_{ji}|) z_i^2(t) \end{aligned}$$

Let $\beta = \sqrt{\frac{\|B\|_\infty}{\|B\|_1}}$, we have

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^n b_{ij} z_i(t) g_j(z_j(t - \tau_{ij})) & + \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \ell_M |b_{ij}| \beta (z_j^2(t) - z_j^2(t - \tau_{ij})) \\ & = \frac{1}{2} \ell_M (\sqrt{\frac{\|B\|_\infty}{\|B\|_1}} \|B\|_1 + \sqrt{\frac{\|B\|_1}{\|B\|_\infty}} \|B\|_\infty) \|z(t)\|_2^2 \\ & = \ell_M \sqrt{\|B\|_1 \|B\|_\infty} \|z(t)\|_2^2 \\ & \leq \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} \|z(t)\|_2^2 \end{aligned} \quad (14)$$

then

$$\begin{aligned} \sum_{i=1}^n \sum_{j=1}^n d_{ij} z_i(t) \int_{t-\sigma}^t g_j(z_j(s)) ds & \leq \sum_{i=1}^n \sum_{j=1}^n \int_{t-\sigma}^t |z(t)| |d_{ij}| |l_j| |z_j(s)| ds \\ & = \int_{t-\sigma}^t \left(\begin{array}{c} |z_1(t)| \\ |z_2(t)| \\ \vdots \\ |z_n(t)| \end{array} \right)^T \frac{D^*}{\sigma} \left(\begin{array}{c} |z_1(s)| \\ |z_2(s)| \\ \vdots \\ |z_n(s)| \end{array} \right) ds \\ & \leq \int_{t-\sigma}^t \frac{1}{2\sigma} \left(\begin{array}{c} |z_1(t)| \\ |z_2(t)| \\ \vdots \\ |z_n(t)| \end{array} \right)^T D^* \left(\begin{array}{c} |z_1(t)| \\ |z_2(t)| \\ \vdots \\ |z_n(t)| \end{array} \right) ds \end{aligned} \quad (15)$$

Using (11)-(16), we have

$$\begin{aligned} \dot{V}(z(t)) & = \dot{V}_1(z(t)) + \dot{V}_2(z(t)) \\ & = -c_m \|z(t)\|_2^2 + \ell_M \|Q\|_2 \|z(t)\|_2^2 \\ & \quad + \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} \|z(t)\|_2^2 + |z(t)|^T D^* |z(t)| \\ & \leq -c_m \|z(t)\|_2^2 + \ell_M \|Q\|_2 \|z(t)\|_2^2 \\ & \quad + \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} \|z(t)\|_2^2 + \lambda_{max}(D^*) \|z(t)\|_2^2 \\ & = -(c_m - \ell_M \|Q\|_2 - \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} \\ & \quad - \lambda_{max}(D^*)) \|z(t)\|_2^2 \\ & = -\varepsilon \|z(t)\|_2^2 \end{aligned}$$

From which we can observe that $\dot{V}(z(t)) < 0$, for all $z(t) \neq 0$. Assume that $z(t) = 0$, we can get $\dot{V}(z(t)) < 0$ easily. If $z(t) = 0$ and $z_j(t - \tau_{ij}) = 0$ for all i, j implying that $\dot{V}(z(t)) = 0$, and $\dot{V}(z(t)) < 0$ otherwise. It is easy to prove that $V(z(t))$ is radially unbound since $V(z(t)) \rightarrow \infty$ as $\|z(t)\| \rightarrow \infty$. So, from Lyapunov theorems, we can conclude that the system (3) or equivalently the equilibrium point of system (1) is globally asymptotically stable. ■

IV. EXAMPLES

Example 1. According to the model (1), giving the the following matrices:

$$\underline{A} = \begin{bmatrix} -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 \end{bmatrix}, \bar{A} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix},$$

$$\bar{B} = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 1 & 2 & 2 & 4 \\ 1 & 2 & 2 & 4 \\ 1 & 2 & 2 & 4 \end{bmatrix}, \underline{B} = \begin{bmatrix} -1 & -2 & -2 & -4 \\ -1 & -2 & -2 & -4 \\ -1 & -2 & -2 & -4 \\ -1 & -2 & -2 & -4 \end{bmatrix},$$

$$D = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

$$\ell_1 = \ell_2 = \ell_3 = \ell_4 = 1, c_1 = c_2 = c_3 = c_4 = 1, \sigma = 1.$$

We can calculate the follow matrices:

$$A^* = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, A_* = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix},$$

$$\hat{A} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} \hat{B} = \begin{bmatrix} 1 & 2 & 2 & 4 \\ 1 & 2 & 2 & 4 \\ 1 & 2 & 2 & 4 \\ 1 & 2 & 2 & 4 \end{bmatrix},$$

$$D^* = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

from which we can directly calculate the norms: $\|A^*\|_2 + \|A_2\|_2 = 4$, $\sqrt{\|A^*\|_2^2 + \|A_*\|_2^2 + 2\|A_*^T | A^*\|_2} = 4$, $\|\hat{A}\|_2 = 4$. So $\|q\|_2 = 4$. We can also get that $\|\hat{B}\|_1 = 16$, $\|\hat{B}\|_\infty = 9$, $\lambda_{max}(D^*) = 4$.

Firstly, we apply the result of Theorem 1 to the neural network employing the network parameters of this example, we get

$$\varepsilon = c_m - \ell_M \|Q\|_2 - \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} - \lambda_{max}(D^*) = c_m - 20.$$

Thus, the constraint condition imposed on c_m by Theorem 1 for robust stability of system (1) is determined to be $c_m > 20$.

Example 2. According to the model (1), giving the the following matrices:

$$\underline{A} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 \\ 0 & -1 & 0 & -1 \\ -1 & -0 & -0 & -1 \end{bmatrix}, \bar{A} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix},$$

$$\underline{B} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} \bar{B} = \begin{bmatrix} -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 \\ -1 & -1 & -1 & -1 \end{bmatrix},$$

$$D = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

$$\ell_1 = \ell_2 = \ell_3 = \ell_4 = 1, \underline{c}_1 = \underline{c}_2 = \underline{c}_3 = \underline{c}_4 = 1, \sigma = 1.$$

We can calculate the follow matrices:

$$A^* = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ -1 & -1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ -1 & 1 & 1 & -1 \end{bmatrix}, A_* = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix},$$

$$\hat{A} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix} \hat{B} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix},$$

$$D^* = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

from which we can directly calculate the norms: $\|A^*\|_2 + \|A_2\|_2 = 3$, $\sqrt{\|A^*\|_2^2 + \|A_*\|_2^2 + 2\|A_*^T | A^*\|_2} = 3.6$, $\|\hat{A}\|_2 = 4$. So $\|q\|_2 = 3$. We can also get that $\|\hat{B}\|_1 = 16$, $\|\hat{B}\|_\infty = 4$, $\lambda_{max}(D^*) = 4$.

Firstly, we apply the result of Theorem 1 to the neural network employing the network parameters of this example, we get

$$\varepsilon = c_m - \ell_M \|Q\|_2 - \ell_M \sqrt{\|\hat{B}\|_1 \|\hat{B}\|_\infty} - \lambda_{max}(D^*) = c_m - 11 > 0$$

Hence, $c_m > 11$ proved to be a sufficient condition for robust stability of the neural network parameters of this example.

V. CONCLUSION

In this letter, a improved delay-dependent global robust exponential stability criterion for uncertain stochastic discrete-time neural networks with time-varying delay is proposed. A suitable Lyapunov functional has been proposed to derive some less conservative delay-dependent stability criteria by using the free-weighting matrices method and the convex combination theorem. Finally, two numerical examples have been given to demonstrate the effectiveness of the proposed method.

REFERENCES

- [1] O.Faydasicok and S.Arik, A new robust stability criterion for dynamical neural networks with multiple time delays, *Neurocomputing*(2012).In press.
- [2] J.L.Shao,T.Z.Huang,X.P.Wang,Improved global robust exponential stability criteria for interval neural networks with time-varying delays, *Expert Syst. Appl.* 38(12)(2011)15587C15593.
- [3] V.Singh,Global robust stability of delayed neural networks:estimating upper limit of norm of delayed connection weight matrix, *Chaos Solitons Fractals* 32(1)(2007)259C263.
- [4] T.Ensari,S.Arik,New results for robust stability of dynamical neural networks with discrete time delays, *Expert Syst.Appl.*27(8)(2010) 5925C5930.
- [5] F.Deng, M.Hua, X.Liu, Y.Peng, J.Fei, Robust delay-dependent exponential stability for uncertain stochastic neural networks with mixed delays, *Neurocomputing* 74(10)(2011)1503C1509.
- [6] T.Huang, Robust stability of delayed fuzzy CohenCGrossberg neural networks, *Comput.Math.Appl.*61(8)(2011)2247C2250.
- [7] Z.Wang, Y.Liu, X.Liu, Y.Shi, Robust state estimation for discrete-time stochastic neural networks with probabilistic measurement delays, *Neurocomputing* 74(1C3)(2010)256C264.
- [8] Q.B. Liu, G.L. Chen, On two inequalities for the Hadamard product and the Fan product of matrices, *Linear Algebra. Appl.*, 431(2009): 974-984.
- [9] X.Li, Global robust stability for stochastic interval neural networks with continuously distributed delays of neutral type, *Appl.Math.Comput.* 215 (12) (2001)4370C4384.
- [10] L.Liu, Z.Han, W.Li, Global stability analysis of interval neural networks with discrete and distributed delays of neutral type, *Expert Syst.Appl.*36(3) (2009) 7328C7331.
- [11] M.Luo, S.Zhong, R.Wang, W.Kang, Robust stability analysis for discrete-time stochastic neural networks systems with time-varying delays, *Appl. Math. Comput.*209(2)(2009)305C313.
- [12] X.Liu, N.Jiang, Robust stability analysis of generalized neural networks with multiple discrete delays and multiple distributed delays, *Neurocomputing*72 (7C9) (2009)1789C1796.
- [13] H.Li, B.Chen, Q.Zhou, S.Fang, Robust exponential stability for uncertain stochastic neural networks with discrete and distributed time-varying delays, *Phys.Lett.*A372(19)(2008)3385C3394.
- [14] H.Zhang, Z.Wang, D.Liu, Robust stability analysis for interval CohenC-Grossberg neural networks with unknown time-varying delays, *IEEETrans. Neural Networks*19(11)(2008)1942C1955.
- [15] Z.Wang, H.Zhang, W.Yu, Robust stability criteria for interval CohenC Grossberg neural networks with time varying delay, *Neurocomputing*72 (4C6) (2009)1105C1110.

Guang Zhou was born in Anhui Province, China, in 1987. He received the B.S. degree from Fuyang University in 2011. He is currently pursuing the M.S. degree from University of Electronic Science and Technology of China. His research interests include the stability of neural networks, pursuing the Ph.D. degree with UESTC. His research interests include chaos synchronization, switch and delay dynamic systems.

Shouming Zhong was born in 1955 in Sichuan, China. He received B.S. degree in applied mathematics from UESTC, Chengdu, China, in 1982. From 1984 to 1986, he studied at the Department of Mathematics in Sun Yat-sen University, Guangzhou, China. From 2005 to 2006, he was a visiting research associate with the Department of Mathematics in University of Waterloo, Waterloo, Canada. He is currently a full professor with School of Applied Mathematics, UESTC. His current research interests include differential equations, neural networks, biomathematics and robust control. He has authored more than 80 papers in reputed journals such as the International Journal of Systems Science, Applied Mathematics and Computation, Chaos, Solitons and Fractals, Dynamics of Continuous, Discrete and Impulsive Systems, Acta Automatica Sinica, Journal of Control Theory and Applications, Acta Electronica Sinica, Control and Decision, and Journal of Engineering Mathematics