

# Effect of Magnetic Field on Seed Germination of Two Wheat Cultivars

Ahmad Gholami<sup>1</sup>, Saeed Sharafi<sup>2</sup> and Hamid Abbasdokht<sup>1</sup>

**Abstract**—The effect of magnetic field on germination characteristics of two wheat Seeds has been studied under laboratory conditions. Seeds were magnetically exposed to magnetic field strengths, 125 or 250mT for different periods of time. Mean germination time and the time required to obtain 10, 25, 50, 75 and 90% of seeds to germinate were calculated. The germination time for each treatment were in general, higher than corresponding control values, in the other word in treated seeds time required for mean seed germination time increased nearly 3 hours in compared non treated control seeds.  $T_{10}$  for doses  $D_5$ ,  $D_6$ ,  $D_{11}$  and  $D_{12}$  significantly higher than the control values for both cultivars. Mean germination time (MGT) in both cultivars significantly increased when the time of seed exposed at magnetic field treatments increased, about 3 and 2 hour respectively for Omid and BCR cultivars.

**Keywords**—wheat, cultivar, germination test, magnetic field

## I. INTRODUCTION

WHEAT (*Triticum aestivum* L.) is one of the world's most important food crops. Seed vigor is defined as the sum total of those properties of the seed which determine the level of activity and performance of the seed or seed lot during germination and seedling emergence [8]. High seed vigor, i.e. rapid, uniform, complete emergence of vigorous seedling, leads to high grain yield of wheat by shortening the days from sowing to complete ground cover, allows the establishment of optimum canopy structure that maximizes crop yield.

Biological stimulation plays an important role in optimizing crops in terms of the maximization of yield, promotion of plant growth, and protection against disease [18]. The main advantage of using electromagnetic stimulation methods over traditional chemical processes is the absence of toxic residues [18]. Savostin [17] observed increase in the rate of wheat seedling elongation under magnetic conditions. Then, Murphy [14] reported changes in seed germination due to a magnetic field. Exposure of seeds to magnetic field for a short time was found to help in accelerated sprouting and growth of the seedlings [19]. They reported the enhancement of plant growth under magnetic conditions.

Aladjajiyan [2] detected that seed exposure to a 150 mT magnetic field stimulated shoot development and led to increase of the germination, fresh weight and shoot length of

maize plants. Studies indicated that suitable magnetic treatment increased the absorption and assimilation of nutrients, and ameliorated photosynthetic activities [10].

Physiological mechanisms of magnetic field on germination and seedling growth are not completely understood. Magnetic field treatment of seeds leads to acceleration of plants growth, proteins biosynthesis and root development [11]. Akoyunoglou [1] and Racuciu and et al. [16] reported that the activities of some enzymes were increased by exposure to magnetic field. Changes in amylase and nitrate reductase activities in germinating seeds treated by electromagnets of different field strengths were also observed [4].

The main objective of this work is to quantify the possible effects of magnetic treatment on the germination characteristics and heterotrophic growth of wheat seeds subjected to a stationary magnetic field.

## II. MATERIAL AND METHODS

### A. Seeds and magnetic treatment

Germination tests were carried out at laboratory conditions with two cultivars wheat seeds named *Omid* and *Backcross Roshan* (BCR). This experiment was conducted as completely randomized design with four replications. Each plot contained 25 seeds of each cultivar. Germination tests were performed according to the guidelines issued by the International Seed Testing Association [9]. Each filter paper with seeds was rolled and placed in a vessel containing distilled water. Four hours later, when seeds were soaked, each roll was subjected to a magnetic treatment.

The procedure of study was conducted according to Florez et al. [6], briefly magnetic treatment was provided as doses (D), varying the exposure time  $t$  and the magnetic field induction (B). the magnetic fields generated by ring magnets with magnetic induction values  $B_1 = 125\text{mT}$  and  $B_2 = 250\text{mT}$ ; the geometric characteristics are 7.5cm external diameter, 3cm internal diameter, 1cm high for  $B_1$  and 1.5cm high for  $B_2$ . The magnet was placed at the top of the vessel to generate each magnetic dose, and each roll containing 25 seeds was placed into hole of the magnet for the time established for each dose.

The doses  $D_1$ - $D_{12}$  applied were obtained by exposing the seeds to each magnetic field induction for different time, as follows: 1-Exposure to 125 mT:  $D_1$  (10min),  $D_2$  (20min),  $D_3$  (30min),  $D_4$  (1h),  $D_5$  (24h),  $D_6$  (continuous exposure). 2-Exposure to 250 mT:  $D_7$  (10min),  $D_8$  (20min),  $D_9$  (30min),  $D_{10}$  (1h),  $D_{11}$  (24h),  $D_{12}$  (continuous exposure). Four rolls carrying 25 seeds were used for each magnetic dose and an additional four rolls for control seeds (non exposed seeds). All the vessels containing rolls with seeds were labeled with

1- Shahrood University of Technology, Shahrood, IRAN  
phone:(+98 274 5224621); fax:(+98 274 5224620);  
e-mail: gholami@shahroodut.ac.ir).

2- Arak University, Arak, IRAN. (sharafi.saeed@gmail.com).

numbers and randomly located to carry out the test. The distance between any two vessels was at least 25 cm, to avoid the influence of each magnet on the other vessels around. Label numbers were not related to the magnetic dose applied, so that scores of germinated seeds and other measurements were blind [6].

### B. Germination test

The artificial light cycle was 12-h light/12-h darkness and the daily temperature  $20 \pm 2$  °C, night temperature  $18 \pm 2$  °C. The number of germinated seeds was recorded 4 times per day for the time necessary to achieve the final number or percentage of germinated seeds ( $G_{max}$ ). Seeds were observed daily for up to 7 days and considered germinated when the radicle was approximately 2 mm long or more. The rate of germination was assessed by determining the mean germinating time (MGT) and time required to germinate 10, 25, 50, 75 and 90% of seeds (parameters  $T_{10}$ ,  $T_{25}$ ,  $T_{50}$ ,  $T_{75}$  and  $T_{90}$ ).

### C. Statistical analyses

The results were subjected to an analysis of variance (ANOVA) to detect differences between mean parameters. Means were compared using with Duncan test to detect differences between parameters of treated plants and control.

## III. RESULTS AND DISCUSSION

### Seed germination

The percentage of germinated seeds ( $G_{max}$ ), time required for germination (parameters MGT,  $T_{10}$ - $T_{90}$ ) were determined for each treatment are presented in tables 1 and 2.

The germination time for each treatment were in general, higher than corresponding control values, in the other word in treated seeds time required for mean seed germination time increased nearly 3 hours in compared non treated control seeds. Thus the rate of germination of treated seeds was lower than the untreated seeds. The time needed to germinate 10%, of the seeds exposed to 125 mT for 10(D1), 20(D2), 30(D3) and 60(D4) minute and exposed to 250 mT for 10(D7), 20(D8), 30(D9) and 60(D10) minute were statistically similar to control (tables 1, 2). Results showed that time required for  $T_{10}$  for doses  $D_5$ ,  $D_6$ ,  $D_{11}$  and  $D_{12}$  were 23.25, 27, 35.25 and 31.50 respectively that significantly higher than the control values for Omid cultivar( table 1). Similar results obtained for BRC cultivar (table 2). As  $T_{10}$  in closely related to the onset of germination, these results indicate no response (for 8 treatments) and the delay of germination (for 4 treatments) of wheat seeds to magnetic field. Mean germination time (MGT) in both cultivars significantly increased when the time of seed exposed at magnetic field treatments increased , about 3 and 2 hour respectively for Omid and BCR cultivars.

Data measured on 7 days after seedlings allow us to corroborate the effect observed in the germination test. seeds final germination significantly affected by treatments (tables 1, 2).in Omid cultivar exposure of seed at magnetic field with  $D_3$ ,  $D_6$ ,  $D_{10}$ ,  $D_{11}$  and  $D_{12}$  caused seed germination significantly reduced but germination at  $D_2$  and  $D_4$

significantly increased. This reduction happened for BCR cultivar only at  $D_{12}$ .  $D_5$ ,  $D_9$  and  $D_{11}$  significantly increased seed germination. The stimulatory effect of the application of different magnetic doses on the germination is in agreement with that obtained by other researchers. Florez et al. [6] observed an increase for initial growth stages and an early sprouting of rice and maize seeds exposed to 125 and 250 mT stationary magnetic fields. Martinez et al. [12, 13] observed similar effects on wheat and barley seeds magnetically treated. Alexander and Doijode [3] reported that pregermination treatment improved the germination and seedling. Vigor of low viability rice and onion seeds. Kavi [10] found that seeds exposed to a magnetic field absorbed more moisture. Carbonell et al. [5] found that magnetic treatment produced a biostimulation of the germination.

The mechanisms at work when plant and other living systems are exposed to a magnetic field are not well known yet, but several theories have been proposed, including biochemical changes or altered enzyme activities by Pirke et al. [15]. Garcia and Arza [7] carried out an experimental study on water absorption by lettuce seeds previously treated in a stationary magnetic field of 1-10 mT; they reported an increase in water uptake rate due to the applied magnetic field, which may be the explanation for increase in the germination speed of treated.




Fig.1. (a) Magnet and (b) vessel containing distilled water. Roll of filter paper with seeds and the hollow cylindrical magnet. N, S: North and South poles of magnet (fig. from Florez et al., [6] ).

## REFERENCES

TABLE I. EFFECT OF MAGNETIC FIELD ON GERMINATION OF WHEAT SEEDS (OMID CULTIVAR)

Dose	T <sub>10</sub> (h)	T <sub>25</sub> (h)	T <sub>50</sub> (h)	T <sub>75</sub> (h)	T <sub>90</sub> (h)	MGT (h)	G <sub>max</sub> (%)
<i>B=125mT</i>							
D <sub>1</sub>	13.50 <sup>d</sup>	17.25 <sup>c</sup>	21.00 <sup>d</sup>	25.5 <sup>d</sup>	33.00 <sup>bc</sup>	22.05 <sup>c</sup>	96.25 <sup>b</sup>
D <sub>2</sub>	15.75 <sup>c</sup>	19.50 <sup>bc</sup>	24.00 <sup>c</sup>	30.75 <sup>c</sup>	36.00 <sup>b</sup>	25.20 <sup>bc</sup>	97.50 <sup>a</sup>
D <sub>3</sub>	15.00 <sup>c</sup>	18.00 <sup>bc</sup>	27.75 <sup>c</sup>	35.25 <sup>b</sup>	39.00 <sup>a</sup>	27.00 <sup>b</sup>	91.25 <sup>c</sup>
D <sub>4</sub>	17.25 <sup>c</sup>	20.25 <sup>b</sup>	25.50 <sup>c</sup>	31.5 <sup>c</sup>	34.50 <sup>b</sup>	25.80 <sup>bc</sup>	98.75 <sup>a</sup>
D <sub>5</sub>	23.25 <sup>b</sup>	29.25 <sup>a</sup>	33.00 <sup>b</sup>	36.75 <sup>b</sup>	40.50 <sup>a</sup>	32.55 <sup>a</sup>	96.25 <sup>b</sup>
D <sub>6</sub>	27.00 <sup>a</sup>	31.50 <sup>a</sup>	36.00 <sup>a</sup>	40.5 <sup>a</sup>	-	27.00 <sup>b</sup>	81.25 <sup>c</sup>
Mean	18.63	22.63	27.88	33.38	36.6	26.6	
<i>B=250mT</i>							
D <sub>7</sub>	12.00 <sup>d</sup>	15.75 <sup>c</sup>	18.75 <sup>c</sup>	22.50 <sup>d</sup>	30.00 <sup>e</sup>	19.80 <sup>c</sup>	96.25 <sup>b</sup>
D <sub>8</sub>	14.25 <sup>d</sup>	18.00 <sup>c</sup>	21.00 <sup>c</sup>	28.50 <sup>c</sup>	31.50 <sup>bc</sup>	22.65 <sup>c</sup>	95.00 <sup>b</sup>
D <sub>9</sub>	14.25 <sup>d</sup>	16.50 <sup>c</sup>	18.75 <sup>c</sup>	24.00 <sup>d</sup>	28.50 <sup>e</sup>	20.40 <sup>c</sup>	96.25 <sup>b</sup>
D <sub>10</sub>	20.25 <sup>c</sup>	25.50 <sup>b</sup>	29.25 <sup>b</sup>	32.25 <sup>b</sup>	35.25 <sup>b</sup>	28.50 <sup>b</sup>	92.50 <sup>c</sup>
D <sub>11</sub>	35.25 <sup>a</sup>	38.25 <sup>a</sup>	41.25 <sup>a</sup>	45.00 <sup>a</sup>	48.75 <sup>a</sup>	41.70 <sup>a</sup>	91.25 <sup>c</sup>
D <sub>12</sub>	31.50 <sup>b</sup>	-	-	-	-	-	-
Mean	20.75	22.8	25.8	30.45	34.2	26.61	
Control	14.25 <sup>cd</sup>	17.25 <sup>c</sup>	24.00 <sup>c</sup>	27.75 <sup>d</sup>	33.00 <sup>bc</sup>	23.25 <sup>c</sup>	96.25 <sup>b</sup>

TABLE II EFFECT OF MAGNETIC FIELD ON GERMINATION OF WHEAT SEEDS (BCR CULTIVAR)

Dose	T <sub>10</sub> (h)	T <sub>25</sub> (h)	T <sub>50</sub> (h)	T <sub>75</sub> (h)	T <sub>90</sub> (h)	MGT (h)	G <sub>max</sub> (%)
<i>B=125mT</i>							
D <sub>1</sub>	7.00 <sup>bc</sup>	10.00 <sup>c</sup>	10.75 <sup>c</sup>	16.75 <sup>b</sup>	21.25 <sup>a</sup>	13.15 <sup>b</sup>	93.75 <sup>ab</sup>
D <sub>2</sub>	7.00 <sup>bc</sup>	10.00 <sup>c</sup>	11.50 <sup>c</sup>	15.25 <sup>b</sup>	19.75 <sup>ab</sup>	12.70 <sup>b</sup>	88.75 <sup>bc</sup>
D <sub>3</sub>	8.50 <sup>b</sup>	10.75 <sup>bc</sup>	13.00 <sup>bc</sup>	16.00 <sup>b</sup>	18.00 <sup>b</sup>	13.25 <sup>b</sup>	90.00 <sup>b</sup>
D <sub>4</sub>	8.50 <sup>b</sup>	13.00 <sup>b</sup>	16.00 <sup>b</sup>	19.00 <sup>a</sup>	23.50 <sup>a</sup>	16.00 <sup>a</sup>	92.50 <sup>b</sup>
D <sub>5</sub>	10.75 <sup>b</sup>	13.75 <sup>b</sup>	16.75 <sup>b</sup>	21.25 <sup>a</sup>	25.00 <sup>a</sup>	17.50 <sup>a</sup>	97.50 <sup>a</sup>
D <sub>6</sub>	15.25 <sup>a</sup>	21.25 <sup>a</sup>	24.25 <sup>a</sup>	-	-	-	
Mean	9.5	13.13	15.38	17.65	21.5	14.52	
<i>B=250mT</i>							
D <sub>7</sub>	5.00 <sup>c</sup>	8.00 <sup>c</sup>	9.50 <sup>b</sup>	12.50 <sup>c</sup>	16.25 <sup>c</sup>	10.25 <sup>c</sup>	91.25 <sup>b</sup>
D <sub>8</sub>	5.00 <sup>c</sup>	7.25 <sup>c</sup>	8.00 <sup>b</sup>	13.25 <sup>c</sup>	17.75 <sup>c</sup>	10.25 <sup>c</sup>	93.75 <sup>ab</sup>
D <sub>9</sub>	5.00 <sup>c</sup>	5.00 <sup>c</sup>	8.00 <sup>b</sup>	12.50 <sup>c</sup>	18.50 <sup>c</sup>	9.80 <sup>c</sup>	96.25 <sup>a</sup>
D <sub>10</sub>	14.00 <sup>b</sup>	17.00 <sup>b</sup>	20.00 <sup>b</sup>	18.00 <sup>b</sup>	23.75 <sup>b</sup>	18.55 <sup>b</sup>	82.50 <sup>c</sup>
D <sub>11</sub>	17.75 <sup>a</sup>	20.75 <sup>a</sup>	23.75 <sup>a</sup>	26.75 <sup>a</sup>	29.00 <sup>a</sup>	23.60 <sup>a</sup>	95.00 <sup>a</sup>
D <sub>12</sub>	17.00 <sup>a</sup>	22.25 <sup>a</sup>	-	-	-	-	
Mean	10.63	13.38	13.83	16.6	21.05	14.49	
Control	7.00 <sup>bc</sup>	10.00 <sup>c</sup>	12.25 <sup>c</sup>	16.00 <sup>b</sup>	19.00 <sup>ab</sup>	12.15 <sup>b</sup>	92.50 <sup>b</sup>

- [1] Akoyunoglou, G. 1964. Effect of a magnetic field on carboxydimutase. Nature 4931, 452-454.
- [2] Aladjadjiyan, A. 2002. Study of influence of magnetic field on some biological characteristics of *zea mays*. J.Central Eur. Agric. 3(2): 89-94.
- [3] Alexander, M.P. and S. D. Doijode. 1995. Electromagnetic field, a novel tool to increase germination and seedling vigor of conserved onion (*Allium cepa* L.) and rice (*Oryza sativa* L.) seeds with low viability. Plant Genet. Resource. Newslett. 104: 1-5.
- [4] Bathnagar, D. and A. Deb. 1977. Some aspects of pre-germination exposure of wheat seeds to magnetic field I. Germination and early growth. Seeds Res. 5: 129-137.
- [5] Carbonell, M.V., E. Martinez. and J.M. Amaya. 2000. Stimulation of germination in rice (*Oryza sativa* L.) by a static magnetic field. Electromagnetobiol. 19(1): 121-128.
- [6] Fl'orez, M., M. V. Carbonell, E. Martinez. 2007. Exposure of maize seeds to stationary magnetic fields: Effects on germination and early growth. Environmental and Experimental Botany 59 : 68-75
- [7] Garcia, F. and L.I. Arza. 2001. Influence of stationary magnetic field on water relations in lettuce seeds. Part I: theoretical considerations. Bioelectromagnetics 22(8): 589-595.
- [8] Hampton, J.G. and D.M. TeKrony. 1995. Handbook of vigor test methods, 3rd ed. International Seed Testing Association, Zurich, Switzerland, 117 pp
- [9] ISTA, International Rules for Seed Testing, 2004. International Seed Testing Assoc. Zurich, Switzerland
- [10] Kavi, P.S. 1977. The effect of magnetic treatment of soybean seed on its moisture absorbing capacity. Sci. Cult. 43, 405-406.
- [11] Kordas, L. 2002. The Effect of Magnetic Field on Growth, Development and the Yield of Spring Wheat, Polish Journal of Environmental Studies 11(5): 527-530
- [12] Martinez, E., M. V. Carbonell. and M. Florez. 2002. Magnetic biostimulation of initial growth stages of wheat (*Triticum aestivum*. L). Electromagnetobiol. Med. 21(1): 43-53.
- [13] Martinez, E., M. V. Carbonell and J. M Amaya. 2000. A static magnetic field of 125 mT stimulates the initial growth stages of barley (*Hordeum vulgare* L.). Electromagnetobiol. 19(3): 271-277.
- [14] Murphy, J.D. 1942. The influence of magnetic field on seed germination. Am. J. Bot. 29 (Suppl.), 15.
- [15] Phirke, P. S., A. B Kubde. and S. P. Umbakar. 1996. The influence of magnetic field on plant growth. Seed Sci. Technol. 24: 375-392.
- [16] Racuciui, M., D. Creanga and I. Horga. 2008. Plant growth under static magnetic field influence. Rom. Journal. Phys. 53(1-2):353-359,
- [17] Savostin, P.W., 1930. Magnetic growth relations in plant. Planta 12, 327.
- [18] Slepser, D., M. S. Pathan, B. Camps-Raga, N. Boriraksantikul, S. Tantong, S. R. Gyawali, P. Kirawanich, and N. E. Islam. Experimental Analysis of Corn Seed Germination Enhancement under the Application of Electromagnetic and Magnetic Fields neos.fontismedia.com/euro\_em/files/e288.pdf
- [19] Yinan, L., Yuan, L., Yongqing, Y., Chunyang, L., 2005. Effect of seed pretreatment by magnetic field on the sensitivity of cucumber (*Cucumis sativus*) seedlings to ultraviolet-B radiation. Environ. Exp. Botany 54, 286- 294