

# Large Deviations for Lacunary Systems

Bainian Li, Kongsheng Zhang

**Abstract**—Let  $X_i$  be a Lacunary System, we established large deviations inequality for Lacunary System. Furthermore, we gained Marcinkiewicz Larger Number Law with dependent random variables sequences.

**Keywords**—Lacunary system, larger deviations, Locally Generalized Gaussian, Strong law of large numbers.

## I. INTRODUCTION

**L**ACUNARY systems is a class of random variables. Lai and Wei[1] gave independent and identically distributed random variable, martingale differences with  $L_p$  bound are Lacunary Systems. Li [2] obtained  $L_p$  bounded dependent is a Lacunary System in 1997.

In this paper, we shall establish large deviations inequality for Lacunary System. Further, we shall get Marcinkiewicz Strong law of large numbers with  $m$ -dependent random variables sequences.

We give defined of Lacunary system as follows:

**Definition 1.1** Given  $p > 0$ , a sequence of real-valued random variables  $\{X_n, n \geq 1\}$  is called a Lacunary System or an  $S_p$  system, if there exists a positive constant  $K_p$  such that

$$E|\sum_{i=m}^n C_i X_i|^p \leq K_p (\sum_{i=m}^n C_i^2)^{p/2} \quad (1)$$

for any sequence of real constant  $\{C_i\}$  and all  $n \geq m$ .

**Definition 1.2** Suppose that  $\{X_n, n \geq 1\}$  is a sequence of random variables on a probability space  $(\Omega, \mathcal{F}, P)$ , set  $\mathcal{F}_a^b = \sigma(X_k, a \leq k \leq b)$ . Denote by the  $\sigma$ -field generated by the random variables  $X_a, X_{a+1}, \dots, X_b$ .

1) Let  $A \in \mathcal{F}_1^k, B \in \mathcal{F}_{k+n}^\infty$  and  $k, n \geq 1, \{X_n, n \geq 1\}$  is called  $\phi$ -mixing if

$$|P(A \cap B) - P(A)P(B)| \leq \phi(n)P(A)$$

for some  $\phi(n) \downarrow 0$ .

2)  $\{X_n, n \geq 1\}$  is called  $\psi$ -mixing, if

$$\psi(n) = \sup_{k \in \mathbf{N}} \psi(\mathcal{F}_1^k, \mathcal{F}_{k+n}^\infty) \rightarrow 0, n \rightarrow \infty,$$

where

$$\psi(\mathcal{A}, \mathcal{B}) = \sup_{A \in \mathcal{A}, B \in \mathcal{B}, P(A)P(B) > 0} \frac{|P(A \cap B) - P(A)P(B)|}{P(A)P(B)}.$$

**Definition 1.3** Let  $X$  be a real-valued random variable, we call a Locally Generalized Gaussian, If there exists  $\alpha > 0$  such that

$$E(\exp(ux)|\mathcal{F}) \leq \exp(u^2 \alpha^2 / 2) \quad a.s. \quad (2)$$

for any  $u \in R$ .

Bainian Li and Kongsheng Zhang are with the School of Statistics and Applied Mathematics, Anhui University of Finance and Economics, Bengbu, 233030 PR China, emails:libainian49@163.com; zks155@163.com

## II. LARGER DEVIATIONS INEQUALITY

In order to prove larger deviations we need the following lemmas.

**Lemma 2.1** Let  $X_n$  be a zero-mean  $\phi$ -mixing and  $\sum_{i=1}^{\infty} \phi^{1/2}(i) < \infty$ , for some  $p \geq 2$ ,  $\sup_i E|X_i|^p < \infty$ . There exists constant  $c > 0$  depending only on  $p$  for any real-valued sequence  $\{a_{ni}\}$ , such that

$$E|\sum_{i=1}^n a_{ni} X_i|^p \leq c (\sum_{i=1}^n a_{ni}^2)^{p/2}. \quad (3)$$

**proof** Let  $a_{ni} = 0, i > n$ , since  $\sum_{i=1}^{\infty} \phi^{1/2}(i) < \infty$ ,  $\sup_i E|X_i|^p < \infty$ , from the proof in [3], we have

$$E|\sum_{i=k+1}^{k+m} a_{ni} X_i|^2 \leq c_1 \sum_{i=k+1}^{k+m} a_{ni}^2 \leq c_1 \sum_{i=1}^n a_{ni}^2,$$

for any  $k \geq 0, n \geq 1, m \leq n$ .

Using the corollary 2.1 in [4], we obtain

$$\begin{aligned} E|\sum_{i=1}^n a_{ni} X_i|^p &\leq c_2 (\sum_{i=1}^n E|a_{ni} X_i|^p + (\sum_{i=1}^n a_{ni}^2)^{p/2}) \\ &\leq c_3 (\sum_{i=1}^n |a_{ni}|^p + (\sum_{i=1}^n a_{ni}^2)^{p/2}). \end{aligned} \quad (4)$$

Since  $p \geq 2$ , it follows that

$$(\sum_{i=1}^n |a_{ni}|^p)^{1/p} \leq (\sum_{i=1}^n |a_{ni}|^2)^{1/2} \Leftrightarrow \sum_{i=1}^n |a_{ni}|^p \leq (\sum_{i=1}^n |a_{ni}|^2)^{p/2}.$$

Then, we deduce (3) from (4).

**Remark 1** Lemma 2.1 implies that  $\phi$ -mixing is a Lacunary System. If  $a_{ni} \equiv 1$ , we have  $E|\sum_{i=1}^n X_i|^p \leq cn^{p/2}$ .

**Lemma 2.2** If  $\{X_n, n \geq 1\}$  is a zero-mean  $\psi$ -mixing, such that

$$\sum_{i=1}^{\infty} \psi(i) < \infty, E|X_i|^p, p \geq 2,$$

then for any real-valued sequence  $a_{ni}$ , (3) holds.

**Proof** From lemma 2.1 and the proof in [5], we can obtain lemma 2.2.

**Theorem 2.1** Let  $\{X_n, n \geq 1\}$  be a Lacunary System, for any  $p > 1, x > 0$ , sequence of real constant  $\{C_i\}$ , then

$$P\{|S_n| \geq nx\} \leq C(p) (\sum_{i=1}^n C_i^2)^{p/2} n^{-p}, \quad (5)$$

where  $S_n = \sum_{i=1}^n C_i X_i$ ,  $C_p = K_p / x^p$ .

**Proof** Since  $\{X_n, n \geq 1\}$  is a Lacunary System, we have

$$E|S_n|^p \leq K_p \left( \sum_{i=1}^n C_i^2 \right)^{p/2}.$$

By using Markov's inequality,

$$P\{|S_n| \geq nx\} \leq \frac{E|S_n|^p}{(nx)^p}$$

for every  $p > 1$ , we can obtain (5).

**Remark 2** (1) If  $\sum_{i=1}^n C_i^2 = O(n)$ , we have

$$E|S_n|^p \leq C(p)n^{-p/2}.$$

(2) If  $C_i \equiv 1, p > 2$ , by Borel-Cantelli lemma:

$$\sum_{n=1}^{\infty} P\left(\left|\sum_{i=1}^n X_i\right| \geq nx\right) \leq \sum_{n=1}^{\infty} C(p)n^{-p/2} < \infty,$$

then

$$\lim_{n \rightarrow \infty} P\left(\left|\sum_{i=1}^n X_i\right| \geq nx\right) = 0. \quad a.s.,$$

**Theorem 2.2** Let  $(X_n, \mathcal{F}_n)$  be a Locally Generalized Gaussian sequence, if  $\sup_n X_n = k < \infty$ , then (5) holds for any  $p \geq 2, x \geq 0$ .

**Proof** Theorem 2.2 holds if only we can prove that Locally Generalized Gaussian sequence is a Lacunary System. Let  $A_n = \sum_{i=m}^n C_i^2, u = x/k^2 A_n$ , by lemma 1 in [6], then

$$\begin{aligned} E(\exp(u \sum_{i=m}^n C_i X_i)) &= E(\exp(u(S_n - S_{m-1}))) \\ &\leq \exp(u^2 k^2 A_n / 2), \end{aligned} \quad (6)$$

where  $S_n = \sum_{i=1}^n C_i X_i$ . Since

$$P(\{|S_n - S_{m-1}| > x\}) \leq 2 \exp(-x^2 / 2k^2 A_n)$$

for  $p \geq 2$ , by Chebyshev's inequality, we get

$$\begin{aligned} E\left|\sum_{i=m}^n C_i X_i\right|^p &= p \int_0^\infty x^{p-1} P(|S_n - S_{m-1}| > x) dx \\ &\leq 2p \int_0^\infty x^{p-1} \exp(-x^2 / 2k^2 A_n) dx \\ &= 2^{p/2} p k^p A_n^{p/2} \int_0^\infty x^{p/2-1} e^{-x} dx \\ &= K_p \left( \sum_{i=m}^n C_i^2 \right)^{p/2}. \end{aligned}$$

where  $K_p = p 2^{p/2} k^p \int_0^\infty x^{p/2-1} e^{-x} dx$ .

### III. THE STRONG LAW OF LARGER NUMBERS

**Theorem 3.1** Assume that  $\{X_n, n \geq 1\}$  is a zero-mean  $\psi$ -mixing, such that

$$\sum_{i=1}^{\infty} \psi(i) < \infty, \quad E|X_i|^p, \text{ for } p \geq 2.$$

If there exists  $1/2 < r \leq 1, \theta = 2r - 1$  and positive constant  $K$  such that  $\sum_{i=1}^n a_{ni}^2 \leq K n^\theta, i = 1, 2, \dots, n$ , then

$$\frac{\sum_{i=1}^n a_{ni} X_i}{n^r} \rightarrow 0, \quad a.s.. \quad (7)$$

**Proof** Denote  $\sum_{i=1}^n a_{ni} X_i$ , by Markov's inequality, we have

$$P(|S_n| \geq n^r \varepsilon) \leq \frac{E(|S_n|^p)}{\varepsilon^p n^{pr}}. \quad (8)$$

From lemma 1.2 and (8), we obtain

$$\begin{aligned} \sum_{n=1}^{\infty} P\left(\left|\sum_{i=1}^n a_{ni} X_i / n^r\right| \geq \varepsilon\right) &= \sum_{n=1}^{\infty} P(|S_n| \geq \varepsilon n^r) \\ &\leq \sum_{n=1}^{\infty} \frac{E(|S_n|^p)}{\varepsilon^p n^{pr}} \leq \sum_{n=1}^{\infty} \frac{c \left(\sum_{i=1}^n a_{ni}^2\right)^{p/2}}{\varepsilon^p n^{pr}} \\ &\leq \sum_{n=1}^{\infty} \frac{c K n^{p\theta/2}}{\varepsilon^p n^{pr}} < \infty. \end{aligned}$$

(3.1) follows from Borel-Cantelli lemma.

**Remark 3.** This result extends independent and identically distributed Marcinkiewicz Law of large numbers for  $\psi$ -mixing.

**Theorem 3.2** Let  $\{X_n\}$  be a zero-mean  $\phi$ -mixing, and  $\sum_{i=1}^{\infty} \phi^{1/2}(i) < \infty, \sup_i E|X_i|^p < \infty$  for some  $p > 2$ . If there exists  $1/2 < r \leq 1, \theta = 1 - 2/p$  and positive constant  $K$  such that  $\sum_{i=1}^n a_{ni}^2 \leq K n^\theta, i = 1, 2, \dots, n$ , then

$$\frac{\sum_{i=1}^n a_{ni} X_i}{\sqrt{n \ln n}} \rightarrow 0, \quad a.s.. \quad (9)$$

**Proof** By lemma 2.1 and (8), we obtain

$$\begin{aligned} \sum_{n=1}^{\infty} P\left(\left|\sum_{i=1}^n a_{ni} X_i / \sqrt{n \ln n}\right| \geq \varepsilon\right) &= \sum_{n=1}^{\infty} P(|S_n| \geq \varepsilon \sqrt{n \ln n}) \\ &\leq \sum_{n=1}^{\infty} \frac{E(|S_n|^p)}{\varepsilon^p n^{p/2} (\ln n)^{p/2}} \\ &\leq \sum_{n=1}^{\infty} \frac{c \left(\sum_{i=1}^n a_{ni}^2\right)^{p/2}}{\varepsilon^p n^{p/2} (\ln n)^{p/2}} \\ &\leq \sum_{n=1}^{\infty} \frac{c K n^{p\theta/2}}{\varepsilon^p n^{p/2} (\ln n)^{p/2}} \\ &= \sum_{n=1}^{\infty} \frac{c K}{\varepsilon^p n (\ln n)^{p/2}} < \infty. \end{aligned}$$

And then, (9) follows from Borel-Cantelli lemma.

### ACKNOWLEDGMENT

This work was supported by the Science Foundation of the Anhui Province (090416222; KJ2010B001), and the Project of PR China for Statistical Research (2008LY048).

### REFERENCES

- [1] T. L. Lai, C. Z. Wei, *Lacunary systems and generalized linear processes*, Stochastic Processes Appl. 1983,14, 187-199.
- [2] B. N. Li, *Sp systems and the nonparametric regression on modal*, 1997,12(3): 265-269.

- [3] P. Billingsley, *Convergence of Probability Measures*. New York: John Wiley & Sons, Inc, 1968.
- [4] Q. M. Shao, *A moment inequality and its applications*, Acta Mathematica Sinica, 1988,31(6): 737 747.
- [5] S. C. Yang, *Almost sure convergence of weighted sums of mixing sequences*, System Science and Math. Sci., 1995,15(3):245 265.
- [6] S. H. Hu, *A large deviation result for the least squares estimators in nonlinear regression*, Stochastic processes and their applications, 1993,47(2): 345-352.