

Trend Analysis for Extreme Rainfall Events in New South Wales, Australia

Evan Hajani, Ataur Rahman, Khaled Haddad

Abstract—Climate change will affect the hydrological cycle in many different ways such as increase in evaporation and rainfalls. There have been growing interests among researchers to identify the nature of trends in historical rainfall data in many different parts of the world. This paper examines the trends in annual maximum rainfall data from 30 stations in New South Wales, Australia by using two non-parametric tests, Mann-Kendall (MK) and Spearman's Rho (SR). Rainfall data were analyzed for fifteen different durations ranging from 6 min to 3 days. It is found that the sub-hourly durations (6, 12, 18, 24, 30 and 48 minutes) show statistically significant positive (upward) trends whereas longer duration (sub-daily and daily) events generally show a statistically significant negative (downward) trend. It is also found that the MK test and SR test provide notably different results for some rainfall event durations considered in this study. Since shorter duration sub-hourly rainfall events show positive trends at many stations, the design rainfall data based on stationary frequency analysis for these durations need to be adjusted to account for the impact of climate change. These shorter durations are more relevant to many urban development projects based on smaller catchments having a much shorter response time.

Keywords—Climate change, Mann-Kendall test, Spearman's Rho test, trends, design rainfall.

I. INTRODUCTION

TREND detection is an active area of interest for both hydrologists and climatologists in order to investigate climate change scenarios and enhance climate impact research to identify impacts of climate change on hydro-meteorological time series data [1]. Trend detection in rainfall time series is crucial for planning and designing many regional water resources management projects. The trend in rainfall pattern has been examined extensively using different methods by many researchers from various parts of the world (e.g. Ashley et al. [2] and Haylock et al. [3] in USA, Burgueno et al. [4] in Spain, Odekunle et al. [5] in Africa, Zhai et al. [6] in China, Fujibe [7] in Japan, Hannaford and Buys [8] in UK, , Nikhil Raj and Azeez [9] in India, Hanif et al. [10] in Pakistan and Burn and Taleghani [11] in Canada). For example, Hamdi et al. [12] adopted several statistical approaches including Mann-Kendall, Linear Regression, CUSUM, Rank Sum, Student's t-test, Rank Difference, Auto Correlation and Skewness-

Kurtosis Normality test to detect trends and quantify evidences of climatic change by analyzing data from six meteorological stations around Jordan. The results indicated that there were no visible trends indicating an increase or decrease in the annual precipitation and maximum temperature. Karpouzou et al. [1] analysed trends of precipitation data covering a 30 year period for six stations located within the prefecture of Pieria in Northern Greece. Various statistical techniques such as the Mann-Kendall test, Sequential version of the Mann-Kendall test and Sen's estimator of slope were adopted to detect possible changes in precipitation in annual, monthly and seasonal basis. Furthermore, CUSUM test was used for evaluating a step change in the precipitation data. Results showed that more significant trends were related to seasonal time series than to annual ones.

In another study, Clarke et al. [13] investigated trends of 5 minutes to 24 hours duration rainfall data from 13 Canadian stations and found that about 30% of the cases demonstrated a statistically significant trend. Rana et al. [14] analyzed trends in precipitation of accumulated daily rainfall data of 48 stations covering 60 years of data during the period from 1951 to 2004 in the cities of Delhi and Mumbai in India. The Mann-Kendall test and linear regression were adopted to evaluate long term trends in rainfall. The analysis revealed great degree of variability in precipitation in the study area. Also it was found significant increasing trends in the higher latitudes and decreasing trends along coastal areas for seasonal rainfall. Mondal et al. [15] analysed the trends using daily rainfall data of 40 years from 1971 to 2010 to find out the monthly variability of rainfall in Birupa River basin located in the north-eastern part of Cuttack district in Orissa state of India. The modified Mann-Kendall test was used for the determination of the spatial variation and temporal trends. The results showed that there were increasing and decreasing trends in monthly data in the study area.

In Australia, many researchers have investigated trends in rainfall data (e.g. Yu and Neil [16], Plummer et al. [17], Haylock and Nicholls [18], Groisman et al. [19], Gallant et al. [20], Alexander et al. [21], Hardwick Jones et al. [22], Li et al. [23], and Chen et al. [24]). Most of these studies have found trends in Australian rainfall data, the common findings are that the northwest had experienced an increase in rainfall over the last 50 years, while much of eastern Australia and the far southwest have experienced a decrease. For example, Chowdhury and Beecham [25] investigated the monthly rainfall trends at ten rainfall stations across Australia covering all state capitals. They found a decreasing trend of June and

E. Hajani is a PhD student in Civil Engineering, School of Computing, Engineering and Mathematics, University of Western Sydney, Australia (phone: 61-2-47360139; e-mail: evan.hajani@gmail.com).

A. Rahman is an Associate Professor in Water Engineering in the School of Computing, Engineering and Mathematics, University of Western Sydney, Australia (phone: 61-2-47360145; e-mail: a.rahman@uws.edu.au).

K. Haddad is a Research Fellow in the School of Computing, Engineering and Mathematics, University of Western Sydney, Australia (phone: 61-2-47360920; e-mail: k.haddad@uws.edu.au).

July rainfall depth at two stations (Perth Airport and Sydney Observatory Hill), while no significant trends in Melbourne, Alice Springs and Townsville rainfall stations, whereas five stations showed an increasing trend on monthly rainfall depth. In another study, Jacob et al. [26] evaluated changes in sub-daily extreme rainfalls, and found that changes in short duration rainfall data are in greater magnitude than the longer duration ones. Chen et al. [24] suggested that for six minutes duration, rainfall intensity would increase despite a decrease in the mean annual rainfall.

Loweridge and Rahman [27] adopted four tests (Mann-Kendall, linear regression, CUSUM and cumulative deviation) to investigate trends in loss parameters using an event-based rainfall-runoff model for four catchments. The results showed that there was strong evidence of an upward trend in the initial loss data for two catchments. Also, Yilmaz and Perera [28] investigated the heavy rainfall trends for storm durations of 6 minutes to 72 hours. The statistical tests used were the Mann-Kendall and Spearman's Rho tests to detect trends in annual maximum rainfall intensities of the selected storm durations. The results of the data sets of hourly storm indicated increasing trends, but for 6 to 72 hours heavy rainfall data sets showed statistically insignificant decreasing trends.

The objective of this study is to investigate trends in sub-hourly, sub-daily and daily rainfall events in New South Wales (NSW), Australia using the latest rainfall data, to compare trends in shorter and longer duration events using two different non-parametric trend tests and the spatial variability of the identified trends.

II. METHOD AND DATA

There are many non-parametric and parametric trend tests available that can be adopted to investigate trends in rainfall data. Non-parametric trend tests were adopted in this study since this is robust with respect to non-normality, nonlinearity, missing values, serial dependency and outliers in the data [16]. The most frequently used non-parametric test for identifying trends in hydrological time series data is the Mann-Kendall (MK) test ([29], [30]), which is a rank-based distribution-free method for identifying trends. In this study, in addition to the MK test, Spearman's Rho (SR) test was also adopted so that the results can be compared by two different tests.

The null hypothesis in the MK test states that the data (X_1, X_2, \dots, X_n) are a sample of n independent and identically distributed random variables. The MK test statistic is given by:

$$S = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{sign}(X_j - X_i) \quad (1)$$

where X represents a univariate time-series, i and j denote the time indices associated with individual values, n is the number of data points and sign is determined as follows:

$$\text{sign}(X_j - X_i) = \begin{cases} +1 & (X_j - X_i) > 0 \\ 0 & \text{if } (X_j - X_i) = 0 \\ -1 & (X_j - X_i) < 0 \end{cases} \quad (2)$$

As documented in Mann [29] and Kendall [30], the statistic S under the null hypothesis is approximately normally distributed for $n \geq 8$ with mean and variance as follows:

$$E(S) = 0 \quad (3)$$

$$\text{Var}(S) = \frac{n(n-1)(2n+5) - \sum_{l=1}^L t_l(l-1)(2l+5)}{18} = \sigma^2 \quad (4)$$

where t_l indicates the number of ties of extent l , and L is the number of tied groups. Under the null hypothesis, the standardized test statistic (Z) defined in (5) and its corresponding p-value are approximately normally distributed as defined below:

$$Z_S = \begin{cases} \frac{S-1}{\sigma} & \text{for } S > 0 \\ \frac{S+1}{\sigma} & \text{for } S < 0 \\ 0 & \text{for } S = 0 \end{cases} \quad (5)$$

The null hypothesis is rejected at a significance level if $|Z_S| > Z_{crit}$, where Z_{crit} is the value of the standard normal distribution with an exceedance probability of $\alpha/2$. In our analysis, the statistically significant trends are evaluated at the 10%, 5% and 1% significance levels (two-tailed test).

In the SR test, the null hypothesis (H_0) is that all the data X_i in the time series are independent and identically distributed, while the alternative hypothesis (H_1) is that X_i increases or decreases with i , that is trend exists [16]. The SR test statistic D is given by:

$$D = 1 - \frac{6 \sum_{i=1}^n (R_i - i)^2}{n(n^2 - 1)} \quad (6)$$

$$Z_{SR} = D \sqrt{\frac{n-2}{1-D^2}} \quad (7)$$

where, R_i is the rank of i observation X_i in the time series and n is the length of the time series. A positive value of Z_{SR} indicates an upward trend; while negative Z_{SR} indicate a downward trend in the time series. When $Z_{SR} > t_{(n-2, 1-\alpha/2)}$, the null hypothesis is rejected indicating a significant trend in the time series data. Here, $t_{(n-2, 1-\alpha/2)}$ is the critical value of t distribution.

The selected study area covers New South Wales (NSW) state in Australia (Fig. 1). The pluviograph stations were

selected which have a recorded length of greater than 10 years, 30 stations with data ranging between 1932 to 2012 were retained which had less than 20% gaps in the annual maximum rainfall series. Rainfall events of six sub-hourly durations (6, 12, 18, 24, 30 and 48 minutes), six sub-daily durations (60 min, 2, 3, 6, 8 and 12 hours), and three daily durations (24, 48 and 72 hours) were extracted from each of the selected 30 pluviograph stations. The geographical

distributions of the selected 30 pluviograph stations are shown in Fig. 1 which shows a uniform density of selected stations over NSW. Table I lists the selected pluviograph stations with record lengths, period of data and latitudes and longitudes. The record lengths of the annual maximum rainfall events series range from 12 to 72 years (average: 36 years) and the mean annual rainfalls in this region are in the range of 183 mm to 1439 mm (average: 580 mm).

TABLE I
SELECTED PLUVIOGRAPH STATIONS AND RECORD LENGTHS

| Station ID | Station | Data Period | Length (Years) | Latitude (Degree) | Longitude (Degree) |
|------------|---------------|-------------|----------------|-------------------|--------------------|
| 46012 | Wilcannia | 2001-2012 | 12 | -31.5194 | 143.3850 |
| 46037 | Tibooburra | 1985-2012 | 28 | -29.4345 | 142.0098 |
| 47058 | Menindee | 1994-2011 | 18 | -32.3902 | 142.4164 |
| 48015 | Brewarrina | 1997-2012 | 16 | -29.9614 | 146.8651 |
| 48027 | Cobar Mo | 1966-2011 | 49 | -31.484 | 145.8294 |
| 48031 | Collarenebri | 1977-2011 | 35 | -29.5407 | 148.5818 |
| 49000 | Ivanhoe | 2001-2012 | 12 | -32.8831 | 144.3088 |
| 51049 | Trangie | 1969-2012 | 44 | -31.9861 | 147.9489 |
| 55054 | Tamworth | 1959-1992 | 34 | -31.0867 | 150.8467 |
| 56018 | Inverell | 1948-2010 | 63 | -29.7752 | 151.0819 |
| 56041 | Bonshaw | 1955-1969 | 15 | -29.1333 | 151.4500 |
| 57033 | Wollomombi | 1960-1982 | 23 | -30.5112 | 152.0427 |
| 58044 | Nimbin | 1964-2009 | 46 | -28.5966 | 153.2233 |
| 58099 | Whiporie | 1974-2010 | 37 | -29.2823 | 152.9886 |
| 61151 | Chichester | 1961-2012 | 52 | -32.2426 | 151.6830 |
| 61174 | Millfield | 1959-1980 | 22 | -32.9000 | 151.2667 |
| 61309 | Milbrodale | 1970-2012 | 43 | -32.6881 | 150.9728 |
| 61351 | Peats | 1982-2010 | 29 | -33.3102 | 151.2443 |
| 63023 | Cowra | 1942-2010 | 69 | -33.8087 | 148.7071 |
| 63063 | Oberon | 1989-2012 | 24 | -33.6682 | 149.8348 |
| 64046 | Coonabarabran | 1972-2011 | 40 | -31.2886 | 149.0687 |
| 65035 | Wellington | 1962-2004 | 43 | -32.5059 | 148.9708 |
| 67035 | Liverpool | 1966-2001 | 36 | -33.9272 | 150.9128 |
| 69049 | Nerriga | 1972-2010 | 39 | -35.1165 | 150.0847 |
| 70014 | Canberra | 1938-2009 | 72 | -35.3049 | 149.2014 |
| 72091 | Tooma | 1957-1973 | 17 | -35.9111 | 148.2167 |
| 74114 | Wagga Wagga | 1947-2003 | 57 | -35.1311 | 147.3091 |
| 74128 | Deniliquin | 1978-2002 | 25 | -35.5269 | 144.9520 |
| 75028 | Griffith | 1932-1988 | 57 | -34.2487 | 146.0695 |
| 75050 | Naradhan | 1971-2012 | 42 | -33.6104 | 146.3161 |




Fig. 1 Selected pluviograph stations in NSW, Australia

III. RESULTS AND DISCUSSION

The results from the trend analysis are presented in Figs. 2 and 3 (for 5% level of significance) and in Figs. 4 and 5 (for 10% level of significance) by both the MK and SR tests. In Fig. 2, it is found that for the MK test (5% level of significance), the percentages of stations with statistically significant positive trends are notably higher for sub-hourly durations (6 to 48 minutes), for 1 hour the positive and negative trends are very similar, for 2 hours to 1 day, the negative trends are much higher and for 2 and 3 days, positive trends are higher than the negative ones.

In Fig. 3, it is found that the SR test (at 5% level of significance) shows all positive trends for 6 minutes to 48 minutes durations and 1 hour to 6 hours durations, and for 8

hours to 1 day there are more positive trends than the negative ones. In Fig. 4, the MK test results (at 10% significance level) show that 6 minutes to 48 minutes durations have more positive trends than the negative ones, for 1 hour to 1 day, there are more negative trends than the positive ones. In the case of SR test at 5% significance level (Fig. 5), there are more positive trends, in particular for shorter durations.




Fig. 2 Percentage of stations with positive and negative trends (at 5% level) for MK test




Fig. 3 Percentage of stations with positive and negative trends (at 5% level) for SR test




Fig. 4 Percentage of stations with positive and negative trends (at 10% level) for MK test




Fig. 5 Percentage of stations with positive and negative trends (at 10% level) for SR test

Table II shows the percentage of stations showing significant positive and negative trend at 1%, 5% and 10% significance levels. This shows that at 1% level of significance, the percentages of stations indicating positive trend vary from 0% to 10% for the MK and SR tests for different durations. However, the percentages of stations indicating negative trend vary from 0% to 7% for the MK test and for SR test, there is no trend.

| Significance level | 1% | | 5% | | 10% | |
|--------------------|-----------|-----------|------------|-----------|------------|-----------|
| Duration | MK test | SR test | MK test | SR test | MK test | SR test |
| 6 min | 10 (0) | 10 (0) | 13 (3) | 13 (0) | 20 (3) | 17 (3) |
| 12 min | 10 (0) | 10 (0) | 27 (3) | 13 (0) | 30 (7) | 23 (3) |
| 18 min | 10 (3) | 10 (0) | 17 (7) | 13 (0) | 23 (7) | 20 (7) |
| 24 min | 10 (3) | 10 (0) | 17 (10) | 13 (0) | 23 (10) | 20 (7) |
| 30 min | 7 (0) | 7 (0) | 20 (10) | 13 (0) | 23 (10) | 20 (7) |
| 48 min | 3 (0) | 3 (0) | 10 (10) | 13 (0) | 13 (13) | 20 (3) |
| 1 hr | 3 (0) | 3 (0) | 7 (7) | 13 (0) | 10 (13) | 17 (3) |
| 2 hr | 3 (7) | 3 (0) | 3 (7) | 10 (0) | 7 (13) | 17 (7) |
| 3 hr | 3 (3) | 0 (0) | 3 (7) | 10 (0) | 3 (10) | 17 (7) |
| 6 hr | 0 (7) | 0 (0) | 0 (7) | 7 (0) | 7 (10) | 13 (7) |
| 8 hr | 0 (7) | 0 (0) | 0 (7) | 7 (3) | 7 (13) | 10 (7) |
| 12hr | 0 (7) | 0 (0) | 0 (10) | 7 (3) | 3 (13) | 10 (7) |
| 24 hr | 0 (0) | 0 (0) | 0 (7) | 7 (3) | 3 (20) | 7 (7) |
| 48 hr | 0 (0) | 0 (0) | 7 (3) | 0 (0) | 10 (10) | 7 (3) |
| 72 hr | 0 (0) | 0 (0) | 10 (7) | 0 (0) | 10 (10) | 7 (3) |

For the 5% significance level, the percentage of stations with positive trend vary from 0% to 27% and 0% to 13% for the MK and SR tests, respectively for different durations and

the percentages of stations indicating negative trend vary from 0% to 7% and 0% to 3% for different durations for MK and SR tests, respectively. While for the 10% significance level, the percentage of stations with positive trend vary from 0% to 30% and 0% to 17% for the MK and SR tests, respectively for different durations and the percentages of stations indicating negative trend vary from 0% to 20% and 0% to 7% for different durations for both the tests. Overall, the numbers of stations indicating positive trends are greater than negative trends for shorter duration events by both the tests and three significance levels considered in this study.

Fig. 6 shows the spatial distributions of the pluviograph stations showing positive and negative trends for 6 minutes and 48 hours durations for the MK test. This figure shows that there are a greater number of stations in NSW state that show positive trends than negative trends for 6 minutes duration.




Fig. 6 Spatial distributions of stations showing positive, negative and no trends (at the 10% level of significance) for 6 minutes (left) and 48 hours (right) durations using MK test. Blue and red triangles represent stations exhibiting significant positive and negative trends respectively and the hollow circles represent stations indicating no trend

IV. CONCLUSION

This paper examines the trends of sub-hourly, sub-daily and daily extreme rainfall events from 30 rainfall stations across NSW state in Australia. Two non-parametric tests (MK and SR) are applied to detect trends at 1%, 5% and 10% significance levels. It is found that the sub-hourly durations (6, 12, 18, 24, 30 and 48 minutes) show statistically significant positive (upward) trends whereas longer duration (sub-daily and daily) events generally show a statistically significant negative (downward) trend. It is also found that the MK test and SR test provide notably different results for some rainfall event durations considered in this study. Since shorter duration sub-hourly rainfall events show positive trends at many stations, the design rainfall data based on stationary frequency analysis for these durations need to be adjusted to account for the impact of climate change. These shorter durations are more relevant to many urban development projects based on smaller catchments having a much shorter response time.

ACKNOWLEDGMENT

We acknowledge the Australian Bureau of Meteorology for providing data for this study.

REFERENCES

- [1] D. Karpouzou, S. Kavalieratou and C. Babajimopoulos, "Trend analysis of precipitation data in Pieria Region (Greece)," *European Water*, 2010, vol. 30, pp. 31-40.
- [2] W. S. Ashley, T. L. Mote, P. G. Dixon, S. L. Trotter, E. J. Powell, J. D. Durkee and A. J. Grundstein, "Distribution of mesoscale convective complex rainfall in the United States," *Monthly Weather Review*, 2003, vol. 131, pp. 3003-3017.
- [3] M. R. Haylock, T. C. Peterson, L. M. Alves, T. Ambrizzi, Y. M. T. Anuncia, J. Baez, V. R. Barros, M. A. Berlatto, M. Bidegain, G. Coronel, V. Corradi, V. J. Garcia, A. M. Grimm, D. Karoly, J. A. Marengo, M. B. Marino, D. Moncunill, D. Nechet, J. Quintana, E. Rebello, M. Rusticucci, J. L. Santos, I. Trebejo and L. A. Vincent, "Trends in total and extreme south American rainfall in 1960-2000 and links with sea surface temperature," *Journal of Climate*, 2006, vol. 19, pp. 1490-1512.
- [4] A. Burgueno, C. Serra and X. Lana, "Monthly and annual statistical distributions of daily rainfall at the Fabra observatory (Barcelona, NE Spain) for the years 1917-1999," *Theoretical and Applied Climatology*, 2004, vol. 77, pp. 57-75, doi: 10.1007/s00704-003-0020-9.
- [5] T. O. Odekunle, E. E. Balogun and O. O. Ogunkoya, "On the prediction of rainfall onset and retreat dates in Nigeria," *Theoretical and Applied Climatology*, 2005, vol. 81, pp. 101-112, doi: 10.1007/s00704-004-0108-x.
- [6] P. Zhai, X. Zhang, H. Wan, and X. Pan, "Trends in Total Precipitation and Frequency of Daily Precipitation Extremes over China," *Journal of Climate*, 2005, vol. 18, pp. 1096-1108.
- [7] F. Fujibe, "Long-Term Changes in Precipitation in Japan," *Journal of Disaster Research*, 2008, vol. 3, pp. 51-52.
- [8] J. Hannaford and G. Buys, "Trends in seasonal river flow regimes in the UK," *Journal of Hydrology*, 2012, vol. 475, pp. 158-174.
- [9] P. P. Nikhil Raj and P. A. Azeez, "Trend analysis of rainfall in Bharathapuzha River basin, Kerala, India," *International Journal of Climatology*, 2012, vol. 32, pp. 533-539, doi: 10.1002/joc.2283.
- [10] M. Hanif, A. Khan and S. Adnan, "Latitudinal precipitation characteristics and trends in Pakistan," *Journal of Hydrology*, 2013, vol. 492, pp. 266-272.
- [11] D. H. Burn and A. Taleghani, "Estimates of changes in design rainfall values for Canada," *Hydrological Processes*, 2013, vol. 27, pp. 1590-1599, doi: 10.1002/hyp.9238.
- [12] M. R. Hamdi, M. Abu-Allaban, A. Al Shayeb, M. Jaber and N. M. Momani, "Climate change in Jordan: A comprehensive examination approach," *American Journal of Environmental Sciences*, 2009, vol. 5, pp. 58-68.
- [13] C. Clarke, M. Hulley, J. Marsalek and E. Watt, "Stationarity of A max series of short-duration rainfall for long-term Canadian stations: detection of jumps and trends," *Canadian Journal of Civil Engineering*, 2011, vol. 38, pp. 1175-1184.
- [14] A. Rana, C. B. Uvo, L. Bengtsson and P. P. Sarthi, "Trend analysis for rainfall in Delhi and Mumbai, India," *Climate Dynamics*, 2012, vol. 38, pp. 45-56, doi: 10.1007/s00382-011-1083-4.
- [15] A. Mondal, S. Kundu and A. Mukhopadhyay, "Rainfall trend analysis by Mann-Kendall test: a case study of north-eastern part of Cuttack District, Orissa," *International Journal of Geology, Earth and Environmental*, 2012, vol. 2, pp. 70-78.
- [16] S. Yue, P. Pilon and G. Cavadias, "Power of the Mann-Kendall and Spearman's rho tests for detecting monotonic trends in hydrological series," *Journal of Hydrology*, 2002, vol. 259, pp. 254-271.
- [17] N. Plummer, M. J. Salinger, N. Nicholls, R. Suppiah, K. J. Hennessy, R. M. Leighton, B. Trewin, C. M. Page and J. M. Lough, "Changes in climate extremes over the Australian region and New Zealand during the twentieth century," *Climatic Change*, 1999, vol. 42, pp. 183-202.
- [18] M. Haylock and N. Nicholls, "Trends in extreme rainfall indices for an updated high quality data set for Australia, 1910-1998," *International Journal of Climatology*, 2000, vol. 20, pp. 1533-1541.

- [19] P. Y. Groisman, R. W. Knight, D. R. Easterling, T. R. Karl, G. C. Hegerl, and V. N. Razuvayev, "Trends in intense precipitation in the climate record," *Journal of Climate*, 2005, vol. 18, pp. 1326-1350.
- [20] A. J. E. Gallant, K. J. Hennessy and J. Risbey, "Trends in rainfall indices for six Australian regions: 1910-2005," *Australian Meteorological Magazine*, 2007, vol. 56, pp. 223-239.
- [21] L. V. Alexander, X. Zhang, T. C. Peterson, J. Caesar, B. Gleason and A. M. G. Klein Tank, "Global observed changes in daily climate extremes of temperature and precipitation," *Journal of Geophysical Research-Atmospheres*, 2006, vol. 111, doi:10.1029/2005JD006290.
- [22] R. Hardwick Jones, S. Westra and A. Sharma, "Observed relationships between extreme sub-daily precipitation, surface temperature and relative humidity," *Geophysical Research Letters*, 2010, vol. 37, pp. 1-5.
- [23] J. Li, J. Feng and Y. Li, "A possible cause of decreasing summer rainfall in northeast Australia," *International Journal of Climatology*, 2012, vol. 32, pp. 995-1005.
- [24] Y. R. Chen, B. Yu and G. Jenkins, "Secular variation in rainfall intensity and temperature in eastern Australia," *Journal of Hydrometeorology*, 2013, vol. 14, pp. 1356-1363. doi: <http://dx.doi.org/10.1175/JHM-D-12-0110.1>.
- [25] R. K. Chowdhury and S. Beecham, "Australian rainfall trends and their relation to the southern oscillation index," *Hydrological Processes*, 2010, vol. 24, pp. 504-514.
- [26] D. Jakob, D. J. Karoly and A. Seed, "Non-stationarity in daily and sub-daily intense rainfall-Part 1: Sydney, Australia," *Natural Hazards and Earth System Science*, 2011, vol. 11, pp. 2263-2271.
- [27] M. Loveridge and A. Rahman, "Trend analysis of rainfall losses using an event-based hydrological model in eastern NSW," in *Proc. 20th International Congress on Modelling and Simulation, Adelaide, Australia*, 1-6 December 2013, pp. 2569-2575.
- [28] A. G. Yilmaz and B. J. C. Perera, "Changes in intensity-frequency-duration relationship of heavy rainfalls at a station in Melbourne," in *Proc. 20th International Congress on Modelling and Simulation, Adelaide, Australia*, 2013, pp. 2834-2840.
- [29] H. B. Mann, "Non-parametric tests against trend," *Econometrica*, 1945, vol. 13, pp. 245-259.
- [30] M. G. Kendall, *Rank Correlation Methods*, 1975, Griffin, London.