

Solving Linear Matrix Equations by Matrix Decompositions

Yongxin Yuan, Kezheng Zuo

Abstract—In this paper, a system of linear matrix equations is considered. A new necessary and sufficient condition for the consistency of the equations is derived by means of the generalized singular-value decomposition, and the explicit representation of the general solution is provided.

Keywords—Matrix equation, Generalized inverse, Generalized singular-value decomposition.

I. INTRODUCTION

WE consider the solution of the linear matrix equations

$$\begin{cases} A_1X = C_1, \\ A_2Y = C_2, YB_2 = D_2, Y = Y^H, \\ A_3Z = C_3, ZB_3 = D_3, Z = Z^H, \\ A_4X + (A_4X)^H + B_4YB_4^H + C_4ZC_4^H = D_4, \end{cases} \quad (1)$$

where

$$\begin{aligned} A_1 &\in \mathbf{C}^{a_1 \times m}, C_1 \in \mathbf{C}^{a_1 \times n}, \\ A_2, C_2 &\in \mathbf{C}^{a_2 \times p}, B_2, D_2 \in \mathbf{C}^{p \times b_2}, \\ A_3, C_3 &\in \mathbf{C}^{a_3 \times q}, B_3, D_3 \in \mathbf{C}^{q \times b_3}, \\ A_4 &\in \mathbf{C}^{n \times m}, B_4 \in \mathbf{C}^{n \times p}, C_4 \in \mathbf{C}^{n \times q}, \end{aligned}$$

and

$$D_4 = D_4^H \in \mathbf{C}^{n \times n}.$$

Solvability and solutions of matrix equations have been one of principle topics in matrix analysis and its applications. For instance, Mitra [1] considered solutions with fixed ranks for the matrix equations $AX = B$ and $AXB = C$, Mitra [2] gave common solutions of minimal rank of the pair of matrix equations $AX = C, XB = D$. Uhlig [3] gave the maximal and minimal ranks of solutions of the equation $AX = B$. Mitra [4] examined common solutions of minimal rank of the pair of matrix equations $A_1XB_1 = C_1$ and $A_2XB_2 = C_2$. In 2006, Lin and Wang in [5] studied the extreme ranks of solutions to the system of matrix equations $A_1X = C_1, XB_2 = C_2, A_3XB_3 = C_3$ over an arbitrary division ring, which was investigated in [6] and [7]. Liu [8] derived the maximal and minimal ranks of least squares

solutions for $AXB = C$ using the matrix rank method and the normal equation. Cvetković-Ilić [9], Peng, Hu and Zhang [10] considered the reflexive and anti-reflexive solutions of the matrix equation $AXB = C$ by means of generalized inner inverse and the generalized singular-value decomposition. In the papers [11–13], necessary and sufficient conditions for the existence of symmetric and anti-symmetric solutions of the equation $AXB = C$ were investigated.

Wu [14] studied Re-pd solutions of the equation $AX = C$ and Wu and Cain [15] found the set of all complex Re-nnd matrices X such that $XB = C$ and presented a criterion for Re-nndness. Größ [16] gave an alternative approach, which simultaneously delivers explicit Re-nnd solutions and gave a corrected version of some results from [15]. Recently, in [17] and [18], the common Re-nnd and Re-pd solutions of the matrix equations $AX = C, XB = D$, where $A, C \in \mathbf{C}^{n \times m}$ and $B, D \in \mathbf{C}^{m \times n}$, are considered by virtue of the maximal and minimal ranks of matrix polynomials. Wang and Yang [19] presented criteria for 2×2 and 3×3 partitioned matrices to be Re-nnd, found necessary and sufficient conditions for the existence of Re-nnd solutions of $AXB = C$ and derived an expression for these solutions. In the special case that A and B are both nonnegative matrices, Cvetković-Ilić [20] put forward a necessary and sufficient condition for the existence of Re-nnd solutions of $AXB = C$ in terms of g-inverses. Zhang, Sheng and Xu [21] generalized the main results of [20] from the finite-dimensional case to the Hilbert space operator case.

Matrix equations such as $A_4X + (A_4X)^H = D_4, B_4YB_4^H = D_4$ and $B_4YB_4^H + C_4ZC_4^H = D_4$, which are the special cases of (1), arise in a number of practical applications in linear system theory, numerical analysis and structural dynamics, and have been studied by Braden [22], Djordjević [23], Yuan [24, 25], Dai and Lancaster [26], Baksalary [27], Größ [28], Liu, Tian and Takane [29], Liao and Bai [30], Deng and Hu [31] and Liu and Tian [32], and so forth.

Very recently, Wang and He [33] derived the solvability conditions and the expression of the general solution to the matrix equations of (1) by virtue of the maximal and minimal ranks of matrix polynomials. In this paper, we will provide a new approach based on the generalized inverses and the generalized singular-value decomposition (GSVD) to solve (1). In Section II, we establish a necessary and sufficient

Yongxin Yuan: School of Mathematics and Statistics, Hubei Normal University, Huangshi, 435002, PR China (e-mail: yuanyx_703@163.com).

Kezheng Zuo: School of Mathematics and Statistics, Hubei Normal University, Huangshi, 435002, PR China (e-mail: Xiangzuo28@163.com).

condition for the existence of the solution of (1) directly by means of the GSVD, and construct the explicit representation of the general solution when it is solvable. Throughout this paper, we denote the complex $m \times n$ matrix space by $\mathbf{C}^{m \times n}$, the conjugate transpose and the Moore-Penrose generalized inverse of a complex matrix A by A^H and A^+ , respectively. I_n represents the identity matrix of size n . Furthermore, for a matrix $A \in \mathbf{C}^{m \times n}$, let E_A and F_A stand for the two orthogonal projectors: $E_A = I_m - AA^+$ and $F_A = I_n - A^+A$.

II. THE SOLUTION OF (1)

Let $G_1 \in \mathbf{C}^{m \times q}, G_2 \in \mathbf{C}^{n \times d}$, then the GSVD (see, e.g., [34–36]) of the matrix pair (G_1, G_2) is of the form

$$G_1 = N\Omega_1P^H, G_2 = N\Omega_2Q^H, \quad (2)$$

where $P \in \mathbf{C}^{q \times q}, Q \in \mathbf{C}^{d \times d}$ are unitary matrices and $N \in \mathbf{C}^{m \times n}$ is a nonsingular matrix, and

$$\Omega_1 = \begin{bmatrix} I & 0 & 0 \\ 0 & S_1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{matrix} h-t \\ t \\ e-h \\ n-e \end{matrix},$$

$$\Omega_2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & S_2 & 0 \\ 0 & 0 & I \\ 0 & 0 & 0 \\ u & t & e-h \end{bmatrix} \begin{matrix} h-t \\ t \\ e-h \\ n-e \end{matrix},$$

$$u = d + h - e - t, \\ h = \text{rank}(G_1), e = \text{rank}[G_1, G_2],$$

and

$$S_1 = \text{diag} \{ \gamma_1, \dots, \gamma_t \}, \\ S_2 = \text{diag} \{ \delta_1, \dots, \delta_t \}$$

with

$$1 > \gamma_1 \geq \dots \geq \gamma_t > 0, \\ 0 < \delta_1 \leq \dots \leq \delta_t < 1, \\ \gamma_i^2 + \delta_i^2 = 1 \quad (i = 1, \dots, t).$$

Lemma 1: [37] If $A \in \mathbf{C}^{m \times n}, H \in \mathbf{C}^{m \times l}$, then

$$AZ = H$$

has a solution $Z \in \mathbf{C}^{n \times l}$ if and only if

$$AA^+H = H.$$

In this case, the general solution of the equation can be described as

$$Z = A^+H + F_A L,$$

where $L \in \mathbf{C}^{n \times l}$ is an arbitrary matrix.

Lemma 2: [38] Let $A, B \in \mathbf{C}^{n \times p}$, then the matrix equation

$$AY = B$$

has a Hermitian solution $Y \in \mathbf{C}^{p \times p}$ if and only if

$$BA^H = AB^H, AA^+B = B,$$

in which case, the general Hermitian solution is

$$Y = A^+B + F_A(A^+B)^H + F_A K F_A,$$

where $K \in \mathbf{C}^{p \times p}$ is an arbitrary Hermitian matrix.

Lemma 3: [22, 24] Let $A \in \mathbf{C}^{m \times n}, D \in \mathbf{C}^{m \times m}$. Then the matrix equation

$$AX^H + XA^H = D$$

has a solution X if and only if

$$D = D^H, E_A D E_A = 0,$$

in which case, the general solution is

$$X = \frac{1}{2}D(A^+)^H + \frac{1}{2}E_A D(A^+)^H + V - \frac{1}{2}V A^+ A \\ - \frac{1}{2}A V^H(A^+)^H - \frac{1}{2}E_A V A^+ A,$$

where $V \in \mathbf{C}^{m \times n}$ is an arbitrary matrix.

It follows from Lemma 1 that the matrix equation $A_1 X = C_1$ has a solution $X \in \mathbf{C}^{m \times n}$ if and only if

$$A_1 A_1^+ C_1 = C_1.$$

In this case, the general solution of the equation can be described as

$$X = A_1^+ C_1 + F_{A_1} L, \quad (3)$$

where $L \in \mathbf{C}^{m \times n}$ is an arbitrary matrix.

Let

$$A^H = [A_2^H, B_2^H]^H, B^H = [C_2^H, D_2^H]^H.$$

By Lemma 2, we know the matrix equations

$$A_2 Y = C_2, Y B_2 = D_2$$

has a Hermitian solution Y if and only if

$$AA^+B = B, AB^H = BA^H,$$

in which case the general Hermitian solution of the equation can be expressed as

$$Y = A^+B + F_A(A^+B)^H + F_A K F_A, \quad (4)$$

where $K \in \mathbf{C}^{p \times p}$ is an arbitrary Hermitian matrix.

Likewise, let

$$C^H = [A_3^H, B_3^H]^H, D^H = [C_3^H, D_3^H]^H,$$

then the matrix equations

$$A_3 Z = C_3, Z B_3 = D_3$$

has a Hermitian solution Z if and only if

$$CC^+D = D, CD^H = DC^H,$$

in which case the general Hermitian solution of the equation can be expressed as

$$Z = C^+D + F_C(C^+D)^H + F_C J F_C, \quad (5)$$

where $J \in \mathbb{C}^{q \times q}$ is an arbitrary Hermitian matrix.

By substituting (3), (4) and (5) into the fourth equation of (1), we can obtain

$$M_1L + L^H M_1^H + M_2K M_2^H + M_3J M_3^H = W_1, \quad (6)$$

where

$$M_1 = A_4 F_{A_1}, \quad M_2 = B_4 F_A, \quad M_3 = C_4 F_C$$

and

$$W_1 = D_4 - A_4 A_1^+ C_1 - (A_4 A_1^+ C_1)^H - B_4 A^+ B B_4^H - B_4 F_A (B_4 A^+ B)^H - C_4 C^+ D C_4^H - C_4 F_C (C_4 C^+ D)^H.$$

According to Lemma 3, the equation of (6) with respect to L is solvable if and only if

$$G_1 K G_1^H + G_2 J G_2^H = W, \quad (7)$$

and the general solution of (6) with respect to L can be expressed as

$$L = \frac{1}{2} M_1^+ \tilde{D} + \frac{1}{2} M_1^+ \tilde{D} E_{M_1} + U - \frac{1}{2} M_1^+ M_1 U - \frac{1}{2} M_1^+ U^H M_1^H - \frac{1}{2} M_1^+ M_1 U E_{M_1}, \quad (8)$$

where

$$\begin{aligned} G_1 &= E_{M_1} M_2, \\ G_2 &= E_{M_1} M_3, \\ W &= E_{M_1} W_1 E_{M_1}, \\ \tilde{D} &= W_1 - M_2 K M_2^H - M_3 J M_3^H \end{aligned}$$

and U is an arbitrary matrix.

By (2), the equation of (7) can be equivalent written as

$$\Omega_1 P^H K P \Omega_1^H + \Omega_2 Q^H J Q \Omega_2^H = N^{-1} W (N^{-1})^H. \quad (9)$$

Write

$$P^H K P = [K_{ij}]_{3 \times 3}, \quad K_{ij} = K_{ij}^H, \quad i, j = 1, 2, 3, \quad (10)$$

$$Q^H J Q = [J_{ij}]_{3 \times 3}, \quad J_{ij} = J_{ij}^H, \quad i, j = 1, 2, 3, \quad (11)$$

$$N^{-1} W (N^{-1})^H = [W_{ij}]_{4 \times 4}, \quad W_{ij} = W_{ij}^H, \quad i, j = 1, 2, 3, 4, \quad (12)$$

and the partitions of the matrices $P^H K P, Q^H J Q$ and $N^{-1} W (N^{-1})^H$ are compatible with those of Ω_1 and Ω_2 . Thus, from (9), we have

$$\begin{aligned} & \begin{bmatrix} K_{11} & K_{12} S_1 & 0 & 0 \\ S_1 K_{12}^H & S_1 K_{22} S_1 + S_2 J_{22} S_2 & S_2 J_{23} & 0 \\ 0 & J_{23}^H S_2 & J_{33} & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \\ &= \begin{bmatrix} W_{11} & W_{12} & W_{13} & W_{14} \\ W_{12}^H & W_{22} & W_{23} & W_{24} \\ W_{13}^H & W_{23}^H & W_{33} & W_{34} \\ W_{14}^H & W_{24}^H & W_{34}^H & W_{44} \end{bmatrix}. \end{aligned} \quad (13)$$

By (13), we can get

$$\begin{aligned} W_{13} &= 0, \quad W_{i4} = 0, \quad i = 1, 2, 3, 4, \\ K_{11} &= W_{11}, \quad K_{12} S_1 = W_{12}, \\ S_1 K_{22} S_1 + S_2 J_{22} S_2 &= W_{22}, \\ J_{33} &= W_{33}, \quad S_2 J_{23} = W_{23}. \end{aligned}$$

In summary of above discussion, we can easily obtain the following result.

Theorem 1: Suppose that

$$\begin{aligned} A_1 &\in \mathbb{C}^{a_1 \times m}, \quad C_1 \in \mathbb{C}^{a_1 \times n}, \\ A_2, C_2 &\in \mathbb{C}^{a_2 \times p}, \quad B_2, D_2 \in \mathbb{C}^{p \times b_2}, \\ A_3, C_3 &\in \mathbb{C}^{a_3 \times q}, \quad B_3, D_3 \in \mathbb{C}^{q \times b_3}, \\ A_4 &\in \mathbb{C}^{n \times m}, \quad B_4 \in \mathbb{C}^{n \times p}, \quad C_4 \in \mathbb{C}^{n \times q} \end{aligned}$$

and

$$D_4 = D_4^H \in \mathbb{C}^{n \times n}.$$

Let

$$\begin{aligned} A^H &= [A_2^H, B_2]^H, \quad B^H = [C_2^H, D_2]^H, \\ C^H &= [A_3^H, B_3]^H, \quad D^H = [C_3^H, D_3]^H, \\ M_1 &= A_4 F_{A_1}, \quad M_2 = B_4 F_A, \quad M_3 = C_4 F_C, \\ W_1 &= D_4 - A_4 A_1^+ C_1 - (A_4 A_1^+ C_1)^H - B_4 A^+ B B_4^H \\ &\quad - B_4 F_A (B_4 A^+ B)^H - C_4 C^+ D C_4^H - C_4 F_C (C_4 C^+ D)^H, \\ G_1 &= E_{M_1} M_2, \quad G_2 = E_{M_1} M_3, \quad W = E_{M_1} W_1 E_{M_1}, \end{aligned}$$

the GSVD of (G_1, G_2) be given by (2) and $N^{-1} W (N^{-1})^H = [W_{ij}]_{4 \times 4}$ be given by (12). Then the equation of (1) has a solution (X, Y, Z) if and only if

$$\begin{aligned} A_1 A_1^+ C_1 &= C_1, \quad A A^+ B = B, \quad A B^H = B A^H, \\ C C^+ D &= D, \quad C D^H = D C^H, \\ W_{13} &= 0, \quad W_{i4} = 0, \quad i = 1, 2, 3, 4, \end{aligned}$$

in which case, the general solution can be expressed as

$$\begin{aligned} X &= A_1^+ C_1 + F_{A_1} L, \\ Y &= A^+ B + F_A (A^+ B)^H + F_A K F_A, \\ Z &= C^+ D + F_C (C^+ D)^H + F_C J F_C, \end{aligned}$$

where L is given by (8) with

$$\tilde{D} = W_1 - M_2 K M_2^H - M_3 J M_3^H,$$

and

$$K = P \begin{bmatrix} W_{11} & W_{12} S_1^{-1} & K_{13} \\ S_1^{-1} W_{12}^H & S_1^{-1} (W_{22} - S_2 J_{22} S_2) S_1^{-1} & K_{23} \\ K_{13}^H & K_{23}^H & K_{33} \end{bmatrix} P^H,$$

$$J = Q \begin{bmatrix} J_{11} & J_{12} & J_{13} \\ J_{12}^H & J_{22} & S_2^{-1} W_{23} \\ J_{13}^H & W_{23}^H S_2^{-1} & W_{33} \end{bmatrix} Q^H,$$

and

$$U, K_{13}, K_{23}, K_{33} = K_{33}^H, J_{11} = J_{11}^H, J_{22} = J_{22}^H, J_{12}, J_{13}$$

are all arbitrary matrices.

REFERENCES

- [1] S. K. Mitra, Fixed rank solutions of linear matrix equations, *Sankhya Ser. A.*, 35 (1972) 387–392.
- [2] S. K. Mitra, The matrix equation $AX = C, XB = D$, *Linear Algebra and its Applications*, 59 (1984) 171–181.
- [3] F. Uhlig, On the matrix equation $AX = B$ with applications to the generators of a controllability matrix, *Linear Algebra and its Applications*, 85 (1987) 203–209.
- [4] S. K. Mitra, A pair of simultaneous linear matrix equations $A_1 X B_1 = C_1, A_2 X B_2 = C_2$ and a matrix programming problem, *Linear Algebra and its Applications*, 131 (1990) 107–123.
- [5] C. Y. Lin, Q. W. Wang, The minimal and maximal ranks of the general solution to a system of matrix equations over an arbitrary division ring, *Math. Sci. Res. J.*, 10 (2006) 57–65.
- [6] P. Bhimasankaram, Common solutions to the linear matrix equations $AX = B, XC = D$, and $EXF = G$, *Sankhya Ser. A.*, 38 (1976) 404–409.
- [7] C. Y. Lin, Q. W. Wang, New solvable conditions and a new expression of the general solution to a system of linear matrix equations over an arbitrary division ring, *Southeast Asian Bull. Math.*, 29 (2005) 755–762.
- [8] Y. Liu, Ranks of least squares solutions of the matrix equation $AXB = C$, *Computers and Mathematics with Applications*, 55 (2008) 1270–1278.
- [9] D. S. Cvetković-Ilić, The reflexive solutions of the matrix equation $AXB = C$, *Computers and Mathematics with Applications*, 51 (2006) 897–902.
- [10] X. Peng, X. Hu, L. Zhang, The reflexive and anti-reflexive solutions of the matrix equation $A^H X B = C$, *Journal of Computational and Applied Mathematics*, 200 (2007) 749–760.
- [11] C. G. Khatri, S. K. Mitra, Hermitian and nonnegative definite solutions of linear matrix equations, *SIAM Journal on Applied Mathematics*, 31 (1976) 579–585.
- [12] Z. Y. Peng, An iterative method for the least squares symmetric solution of the linear matrix equation $AXB = C$, *Applied Mathematics and Computation*, 170 (2005) 711–723.
- [13] Y. X. Peng, X. Y. Hu, L. Zhang, An iteration method for the symmetric solutions and the optimal approximation solution of the matrix equation $AXB = C$, *Applied Mathematics and Computation*, 160 (2005) 763–777.
- [14] L. Wu, The Re-positive definite solutions to the matrix inverse problem $AX = B$, *Linear Algebra and its Applications*, 174 (1992) 145–151.
- [15] L. Wu, B. Cain, The Re-nonnegative definite solutions to the matrix inverse problem $AX = B$, *Linear Algebra and its Applications*, 236 (1996) 137–146.
- [16] J. GröB, Explicit solutions to the matrix inverse problem $AX = B$, *Linear Algebra and its Applications*, 289 (1999) 131–134.
- [17] Z. Xiong, Y. Qin, The common Re-nnd and Re-pd solutions to the matrix equations $AX = C$ and $XB = D$, *Applied Mathematics and Computation*, 218 (2011) 3330–3337.
- [18] X. Liu, Comments on “The common Re-nnd and Re-pd solutions to the matrix equations $AX = C$ and $XB = D$ ”, *Applied Mathematics and Computation*, 236 (2014) 663–668.
- [19] Q. Wang, C. Yang, The Re-nonnegative definite solutions to the matrix equation $AXB = C$, *Comment. Math. Univ. Carolinae*, 39 (1998) 7–13.
- [20] D. S. Cvetković-Ilić, Re-nnd solutions of the matrix equation $AXB = C$, *Journal of the Australian Mathematical Society*, 84 (2008) 63–72.
- [21] X. Zhang, L. Sheng, Q. Xu, A note on the real positive solutions of the operator equation $AXB = C$, *Journal of Shanghai Normal University (Natural Sciences)*, 37 (2008) 454–458.
- [22] H. W. Braden, The equation $A^T X \pm X^T A = B$, *SIAM J Matrix Anal Appl.*, 20 (1998) 295–302.
- [23] D. S. Djordjević, Explicit solution of the operator equation $A^* X \pm X^* A = B$, *Journal of Computational and Applied Mathematics*, 200 (2007) 701–704.
- [24] Y. Yuan, On the symmetric solutions of a class of linear matrix equation, *Chinese Journal of Engineering Mathematics*, 15 (1998) 25–29.
- [25] Y. Yuan, The minimum norm solutions of two classes of matrix equations, *Numer. Math. J. Chinese Univ.*, 24 (2002) 127–134.
- [26] H. Dai, P. Lancaster, Linear matrix equations from an inverse problem of vibration theory, *Linear Algebra Appl.*, 246 (1996) 31–47.
- [27] J. K. Baksalary, Nonnegative definite and positive definite solutions to the matrix equation $AXA^* = B$, *Linear Multilinear Algebra*, 16 (1984) 133–139.
- [28] J. GröB, Nonnegative-definite and positive-definite solutions to the matrix equation $AXA^* = B$ revisited, *Linear Algebra Appl.*, 321 (2000) 123–129.
- [29] Y. H. Liu, Y. G. Tian, Y. Takane, Ranks of Hermitian and skew-Hermitian solutions to the matrix equation $AXA^* = B$, *Linear Algebra Appl.*, 431 (2009) 2359–2372.
- [30] A. P. Liao, Z. Z. Bai, The constrained solutions of two matrix equations, *Acta Math Sin English Ser.*, 18 (2002) 671–678.
- [31] Y. B. Deng, X. Y. Hu, On solutions of matrix equation $AXA^T + BYB^T = C$, *J Comput Math.*, 23 (2005) 17–26.
- [32] Y. H. Liu, Y. G. Tian, A simultaneous decomposition of a matrix triplet with applications, *Numer Linear Algebra Appl.*, 18 (2011) 69–85.
- [33] Q.-W. Wang, Z.-H. He, A system of matrix equations and its applications, *Sci China Math.*, 56 (2013) 1795–1820.
- [34] G. H. Golub, C. F. Van Loan, *Matrix Computations*, The Johns Hopkins University Press, Baltimore, 1983.
- [35] C. C. Paige, M. A. saunders, Towards a generalized singular value decomposition, *SIAM J Numer. Anal.*, 18 (1981) 398–405.
- [36] G. W. Stewart, Computing the CS-decomposition of a partitioned orthogonal matrix, *Numer Math.*, 40 (1982) 297–306.
- [37] A. Ben-Israel, T. N. E. Greville. *Generalized Inverses. Theory and Applications* (second ed). New York: Springer, 2003.
- [38] F. J. H. Don, On the symmetric solutions of a linear matrix equation, *Linear Algebra Appl.*, 93 (1987) 1–7.