

# Assessment of Drought Tolerance Maize Hybrids at Grain Growth Stage in Mediterranean Area

Ayman El Sabagh, Celalettin Barutçular, Hirofumi Saneoka

**Abstract**—Drought is one of the most serious problems posing a grave threat to cereals production including maize. Maize improvement in drought-stress tolerance poses a great challenge as the global need for food and bio-energy increases. Thus, the current study was planned to explore the variations and determine the performance of target traits of maize hybrids at grain growth stage under drought conditions during 2014 under Adana, Mediterranean climate conditions, Turkey. Maize hybrids (Sancia, Indaco, 71May69, Aaccel, Calgary, 70May82, 72May80) were evaluated under (irrigated and water stress). Results revealed that, grain yield and yield traits had a negative effects because of water stress conditions compared with the normal irrigation. As well as, based on the result under normal irrigation, the maximum biological yield and harvest index were recorded. According to the differences among hybrids were found that, significant differences were observed among hybrids with respect to yield and yield traits under current research.

Based on the results, grain weight had more effect on grain yield than grain number during grain filling growth stage under water stress conditions. In this concern, according to low drought susceptibility index (less grain yield losses), the hybrid (Indaco) was more stable in grain number and grain weight. Consequently, it may be concluded that this hybrid would be recommended for use in the future breeding programs for production of drought tolerant hybrids.

**Keywords**—Drought susceptibility index, grain filling, grain yield, maize, water stress.

## I. INTRODUCTION

**D**ROUGHT stress is considered one of the most common factors of limiting plant growth in arid and semi-arid regions [1]. The susceptibility of plants to drought stress varies in dependence of stress degree, different accompanying stress factors, plant species, and their developmental stages [2]. Acclimation of plants to water deficit is the result of different events, which lead to adaptive changes in plant growth and physico-biochemical processes, such as changes in plant structure, growth rate, tissue osmotic potential, and antioxidant defenses [3].

Maize crop plays an important role in the world economy and is valuable ingredient in manufactured items that affect a large proportion of the world population [4]. 20-25 percent of the planting area of maize is affected by drought pressure in the world [5]. Maize is one of the most important cereal crops

in Turkey [6]. The cultivated area is about 0.60 million hectares and the production in Turkey are about 4.25 million tons and it covers about 95% of corn consumption in the country [7]. Grain yield reduction of maize due to the drought pressure is varied between 1% to 76% depending on the severity, timing, and stage of occurrence [8], [9].

Loss of yield is the main concern of plant breeders and they hence emphasize on yield performance under stress conditions. Thus, drought indices, which provide a measure of drought based on loss of yield under drought-conditions in comparison to normal conditions, have been used for screening drought/tolerant genotypes [10], [11]. Khalili et al. [12] reported that the yield decrease under drought stress at the reproductive stage was greater than that at the vegetative and grain filling stages. The flowering and grain setting stages appear to be the most sensitive stages to water stress. Limited irrigation at critical stages of growth and development may be crucial for recognition of tolerant maize hybrids. Limited irrigation at critical stages of growth and development may be crucial for recognition of tolerant maize hybrids. Thus, the objective of this research was to evaluate the performance of maize hybrids under water deficit stress at grain filling stage.

## II. THE STUDY AREA

This project was performed in the experimental area of Agricultural Faculty in Cukurova University, Turkey.

## III. MATERIALS AND METHODS

### A. Plant Material and Water Stress Treatments

Seven accessions of maize were chosen based on their morphological and agronomic diversity to formulate the representative core sample of the hybrids. Two factors (F) under current study, F1 hybrids, Sancia, Indaco, 71May69, Aaccel, Calgary, 70May82, 72May80 and F2, water stress control and deficit of water during the grain growth stage in Table I.

The experiment was laid out in strip-split design having four replications. Each plot was of 10 m length and 5.6 m width containing plant density (Intrarow: 70 cm, Interrow: 17 cm). In addition, in order to prevent adjacent treatments from overlapping with each other, 2 row was planted as marginal one between minor plots. Hybrids were sown on June 28, 2014. The regular tillage and agricultural operations of growing maize of the location were followed. All other agronomic practices were kept normal and uniform for all the treatments. Data on various yield components was collected by using standard procedures. During experiments,

A. El Sabagh is with the Department of Agronomy, Faculty of Agriculture, Kafrelsheikh University, Egypt (Corresponding author, e-mail: aymanelsabagh@gmail.com).

C. Barutçular is with Department of Field Crops, Faculty of Agriculture, Cukurova University, Turkey (e-mail: cebar@cu.edu.tr).

H. Saneoka is with Plant Nutritional Physiology, Graduate School of Biosphere Science, Hiroshima University, Japan (e-mail: saneoka@hiroshima-u.ac.jp).

nitrogenous fertilizer was utilized within 2 times of planting, sowing 100 kg N and P<sub>2</sub>O<sub>5</sub>ha<sup>-1</sup> (20-20-0) and V6-growth stage 200 kg N ha<sup>-1</sup> (Urea). Planting was performed with drill.

TABLE I  
IRRIGATION DATE AND AMOUNT IN MAIZE

| Irrigation No    | Date       | Control, mm | Deficit irrigation, mm |
|------------------|------------|-------------|------------------------|
| 1                | 30.06.2014 | 48          | 48                     |
| 2                | 11.07.2014 | 48          | 48                     |
| 3                | 24.07.2014 | 48          | 48                     |
| 4                | 06.08.2014 | 48          | 48                     |
| 5                | 16.08.2014 | 48          | 24                     |
| 6- (Pollination) | 23.08.2014 | 48          | 24                     |
| 7                | 30.08.2014 | 48          | 24                     |
| 8                | 06.09.2014 | 48          | 24                     |
| 9                | 14.09.2014 | 48          | 24                     |
| Total            |            | 432         | 312                    |

#### B. Plant Sampling and Measurements

- Measurements: grain weight (mg), grain number per m<sup>2</sup>, grain yield (t ha<sup>-1</sup>), biomass (t ha<sup>-1</sup>) and harvest index (%).
- Estimation of Drought Susceptibility Index: A drought susceptibility index (DSI) for grain yield and its components was calculated according to Fischer and Maurer [13].
- Statistical analysis: All data collected for both seasons were subjected to analysis of variance according to Gomez and Gomez [14]. Treatment means were compared using Duncan Multiple Range Test [15]. All statistical analysis performed using analysis of variance technique by "MSTAT-C" computer software package

TABLE II  
MEANS OF GRAIN WEIGHT, SEEDS PERM<sup>2</sup>, AS INFLUENCED BY WATER STRESS IN MAIZE HYBRIDS

| Factor         | Grain weight (mg) | Grains per m <sup>2</sup> | Grain yield (t ha <sup>-1</sup> ) | Biological yield (t ha <sup>-1</sup> ) | Harvest index (%) |
|----------------|-------------------|---------------------------|-----------------------------------|--|-------------------|
| Irrigation (A) |                   |                           |                                   |  |                   |
| Control        | 258 a             | 4985                      | 12.72 a                           | 2271 a                                 | 56.0 a            |
| Stress         | 232 b             | 4586                      | 10.50 b                           | 1954 b                                 | 53.7 b            |
| F. test        | **                | NS                        | **                                | **                                     | *                 |
| Hybrid (B)     |                   |                           |                                   |  |                   |
| Sancia         | 236 bc            | 4869 bc                   | 11.30 b                           | 2082                                   | 54.2 bc           |
| Indaco         | 271 a             | 4614 bc                   | 12.52 a                           | 2280                                   | 54.8 abc          |
| 71 May 69      | 232 bc            | 4972 ab                   | 11.45 ab                          | 2036                                   | 56.2 ab           |
| Aaccel         | 263 ab            | 4453 bc                   | 11.65 ab                          | 2054                                   | 56.6 a            |
| Calgary        | 223 c             | 5478 a                    | 12.12 ab                          | 2198                                   | 55.1 abc          |
| 70May82        | 257 ab            | 4310 c                    | 11.12 b                           | 2075                                   | 53.3 c            |
| 72May80        | 234 bc            | 4802 bc                   | 11.13 b                           | 2060                                   | 54.1 bc           |
| F. test        | **                | **                        | *                                 | NS                                     | *                 |
| Interaction    |                   |                           |                                   |  |                   |
| A x B          | NS                | NS                        | NS                                | NS                                     | NS                |

\*, \*\* and NS indicate  $P < 0.05$ ,  $P < 0.01$  and not significant, respectively. Means within the same column of each factor followed by a common letter are not significantly different at the 5% level, according to Duncan's multiple range test.

#### B. Grain Yield

According to the results for grain yield show that decreased significantly with under water stress and a greater yield was observed in indaco, 71May69, Aaccel and Calgary in Table II. Grain yield is the result of the expression and association of several plant growth components. The deficiency of water leads to severe decline in yield of crop plants probably by

1990.

#### IV. RESULTS AND DISCUSSION

Detailed discussions on the results presented in this chapter have been made under the following heads.

##### A. Agronomic Traits

Observations of yield traits for plants are summarized in Table II. Data showed that for yield components of maize hybrids were significantly affected by irrigation treatments. Data in Table II indicated that water stress lead to a significant reduction in seed weight and seed per m<sup>2</sup>. The results showed that the stress resulted in higher value for normal irrigation compared with the water stress treatment. According to The hybrids, the results showed significant differences in grain yield and yield traits. The hybrids 71May69 and Calgary had higher values of grains per m<sup>2</sup> under the stressed treatment. In this concern, 70May82 and Aaccel more positive effect of grain weight. These results agree with [16], which reported that irrigation disruption at grain filling stage causes a decrease in the leaf area duration and photosynthate mobilization to seeds and thereby decreasing grain weight. Similar results were reported by [17] that kernel number per plant is moisture stress-dependent and concluded that kernel number decrease is the primary effect of water deficit on corn grain yield. Zinselmeier et al. [18] stated that the drought effect on number of grains and grain weight, grain yield was reduced.

disrupting leaf gas exchange properties which not only limited the size of the source and sink tissues but the phloem loading, assimilate translocation and dry matter partitioning are also impaired [19]. Pandey et al. [20] reported that yield reduction (22.6–26.4%) caused by deficit irrigation was associated with a decrease in kernel number and weight. The grain filling stage it had the maximum effect on this trait, showing the

severe effects of drought stress at this stage of growth, results that are in agreement with [21].

### C. Biological Yield

The resulted in irrigated treatment higher biomass, compared with the stressed treatment in Table II. Moreover, there is significant different between hybrids in biomass yield. The highest of biological yield was achieved in control with compare water stress treatments. In this concern, the hybrids Indiaco and Calgary were achieved the best value of biomass yield. These results agree with [22], which reported that biomass was reduced by moisture stress. Stone et al. [23] stated that yield was related strongly to biomass especially that accumulated after silking and reported that biomass was reduced by moisture stress.

### D. Harvest Index

The physiological efficiency of a crop to convert dry matter into economic yield is determined by the harvest index. The results show that water stress leads to a decrease in dry material yield and grain yield, so the harvest index does not differ so much. The highest of harvest index was achieved in control with significant differences from other treatment in Table II. According to hybrids, Aaccel, Indaco, Calgary and 71May69 produced the highest values compared with other hybrids in harvest index. The decrease in harvest index under water deficit stress showed the fact that both grain yield decreased under drought stress, but grain yield decreased more than decreased harvest index. With drought, stress had decreased seed yield and harvest index. Therefore, difference between hybrids in control was little and under drought stress, this difference had expanded. These results are in line with the findings of [24] and [25] that skipping irrigation at different crop growth stages significantly influenced different maize hybrids for the parameter of harvest index.

### E. Drought Susceptibility Index

The drought susceptibility index (DSI) estimates the rate of change for each hybrid in yield, between the stress and non-stress conditions relative to the mean change for all hybrids. Values of DSI lower than 1 denotes low drought susceptibility (or drought tolerance) and values higher than 1 indicate high drought susceptibility or poor yield in Fig. 1.




Fig. 1 The drought susceptibility index for grain yield, grain number and grain weight

Depending on the mean DSI value based on grain yield, varieties could be ordered according to water stress tolerance as follow; Indaco, 71May69, Calgary and 72May80, respectively. From the result of DSI (less grain yield losses) this hybrids (Indaco) was more stable in grain number and grain weight. Additionally, it is less grain yield losses under drought stress. Using DSI, Indaco was selected as more stable hybrid in grain growth stages of stress (Fig. 1). Indaco yield was relatively high in all conditions. This finding is consistent with that reported by [26]. Fayaz and Arzani [27] reported cultivars with low DSI values are drought tolerance because they have lesser reduction in grain yield under stress compared with non-stress condition. Moreover, [28] indicated that genotypes with low DSI values might represent a valuable genetic resource for enlarging the genetic variation of barley breeding programs for drought tolerance.

### V. CONCLUSION

This result indicated that grain weight had more effect on grain yield than grain number during grain filling growth stage under water stress conditions. Accordingly, drought susceptibility stress index for grain weight could be recommended in future breeding programs for a selection as criteria in identifying high yielding hybrids under stress conditions. In this concern, the low drought susceptibility index (less grain yield losses) the hybrid (Indaco) was more stable in grain number and grain weight. Consequently, it may be concluded that this hybrid would be perform better under conditions of poor water supply and it could be recommended for use in future breeding programs for production of drought tolerant hybrids.

### ACKNOWLEDGMENT

This research was supported by The Scientific and Technological Research Council of Turkey (TUBITAK-BIDEB-1059B161400153).

### REFERENCES

- [1] H. Turhan, I. Baser, "In vitro and in vivo waterstress in sunflower (*Helianthus annuus* L.)". *Helia*, 27: 227-236, 2004.
- [2] K. Demirevska, D. Zasheva, R. Dimitrov, L. Simova-Stoilova, M. Stamenova, U. Feller, "Drought stress effects on Rubisco in wheat: changes in the Rubisco large subunit". *Acta Physiol. Plant.*, 31: 1129-1138, 2009.
- [3] B. Duan, Y. Yang, Y. Lu, H. Korpelainen, F. Berninger, C. Li, "Interactions between drought stress, ABA and genotypes in *Picea asperata*". *J. Exp. Bot.*, 58: 3025-3036, 2007.
- [4] M.B.M. Alvi, M. Rafique, S. Tariq, A. Hussain, T. Mahmood, M. Sarwar, "Character association and path coefficient analysis of grain yield and yield components maize (*Zea mays* L.)". *Pakistan Journal Biology Science*, 6: 136-138, 2003.
- [5] M. Golbashi, M. Ebrahimi, S. KhavariKhorasani, R. Choukan, "Evaluation of droughttolerance of some corn (*Zea mays* L.) hybrids in Iran". *African Journal of Agricultural Research*, 5: 2714-2719, 2010.
- [6] E. İlker, "Correlation and path coefficient analyses in sweet corn". *Turk. J. Field Crops* 16(2): 105-107, 2011.
- [7] FAO. "http://faostat.fao.org (accessed 11.21.11)". 2009.
- [8] K.H. Mostafavi, H. Sadeghi Geive, M. Dadresan, "Effects of drought stress on germination indices of corn hybrids (*Zea mays* L.)". *International J. Agric. Sci.*, 1 (2):10-18, 2011.
- [9] M. Zarabi, I. Alahdadi, G.A. Akbari, G. A. Akbari, "A study on the effects of different biofertilizer combinations on yield, its components

- and growth indices of corn (*Zea mays* L.) under drought stress condition". Afr J Agric Res 6(3): 681-685, 2011.
- [10] J. Mitra, "Genetics and genetic improvement of drought resistance in crop plants". Curr Sci India 80:758-762, 2001.
- [11] M.R. Naghavi, A.R. Pour-Aboughadareh, M. Khalili, "Evaluation of drought tolerance indices for screening some of corn (*Zea mays* L.) cultivars under environmental conditions". Notulae Scientia Biologicae, 5(3):388-393, 2013.
- [12] M. Khalili, M. Moghaddam, H. Kazemi Arbat, M.R. Shakiba, H. Kanooni, R. Choukan, "Effect of drought stress on different corn genotypes". J. Agric. Sci. 2(20): 67-84, (010)
- [13] R.A. Fischer, R. Maurer, "Drought resistance in spring wheat varieties. 1: Grain yield response". Australian Journal of Agricultural Research. 29, 897- 912, 1978.
- [14] K.A. Gomez, A.A. Gomez. "Statistical procedures for Agricultural Research", 2nd Ed. Johwiley and sons, Inc. New York. 1984.
- [15] B.D. Duncan "Multiple range and multiple F. Test". Biometrics, II: 1-42, 1955.
- [16] K. Ghassemi-Golezani, R. Lotfi, "Response of soybean cultivars to water stress at reproductive stages". International Journal of Plant, Animal and Environmental Sciences, 17: 198-202, 2012.
- [17] A. Yazar, T.A. Howell, D.A. Dusek, K.S. Copeland, "Evaluation of crop water stress index for LEPA irrigated corn". Irrig. Sci. 18, 171-180, 1999.
- [18] C. Zinselmeier, M.J. Lauer, J.S. Boyer, "Reversing drought-induced losses in grain yield: sucrose Maintains embryo growth in maize". Crop Sci., 35: 1390-1400, 1995.
- [19] M. Farooq, A. Wahid, N. Kobayashi, D. Fujita, S.M.A. Basra, "Plant drought stress: effects, mechanisms and management". Agron.Sustain. Dev., 29: 185-212, 2009.
- [20] R.K. Pandey, J.W. Maranville, A. Admou, "Deficit irrigation and nitrogen effects on maize in a Sahelian environment. I. Grain yield and yield components". Agric. Water Manage. 46(1): 113, 2000.
- [21] S. Karimian, S.A.M. Modares Sanavi, A. Sepehri "Effect of water stress at vegetative and generative growth stages in leafy hybrids and commercial corn". Agric. Res. Water, Soil Plant, 5(3): 39-51, 2005.
- [22] S.B. Moser, B. Feil, S. Jampatong, P. Stamp, "Effect of pre-anthesis drought nitrogen fertilizer rate and variety on grain yield, yield components and harvest index of tropical maize". Agric. Water Manage., 81: 41-58, 2006.
- [23] P.J. Stone, D.R. Wilson, J.B. Reid, R.N. Gillespie, "Water deficit effects on sweet corn. I. Water use, radiation use efficiency growth and yield". Aust. J. Agr. Res., 52(1): 103-113, 2001.
- [24] T.Y.M. Bayoumi, H. Eid, E.M. Metwali, "Application of physiological and biochemical indices as a screening technique for drought tolerance in wheat genotypes". African J. Biotech.7: 2341-2352, 2008.
- [25] P. Manivannan, C.A. Jaleel, A. Kishorekumar, B. Sankar, R. Somasundaram, R. Sridharan, R. Panneerselvam, "Changes in antioxidant metabolism of *Vigna unguiculata* (L.) Walp. by propiconazole under water deficit stress". Colloids Surf. B Biointerfaces, 57: 69- 74, 2007.
- [26] Z. Khodarahmpour, R. Choukan, M.R. Bihamta, E. MajidiHarvan, "Determination of the best heat stress tolerance indices in maize (*Zea mays* L.) inbred lines and hybrids under Khuzestan province conditions". J. Agric. Sci. Tech. 13: 111-121, 2011.
- [27] N. Fayaz, A. Arzani, "Moisture stress tolerance in reproductive growth stages in triticale (X *Triticosecale* Wittmack) cultivars under field conditions". Crop Breeding J. 1,1., 2011.
- [28] M. Haddadin, A. Abdel-Ghani, N. Al-Majali, "Response of barley varieties to drought stress imposed at different developmental stages". Jordan J Agric Sci. 9:507-524, 2013.