

# Adverse Reactions from Contrast Media in Patients Undergone Computed Tomography at the Department of Radiology, Srinagarind Hospital

Pranee Suecharoen, Jaturat Kanpittaya

**Abstract**—Background: The incidence of adverse reactions to iodinated contrast media has risen. The dearth of reports on reactions to the administration of iso- and low-osmolar contrast media should be addressed. We, therefore, studied the profile of adverse reactions to iodinated contrast media; viz., (a) the body systems affected (b) causality, (c) severity, and (d) preventability. Objective: To study adverse reactions (causes and severity) to iodinated contrast media at Srinagarind Hospital. Method: Between March and July, 2015, 1,101 patients from the Department of Radiology were observed and interviewed for the occurrence of adverse reactions. The patients were classified per Naranjo's algorithm and through use of an adverse reactions questionnaire. Results: A total of 105 cases (9.5%) reported adverse reactions (57% male; 43% female); among whom 2% were iso-osmolar vs. 98% low-osmolar. Diagnoses included hepatoma and cholangiocarcinoma (24.8%), colorectal cancer (9.5%), breast cancer (5.7%), cervical cancer (3.8%), lung cancer (2.9%), bone cancer (1.9%), and others (51.5%). Underlying diseases included hypertension and diabetes mellitus type 2. Mild, moderate, and severe adverse reactions accounted for 92, 5 and 3%, respectively. The respective groups of escalating symptoms included (a) mild urticaria, itching, rash, nausea, vomiting, dizziness, and headache; (b) moderate hypertension, hypotension, dyspnea, tachycardia and bronchospasm; and (c) severe laryngeal edema, profound hypotension, and convulsions. All reactions could be anticipated per Naranjo's algorithm. Conclusion: Mild to moderate adverse reactions to low-osmolar contrast media were most common and these occurred immediately after administration. For patient safety and better outcomes, improving the identification of patients likely to have an adverse reaction is essential.

**Keywords**—Adverse reactions, contrast media, computed tomography, iodinated contrast agents.

## I. INTRODUCTION

THE number of people receiving services at Srinagarind Hospital has increased annually, so that by 2013 it served 850,000 patients: a rise of 45,000 from the previous year [1]. Consequently, the hospital has had to improve its preparedness in all medical disciplines; particularly vis-à-vis rapid, accurate, modern diagnostic radiology so as to quickly diagnose the large number of incoming patients.

Currently, radiologic diagnosis with modalities (contrast media) plays a major role in modern medicine. Computed tomography (CT Scan) employs multiple X-ray slices, that

Pranee Suecharoen is with the Department of Pharmacy, Srinagarind Hospital, Faculty of Medicine, Khon Kaen 40002 (e-mail: pharmpupu@gmail.com).

Jaturat Kanpittaya is with the Department of Diagnostic Radiology, Srinagarind Hospital, Faculty of Medicine, Khon Kaen 40002.

pass through the body in various planes, creating 3D, cross-sectional, computer-generated images of the desired organ that can be viewed from various angles on screen. These images are used for assessing anatomy, detecting abnormalities, and diagnosing disease (including congenital disorders) and determining the status of infectious disorders, vascular tumors, and accidents. In conjunction with injection modalities (using iodine compounds), CT is invaluable for: (a) Detecting tumors in various organs: (b) determining tumor spread to lymph nodes: (c) assessing retention of blood in the brain, abdomen, and pelvis: (d) detecting coronary artery aneurysms and embolisms: and, (e) assessing bone disorders including, arthritis or fractures. Owing to these multiple uses and evident benefits, the respective number of patients receiving CT Scan in 1998 and 1999 was 6,645 and 7,680, which rose to 9,953 in 2012 [1].

Modalities using iodine (i.e., iodinated contrast media) help to visualize kidneys, blood vessels, gall bladder, bile ducts, urinary tract, and lymph nodes, but careful consideration must be given to the dosages given as the contrast media are delivered directly into the body (i.e., orally, rectally, intravenously, or through a colostrom line), and there can be adverse reactions.

Idiosyncratic or general- reactions occur within 20 minutes after injection [2] (i.e., anaphylactic shock). An immunoglobulin E (IgE) adverse reaction has non-specific symptoms but these directly affect organs (including heart rate and low blood pressure). Since modalities can exacerbate the severity of disease, people with medical conditions (i.e., cardiac arrhythmias and ischemic) are at risk. The deterioration of the kidney due to the adverse non-specific effects from contrast modalities (i.e., contrast induced nephropathy, CIN) can be categorized into 3 levels (mild, moderate, and serious). Mild symptoms include rash, allergic reactions, distributed generally, and followed by itching, nausea, vomiting, sweating, coughing and dizziness. Moderate symptoms include rash all over the body, facial swelling, bronchoconstriction, tachycardia, palpitations, wheezing, abdominal pain, abdominal cramp requiring diagnosis and immediate treatment. Serious symptoms include severe cardiac arrhythmias, low blood pressure, bronchospasm, severe throat swelling, pneumonia, seizures, coma and possibly death. This category could also have non-idiosyncratic reactions and organ-specific types of adverse reactions caused by a direct effect on cell structure. All patients should be closely monitored.

One of the mechanisms of actions from contrast media may be the results of enzymes which stimulate the complementary system while other causes have different potential from mild to life-threatening that affect organs (i.e., bradycardia, hypotension). Vasovagal reactions induce the parasympathetic system and may be associated with symptoms of the autonomic nervous system including nausea, vomiting, diarrhea, sweating, confusion, a reduction in the rate of discharge from the sino-atrial node and atrial-ventricular node making the heart beat slower. The peripheral expansion causes low blood pressure, cardiovascular reactions, including severe hypotension and bradycardia. These symptoms are caused by compounds that act on the heart and blood vessels caused by anaphylactoid reaction from iodinated contrast media.

Accidental extravascular or contrast extravasation is caused by contrast media injection leakage from the vessel puncture site or leaks into the tissue around vein. Contrast extravasation is a well-recognized complication; the respective reported frequencies where power injectors were used in three large CT series was 0.25% (56/22,254), 0.7% (475/69,657), and 0.9% (48/5,106) [3]-[5].

In the current study, reactions were not regarded adverse reactions from contrast extravasation as they occurred after administration of the contrast media. Nueaithong reviewed 2,176 patients injected with contrast media for the diagnosis of special urology and found 9.3 percent of the patients had adverse reactions: 8.9% mild, 0.4% moderate, and no severe adverse reactions [6]. In another study by a Japanese committee on the safety of contrast media, of 337,647 patients, 0.02% had serious reactions, while 0.04% had severe reactions to ionic contrast media while 0.004% had severe reactions to nonionic contrast media [7]. Few studies provide information on the operating result (i.e. the intensity and duration of the adverse reactions and the body systems involved).

Since the 1990s, iodinated contrast media have been used mostly in case of unclear diagnosis [2]-[4]. The lack of reports on reactions to the administration of iso- and low-osmolar contrast media needs to be addressed in the interests of patient safety and cost to radiologic services. It is important to (a) quantify adverse reactions when iodine is used as an ingredient, and (b) find ways to prevent and reduce this adverse reaction. We, therefore, studied the profile of adverse reactions to iodinated contrast media: (a) body systems affected; (b) causality; (c) severity; and, (d) preventability.

## II. METHOD

The current study was designed to determine the incidence and severity of adverse reactions due to the contrast media used during computed tomography at the Department of Radiology, Srinagarind Hospital, Khon Kaen University between March and July, 2015. Data collection was based on inclusion criteria: 1,101 patients diagnosed using CT scan and their major features are observed and recorded by a nurse. All patients were classified per Naranjo's algorithm as (i.e. serious, moderate, or mild) after which a pharmacist completed a questionnaire on their symptoms and adverse

reaction. SPSS version 17 was used to complete the statistical analyses.

## III. RESULTS

TABLE I  
CHARACTERISTICS OF CONTRAST MEDIA ADVERSE EVENT REPORTS

Characteristic	Total of reports (1101)		Adverse reports (105)	
	amount	percentage	amount	percentage
Sex				
male	623	57	49	47
female	478	43	56	53
Age (year, mean±SD)	55.5±15.8		54.7±11.6	
20-30	28	2.54	5	4.8
31-40	98	8.9	10	9.5
41-50	140	12.7	22	21
51-60	399	36.24	35	33.34
61-70	295	26.9	28	26.7
71 up	141	12.8	5	4.8
Medical history				
Antidiabetic drugs	385	35	40	40
Antihypertensive drugs	227	20.6	30	30
Cardiovascular drugs	195	17.7	8	8
Antithyroid drugs	98	8.9	3	3
Antiepileptic drugs	98	8.9	3	3
Antiviral drugs	98	8.9	16	16
Diagnosis				
Hepatoma and Cholangiocarcinoma	358	32.5	26	25
Colorectal cancer	143	13	10	9.5
Breast cancer	45	4.1	6	5.7
Cervical cancer	89	8.1	4	3.8
Lung cancer	144	13.1	3	2.9
Bone cancer	5	0.45	2	1.9
Abdominal and stomach cancer	185	16.80	0	0
Brain cancer	34	3.1	0	0
Gallbladder cancer	24	2.2	0	0
Cardiovascular disease	18	1.64	0	0
Kidney cancer	32	2.9	0	0
Others	24	2.2	54	51.5

A total of 1,101 patients were included in the study: 623 (57%) males and 478 (43%) females. Adverse reactions were reported by 105 cases (9.5%). The mean age ± SD of all patients was 55.5±15.8 vs. 54.7±11.6 for those who developed adverse reactions. Most of those having an adverse reaction from contrast media were between 51 and 60 years of age (33.34%). The other respective age ranges (and proportion suffering adverse reactions) were 61-70 (26.7%), 41-50 (21%), 31-40 (9.5%), 20-30 (4.8%), and 70 and over (4.8%). The medical histories indicated the rank of medications being taken was: Anti-diabetic medications, anti-hypertensives, cardiovascular medications, anti-hyper/hypothyroid medications, anti-epileptics, and anti-virals. The rank of diagnosed diseases included: Hepatoma and cholangiocarcinoma, abdominal and stomach cancer, lung cancer, colorectal cancer, cervical cancer, breast cancer, brain cancer, kidney cancer, gallbladder cancer, cardiovascular disease, bone cancer, and others. The rank of diseases associated with an adverse reaction included: Hepatoma and

cholangiocarcinoma (24.8%), colorectal cancer (9.5%), breast cancer (5.7%), cervical cancer (3.8%), lung cancer (2.9%), bone cancer (1.9%), and others (51.5%). There was no significant difference in the occurrence/non-occurrence of adverse reactions to iodinated contrast media due to sex, mean age, or medical history (Table I).

Among the 1,101 cases who received contrast media, 105 had adverse reactions. Among those who with an adverse reaction from contrast media for diagnosis, 2% were iso-osmolar and 98% low-osmolar. In the current study, the respective rank of those having majority who had an adverse reaction from iodinated contrast media using low-osmolarity for diagnosis were iopromide (31%), iohexol (26%), iopamidol (21%), iobitridol (20%), and iso-osmolarity as iodixanol (2%), (Table II).

TABLE II  
FREQUENCY OF REPORTS USING IODINATED CONTRAST MEDIA WITH AN ADVERSE REACTION

Class	Contrast media	Total of cases (1101)		Adverse reports (105)	
		amount	percentage	amount	percentage
Low-osmolarity	Iobitridol	214	19.4	21	20.0
	Iohexol	327	29.7	27	25.7
	Iopromide	278	25.24	33	31.4
	Iopamidol	257	23.4	22	21.0
Iso-osmolar	Iodixanol	25	2.3	2	1.9

105 of 1,101 patient's reactions were classified using Naranjo's algorithm for an adverse reaction with contrast media; then, they were assessed per their symptoms using specific questionnaire administered by a pharmacist. Their symptoms were assessed; all reactions were deemed to be probable. The respective incidence of mild, moderate and severe adverse reactions accounted was 92, 5, and 3%. The most common adverse reaction after iodinated contrast media administration was immediate; occurring within an hour (both mild and severe). Incidences of delayed adverse reactions (occurring hours to weeks after the administration of ionated contrast media) are classified in (Table III). These reactions are usually self-limiting and cutaneous though they can be classified as moderate adverse reactions.

TABLE III  
INCIDENCE OF ADVERSE REACTIONS TO NON-IODINATED CONTRAST MEDIA

Classification	Amount (105 cases)	Percentage
Mild	97	92
Moderate	5	5
Severe	3	3
Total	105	100

The respective groups of escalating symptoms included: mild—urticaria, itching, rash, nausea, vomiting, dizziness, and headache; moderate—hypertension, hypotension, dyspnea, tachycardia and bronchospasm; and, severe—laryngeal edema, profound hypotension, and convulsions (Table IV). The most common adverse reactions included: Cutaneous urticaria, itching, and immediate MP rash. A few cases had nausea/vomiting, dizziness, and headache. Among the severe

reactions, symptoms included laryngeal edema, profound hypotension, and convulsions, but the incidence was low compared to mild adverse reactions.

TABLE IV  
SYMPTOMS OF AN ADVERSE REACTION FROM NON-IODINATED CONTRAST MEDIA

Classification	Symptoms*	Amount	Percentage
Mild	Urticaria	82	78.1
	Itching	19	18.1
	MP rash	18	17.14
	Nausea / Vomiting	2	2.9
	Dizziness	1	≤1.0
	Headache	1	≤1.0
Moderate	Hypertension	2	1.9
	Hypotension	1	≤1.0
	Dyspnea	1	≤1.0
	Tachycardia	1	≤1.0
	Bronchospasm	1	≤1.0
	Laryngeal edema	2	1.9
Severe	Profound hypotension	1	≤1.0
	Convulsions	2	1.9

\* Patient could have more than one symptom

#### IV. DISCUSSION

The present study comprised out-patients from the Radiology Department of Srinagarind Hospital. We focused on patients who developed an adverse reaction due to administration of non-iodinated contrast media. All other reports of adverse reactions from contrast media were due to non-iodinated contrast media over against the 9.5% in our study. Several studies reported the prevalence of adverse reactions from both iodinated and non-iodinated contrast media [8]-[12]. Reports on the characteristics of patients after contrast media administration with an adverse event indicate no significant difference among adverse reactions vis-à-vis particularly sex and age. In an Indian study, neither sex nor age yielded any significant association with development of adverse reactions [10].

In the current study, adverse reactions were classified into 3 types (i.e., mild, moderate, and serious). The most common adverse reactions were mild (i.e. urticaria, itching, MP rash, nausea, vomiting, dizziness, and headache) and these developed within 30 minutes of administration; these were also classified as immediate. Some studies of low osmolarity contrast media [6]-[11] found that less than 3% of patients had reactions which were immediate.

The reported reactions include mild reactions such as urticaria, itching, MP rash, and nausea. Moderate and severe reactions include bronchospasm, cardiac arrhythmia, and loss of consciousness. Mortality due to contrast media reactions is low or ≤1 one death per 100,000 patients [7], [13], [14]. Incidence of immediate reactions to nonionic contrast media ranged from mild (i.e., 3%) to severe (i.e., 0.01–0.04%) [12]. Non-ionic agents are associated with a decreased risk of adverse reactions [11]. The prevalence of adverse reaction with ionic contrast media was 12.7% compared to 3.13% when using non-ionic contrast media [2], [6], [12]. By

comparison, high-osmolality contrast media have 5-8 times the osmolality of plasma. At present, both low-osmolality and iso-osmolality are used in diagnosis increasingly. The incidences of mild and moderate contrast media reactions are higher for high osmolality contrast media (6-8%) than low osmolality contrast media 0.2% [14], [15]. Similarly, reactions are infrequent and range between 5% and 12% for high osmolality and low osmolality (1-3%) for contrast media [15], [16].

Although, our study used non-iodinated contrast media, the most common reactions were mild albeit immediate reactions are akin to hypersensitivity reactions; including both E and non-Ig E-mediated anaphylaxis, with activation of mast cells, coagulation, kinin and complementary mechanisms, (i.e. inhibition of enzymes, and platelet aggregation) [17]. As for severe adverse reactions, one case was given iopamidol and 2 cases were given iopromide. The symptoms included anaphylactic shock, severe hypotension, confusion, dyspepsia, desaturation and generalized urticaria [17]-[19]. Finally, anti-histamines are initially used to manage common reactions from iodinated contrast media in mild reactions. Severe reactions [20], [21] are treated by guidelines and a hospital multidisciplinary team.

#### V.CONCLUSION

Mild to moderate adverse reactions to low-osmolality contrast media were the most common and these occurred immediately after administration of the media. Improving the identification of patients likely to have an adverse reaction is critical to avoiding life-threatening outcomes.

#### REFERENCES

- [1] Department of Radiology, Srinagarind Hospital. Summary statistic of contrast media in Srinagarind Hospital January 2008 – December 2014. Khon Kaen: Faculty of Medicine, Khon Kaen University; 2015.
- [2] Neuaithong K. Prevention guidelines for reducing adverse reaction from intravenous administration of Iodinated Contrast Medium (ICM). *Srinagarind Med J.* 2009;24(1):91–101.
- [3] Cohan RH, Bullard MA, Ellis JH, Jan SC, Francis IR, Garner WL, et al. Local reactions after injection of iodinated contrast material: detection, management, and outcome. *Acad Radiol.* 1997;4(11):711–8.
- [4] Wang CL, Cohan RH, Ellis JH, Adusumilli S, Dunnick NR. Frequency, management, and outcome of extravasation of nonionic iodinated contrast medium in 69,657 intravenous injections. *Radiology.* 2007;243(1):80–7.
- [5] Federle MP, Chang PJ, Confer S, Ozgun B. Frequency and effects of extravasation of ionic and nonionic CT contrast media during rapid bolus injection. *Radiology.* 1998;206(3):637–40.
- [6] Neuaithong K. Adverse Reaction from intravenous administration of Iodinated Contrast Material in patient undergoing intravenous urography. *Srinagarind Med J.* 2012;27(3):288–98.
- [7] Katayama H, Yamaguchi K, Kozuka T, Takashima T, Seez P, Matsuura K. Adverse reactions to ionic and nonionic contrast media. A report from the Japanese Committee on the Safety of Contrast Media. *Radiology.* 1990;175(3):621–8.
- [8] Seong J-M, Choi N-K, Lee J, Chang Y, Kim Y-J, Yang BR, et al. Comparison of the safety of seven iodinated contrast media. *J Korean Med Sci.* 2013;28(12):1703–10.
- [9] Subathra A, Sandhiya S, Kesavan R. An analysis of adverse drug reactions to radiographic contrast media reported during a 3 year period in a tertiary care hospital in south India. *Indian J Physiol Pharmacol.* 2014;58(1):45–50.
- [10] Modi N, Vaidya K, Dudhia S, Shah R, others. A study of adverse drug reactions to radio contrast media in a tertiary care teaching rural hospital. *Int J Med Sci Public Health.* 2014;3(2):133–6.
- [11] Bhowmick S, Bhat E, Panja B, Mukherjee S, Sikdar S, Biswas A, et al. A study on adverse drug reactions to non-ionic contrast medium in an Indian population: a 1-year experience. *Int J Basic Clin Pharmacol.* 2014;3:1066–71.
- [12] Namasivayam S, Kalra MK, Torres WE, Small WC. Adverse reactions to intravenous iodinated contrast media: a primer for radiologists. *Emerg Radiol.* 2006;12(5):210–5.
- [13] Caro JJ, Trindade E, McGregor M. The risks of death and of severe nonfatal reactions with high- vs low-osmolality contrast media: a meta-analysis. *AJR Am J Roentgenol.* 1991;156(4):825–32.
- [14] Cochran ST, Bomyea K, Sayre JW. Trends in adverse events after IV administration of contrast media. *AJR Am J Roentgenol.* 2000;176(6):1385–8.
- [15] Bush WH, Swanson DP. Acute reactions to intravascular contrast media: types, risk factors, recognition, and specific treatment. *AJR Am J Roentgenol.* 1991;157(6):1153–61.
- [16] Thomson KR, Varma DK. Safe use of radiographic contrast media. *Aust Prescr.* 2010;33(1):19–22.
- [17] Cutroneo P, Polimeni G, Curcuruto R, Calapai G, Caputi AP. Adverse reactions to contrast media: an analysis from spontaneous reporting data. *Pharmacol Res.* 2007;56(1):35–41.
- [18] Singh J, Daftary A. Iodinated contrast media and their adverse reactions. *J Nucl Med Technol.* 2008;36(2):69-74-; quiz 76-7
- [19] Zhang B, Dong Y, Liang L, Lian Z, Liu J, Luo X, et al. The incidence, classification, and management of acute adverse reactions to the low-osmolar iodinated contrastmedia isovue and ultravist in contrast-enhanced computed tomography scanning. *Medicine.* 2016;95(12):1-9.
- [20] Maddox T G. Adverse reactions to contrast material: recognition, prevention, and treatment. *Missouri: Am Fam Physician.* 2002;66(7):1229-34.
- [21] Tramèr MR, von Elm E, Loubeyre P, Hauser C. Pharmacological prevention of serious anaphylactic reactions due to iodinated contrast media: systematic review. *BMJ.* 2006;333(7570):675-80.