# Influence of Organic Supplements on Shoot Multiplication Efficiency of *Phaius tankervilleae* var. *alba*

T. Punjansing, M. Nakkuntod, S. Homchan, P. Inthima, A. Kongbangkerd

Abstract-The influence of organic supplements on growth and multiplication efficiency of Phaius tankervilleae var. alba seedlings was investigated. 12 week-old seedlings were cultured on halfstrength semi-solid Murashige and Skoog (MS) medium supplemented with 30 g/L sucrose, 8 g/L agar and various concentrations of coconut water (0, 50, 100, 150 and 200 mL/L) combined with potato extract (0, 25 and 50 g/L) and the pH was adjusted to 5.8 prior to autoclaving. The cultures were then kept under constant photoperiod (16 h light: 8 h dark) at 25 ± 2 °C for 12 weeks. The highest number of shoots (3.0 shoots/explant) was obtained when cultured on the medium added with 50 ml/L coconut water and 50 g/L potato extract whereas the highest number of leaves (5.9 leaves/explant) and roots (6.1 roots/explant) could receive on the medium supplemented with 150 ml/L coconut water and 50 g/L potato extract. with 150 ml/L coconut water and 50 g/L potato extract. Additionally, plantlets of P. tankervilleae var. alba were transferred to grow into seven different substrates i.e. soil, sand, coconut husk chip, soil-sand mix (1: 1), soil-coconut husk chip mix (1: 1), sand-coconut husk chip mix (1: 1) and soil-sand-coconut husk chip mix (1: 1: 1) for four weeks. The results found that acclimatized plants showed 100% of survivals when sand, coconut husk chip and sand-coconut husk chip mix are used as substrates. The number of leaves induced by sand-coconut husk chip mix was significantly higher than that planted in other substrates (P > 0.05). Meanwhile, no significant difference in new shoot formation among these substrates was observed (P < 0.05). This precursory developing protocol was likely to be applied for more large scale of plant production as well as conservation of germplasm of this orchid species.

*Keywords*—Acclimatization, coconut water, orchid, *Phaius tankervilleae* var. *alba.*, potato extract.

### I. INTRODUCTION

THE *Phaius* is a genus of terrestrial orchids belonging to Collabieae tribe of Epidendroideae subfamily. Less than 50 described species were recorded and widely distributed in Africa, Australia, Pacific and Indian Ocean islands and Asia including Thailand [1]. Up to date, six known species of *Phaius* i.e. *P. tankervilleae*, *P. mishmensis*, *P. indigoferus*, *P.* 

T. Punjansing is Ph.D. student in Plant Tissue Culture Research Unit, Department of Biology, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand.

M. Nakkuntod and S. Homchan are with the Department of Biology, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand.

P. Inthima and A. Kongbangkerd are with the Plant Tissue Culture Research Unit, Department of Biology, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand (e-mail:anupank@nu.ac.th).

A. Kongbangkerd is also take part in Center of Excellence in Research for Agricultural Biotechnology, Faculty of Agriculture, Natural Resources and Environment, Naresuan University, Phitsanulok 65000, Thailand. takeoi, P. flavus and P. epiphticus have been morphologically classified and exhibited throughout Thailand [2]. Among them, P. tankervilleae is one of the most ornamentally attractive species. Morphologically, their inflorescences are large, showy and distinctive. They flower easily and have a long duration of blooming which is appropriate to use as potted plants and cut flower [3], [4]. In addition, some countries have applied P. tankervilleae as folk medicines to treat rash, freckle, blemish and blister [5]. This orchid species in natural habitat is stated as rare and endangered in Japan [6], India [7] and Australia [8] and is now listed in CITES Appendix II which somehow can be traded under a strict regulation [9]. The extinction is much more concerned with P. tankervilleae var. alba which is rarely found only in few natural population (Fig. 1 (a)). Therefore, an efficient method for rapid propagation of this variety is needed to study for conservation and sustainable utilization. In vitro culture technique is an effective way of propagation and has been successful in propagating many orchid species [10].

The composition of culture medium is one of the key factors affecting the result of *in vitro* orchid culture. Growth and multiplication rate of many orchid species can be enhanced by adding some organic supplements like coconut water (CW) or potato extract (PE) to culture medium [11], [12]. However, the accomplishments are depended on explants type and orchid species [13]. At present, only few publications [14], [15] study about the effect of organic supplements on shoot regeneration capability in the *Phaius* species especially in *P. tankervilleae* var. *alba*. Therefore, this study attempted to observe the influence of organic supplements on shoot multiplications efficiency of *P. tankervilleae* var. *alba*.



Fig. 1 Flowers (a) and pod (b) of P. tankervilleae var. alba

50 g/L).

### II. MATERIALS AND METHODS

## A. In vitro Culture of P. tankervilleae var. alba

The pod of P. tankervilleae var. alba (Fig. 1 (b)) obtained from Romklao Botanical Garden under the Royal Initiative, Phitsanulok, Thailand was sterilized with 15% (v/v) Clorox® solution for 20 minutes. The seeds were collected from sterilized pod and cultured on modified VW [16] medium supplemented with 150 mL/L CW and 50 g/L PE. The media was solidified by 8 g/L agar and pH was adjusted to 5.2. After 12 weeks of seed culture, the seedlings were cultured on halfstrength MS medium [17] supplemented with different concentrations of CW and PE. The media were added with 15 g/L sucrose, 8 g/L agar and adjusted pH to 5.8. The seedlings were cultivated under cool-white LED lamp (20  $\mu$ mol/m<sup>2</sup>/s) for 16 h/day at 25 ± 2 °C. Growth and development of seedlings were measured and recorded after 12 weeks of culture. Each treatment was performed in 10 replicates and repeated 3 times. The statistical differences of data were analyzed by One-way ANOVA and Duncan's multiple range tests (DMRT).

## B. Acclimatization of P. tankervilleae var. alba Plantlets

The *in vitro* rooted plantlets were washed with sterile water to remove the agar medium and immersed in fungicide solution for 20 min. The treated plantlets were then transplanted to plastic pot (5 cm wide and 5 cm height) containing different planting substrates and cultured under greenhouse (28–34 °C air temperature, relative 50–60% humidity and natural sunlight with 70% shade nets). After 4 weeks of transplantation, survival and growth of transplanted plants were recorded. The experiment was conducted with 3 replications of 10 plantlets per replication

## III. RESULTS AND DISCUSSION

A. In vitro Growth and Multiplication Efficiency



Fig. 2 *In vitro* plantlets of *P. tankervilleae* var. *alba* 12 weeks after cultured on (a) ½MS medium, (b) ½ MS supplemented with 150 mL/L CW, (c) 50 g/L PE and (d) 50 mL/L CW combined with 50 g/L PE

For *in vitro* orchid culture, CW and PE are extensively used because they effectively promote growth and development of orchids [18], [19]. Therefore, CW and PE were applied in this study. The seedlings were cultured on half-strength semi-solid MS medium supplemented with various concentrations of CW (0, 50, 100, 150 and 200 mL/L) combined with PE (0, 25 and

After 12 weeks of culture, the seedlings were all survived in every tested media. However, diverse growth performances were found among the different PE and CW supplementations (Fig. 2).



Fig. 3 Effect of CW and PE supplementation on number of shoot (a), leaves (b) and root (c) per explant of *P. tankervilleae* var. *alba* after culture for 12 weeks. The bars and error bars represent means and SE of 3 replicates (each with 10 plants), respectively. Different letters above the bar show significant differences analyzed by DMRT at  $p \le 0.05$ 

Different concentrations of CW affected shoot production of seedling randomly (Fig. 3 (a)). At low concentration of CW supplementation (0 and 50 mL/L), shoot number had a tendency to increase with the concentration of PE supplementation. On the other hand, shoot numbers tend to decrease with increasing PE content at higher CW

## International Journal of Biological, Life and Agricultural Sciences ISSN: 2415-6612 Vol:13, No:4, 2019

supplementation (100 and 150 mL/L). Nevertheless, different PE supplementation did not affect shoot number at the highest CW addition (200 mL/L). In summary, the most effective combination for shoot multiplication of *P. tankervilleae* var. *alba* seedling was 50 mL/L CW with 50 g/L PE (Figs. 2 (d) and 3 (a)). Although PE and CW have been used in orchid culture media, combination effect has been observed in only few orchid species [11]. Previously, enhancement growth of *Calanthe* hybrid plantlet by CW supplement has been reported [20]. The optimum concentration was 50 mL/L which was consistent with this study. However, the higher CW concentration promotes better growth and shoot multiplication in some orchid species [11], [21]. So, the effect of CW may be depending on species and explants.

Numbers of leaves of seedling are obviously affected by the supplementation of CW and PE (Fig. 3 (b)). Without PE, adding high concentration (more than 150 mL/L) of CW inhibited leaves production. However, the leave production was obviously improved along with the increasing concentration of PE augmentation in high CW containing media. On the contrary, PE combination did not significantly increase the number of leave at CW supplementation lower than 150 mL/L. The highest average leaves number per explant was achieved from the medium containing 150 mL/L CW and 50 g/L PE. The combination effect of CW and PE on number of leaves was also formerly reported in *Bulbophyllum nipondhii* [11]; however the number of leave did not decrease in high concentration of CW supplementation.

In vitro plantlet with root is preferred before transfer to ex vitro environment. The plants without root or poor root quality hardly survive under ex vitro condition [22]. In this study, roots were successfully produced in every plantlet. Like leaves production, without PE, root was reduced with the increasing of CW concentration (Fig. 3 (c)). Similarly, root was also decreased with the increasing of PE concentration in media without CW. However, PE combinations were significantly improved root production in media holding CW, especially at high CW concentration (more than 150 mL/L). Therefore, root productions appear to be induced by the combination effect of CW and PE. This effect was aforementioned in *Bulbophyllum nipondhii* [11].

The main composition of CW is sugars, amino acids, minerals, vitamins and particularly plant hormones like auxin and cytokinin [23] which has been proposed as the key ingredient affecting explant organogenesis, growth and development. For PE, although the main composition is carbohydrate, protein, and vitamin [25], the important composition influenced plant growth and development was still doubtful. Previously, it was proposed that organic nitrogen was easier to process by orchid protocorms in comparison to inorganic nitrogen because it may by pass some step in nitrogen metabolic pathways [25]. Thus, the improvement of shoot, leaves and root number of P. tankervilleae var. alba probably due to the action of some phytohormones, organic nitrogen compounds, phosphorus, or vitamins from both CW and PE. In summary, the most suitable culture medium for in vitro shoot multiplication of P.

*tankervilleae* var. *alba* was half-strength MS supplemented with 50 mL/L CW and 50 g/L PE. This medium provides the highest shoot number with fair leaves and roots number.

B. Survival and Growth of Acclimatized Plant



Fig. 6 Growth of *P. tankervilleae* var. *alba* planting with soil (a), sand (b), coconut husk chip (c), soil-sand mix, 1: 1 (d), soil-coconut husk chip mix, 1: 1 (e), sand-coconut husk chip mix, 1: 1 (f), coconut husk chip mix, 1: 1: 1 (g). The pictures were taken after 4 weeks of greenhouse acclimatization

Acclimatization is an important step for in vitro plant propagation. This process enhances the survival of in vitro plantlet after transfer to ex vitro environment [26]. The in vitro plantlets of P. tankervilleae var. alba were successfully transplanted to greenhouse and grew healthily (Fig. 6). Up to 90% of them were survived in all tested substrates, however there were no statistically significant different among the tested substrates (Table I) although survival variant between different substrates have been previously observed in some orchids [27], [28]. This may be due to all substrates were able to keep the minimum requirement of ground moisture for P. tankervilleae var. alba plants. Nonetheless, different growth of acclimatized plant was observed among the tested substrates. Highest leaves number was obtained from the plants culturing with the sand-coconut husk chip mix (Fig. 6 (f)). The longest leaf length was gained from the plants growing with the soilsand mix (Fig. 6 (f)) while the broadest leaf width was obtain

## International Journal of Biological, Life and Agricultural Sciences ISSN: 2415-6612 Vol:13, No:4, 2019

form soil-grown plants (Fig. 6 (a)). However, regarding the overall performances, sand-coconut husk chip mix seem to be the suitable substrate for P. *tankervilleae* var. *alba* acclimatization.

### IV. CONCLUSIONS

This research presents the first protocol for *in vitro* shoot multiplication of *P. tankervilleae* var. *alba*. The most suitable

culture medium was half-strength MS supplemented with 50 mL/L CW and 50 g/L PE. The *in vitro* plantlets were successfully acclimatized and survived under the greenhouse. The mixture of sand and coconut husk ship (1: 1) was the most appropriate substrate for acclimatization. This study provides the useful information applicable for commercial propagation and conservation.

TABLE I
EFFECT OF PLANTING SUBSTRATES ON ACCLIMATIZATION OF P. TANKERVILLEAE VAR. ALBA IN GREENHOUSE FOR 4 WEEKS

	Substrates	Survival (%)	No. of leaves per plant	Leaf size (cm) <sup>1</sup>		New shoot
				Length	Width	Formation (%)
_	Soil	$92.5\pm15.0\ ns$	$2.6\pm1.0\ d$	$5.5\pm2.4\ abc$	$1.8\pm3.0\;a$	$2.4\pm4.0\ ns$
	Sand	$100.0\pm0.0$	$3.6\pm1.1\ b$	$5.6 \pm 1.9 \text{ abc}$	$0.7\pm0.2\;b$	$1.7\pm2.8$
	Coconut husk chip	$100.0\pm0.0$	$3.3\pm0.8\ bc$	$5.5\pm2.0\ bc$	$0.9\pm0.2\;b$	$1.9\pm3.2$
	Soil + Sand $(1: 1)$	$87.5\pm5.0$	$3.3\pm0.9\ bc$	$6.8\pm2.4~a$	$0.8\pm0.2\;b$	$2.4\pm4.0$
	Soil + Coconut husk chip (1: 1)	$92.5\pm15.0$	$2.7\pm0.5\ d$	$6.6 \pm 1.9 \text{ ab}$	$1.0\pm1.0\ b$	$0.0\pm0.0$
	Sand + Coconut husk chip (1: 1)	$100.0\pm0.0$	$4.2\pm1.0\ a$	$4.9\pm1.8\ c$	$0.8\pm0.3\;b$	$0.0\pm0.0$
	Soil + Sand + Coconut husk chip (1: 1: 1)	$95.5\pm2.8$	$3.0\pm1.0\ cd$	$6.6\pm2.6\;ab$	$0.8\pm0.2\;b$	$6.6\pm7.2$

Values are means  $\pm$  SE of 3 replications (10 plants per replication). Different letters within the same column show significant differences analyzed by DMRT at  $p \le 0.05$ . ns = not significant. <sup>1</sup> Data were obtained from the 3<sup>rd</sup> leaf of each plant.

## ACKNOWLEDGMENT

This work was financial supported by Capacity Building Program for New Researcher 2018, National Research Council of Thailand (NRCT). The authors are grateful to Romklao Botanical Garden under the Royal Initiative, Phitsanulok, Thailand for plant materials. T. Punjansing offers thanks to Udon Thani Rajabhat University, Thailand for scholarship and to Professor Dr. Duncan R. Smith (Institute of Molecular Biosciences, Mahidol University, Thailand) for English editing and scientific proofreading of this manuscript.

#### REFERENCES

- [1] R. L. Dressler, *Phylogeny and classification of orchid family*. Portland, OR: Dioscorides, 1993, ch1.
- [2] H. Kurzweil, "A precursory study of the Calanthe group Orchidaceae in Thailand," Adasonia ser, vol. 3, no. 1, pp. 92, 2010.
- [3] L.V. Averyanov and E.A. Christenson, "Orchids of Vietnam. *Phaius tankervilleae*," Orchids. *Mag. Amer. Orch. Soc.*, vol. 67, no 4, pp. 368-370, 1998.
- [4] R Ngapui, P. Gogoi, K. Chowlu, and S. P. Vij, "Effects of NPK and 6-Benzylaminopurine on Growth and Flowering of Two Orchid Genera," *Pl sci Res*, vol. 3, no 1, pp. 93-99, 2018.
- [5] M. M. Hossain, "Traditional therapeutic uses of some orchids of Bangladesh," *Medicinal Aromatic Plant Sci. Biotech.*, vol.3, no1, pp.100-106, 2009.
- [6] T. Hirano, T. Godo, K. Miyoshi, K. Ishikawa M. Ishikawa and M. Mii, "Cryopreservation and low temperature storage of seeds of *Phaius tankervilleae*," *Plant Biotechnol Rep.*, vol.3, pp. 103-109, 2009.
- [7] R. Thokchom, S. Maitra and S. Sharma, "In vitro Mass Propagation of Endangered Terrestrial Orchid - Phaius tankervilleae (L'Her.) Blume through Green Seed Pod Culture," Microbiol. App. Sci., vol. 6, no.5, pp.722-728, 2017.
- [8] D. J. Metcalfe and T.J. Lawson, "An International Union for Conservation of Nature risk assessment of coastal lowland rainforests of the Wet Tropics Bioregion, Queensland," *Austral Ecology.*, vol. 40, no. 4, pp.373-385, 2015.
- [9] UNEP-WCMC (Comps.), Checklist of CITES species. CITES Secretariat, Geneva, Switzerland, and UNEP-WCMC, Cambridge, United Kingdom. Accessed on September 15, 2018. <u>http://checklist.cites.org.</u> 2014.
- [10] S. Chugh, S. Guha, and I. U. Rao, "Micropropagation of orchids: A

review on the potential of different explants," *Sci. Hortic.*, vol. 122, no. 4, pp. 507-520, 2009.

- [11] W. Pakum, S. Watthana, K. Srimuang, and A. Kongbangkerd, "Influence of Medium Component on *In vitro* Propagation of Thais Endangered Orchid: *Bulbophyllum nipondhii* Seidenf," *Plant Tissue Cult. & Biotech.*, vol. 26, no.1, pp.37-46, 2016.
- [12] B. Chen, S. J. Trueman, L. Jianmin, L. Qianzhen, H. Fan and J. Zhang, "Micropropagation of the endangered medicinal orchid, *Dendrobium* officinale," *Life Sci.*, vol.11, no.9, pp.526-530, 2014.
- [13] Z. Molnár, E. Virág and V. Ördög, "Natural substances in tissue culture media of higher plants,". Acta Biol Szeged., vol.55, no.1, pp. 123-127, 2011.
- [14] B. Pant, S. Shrestha and S. W. Pradhan, "*In Vitro* seed Germination and Seeding Development of *Phaius tancarvillere* (L' Her.) Blume. *Scientific World*, vol. 9, pp. 50 – 52, 2011.
  [15] B. Pant and S. Shrestha "*In Vitro* Mass Propagation of a Ground Orchid
- [15] B. Pant and S. Shrestha "In Vitro Mass Propagation of a Ground Orchid Phaius tancarvillere (L' Her.) Blume through Shoot Tip Culture. Plant Tissue Cult. & Biotech, vol. 21, pp. 181-188, 2011.
- [16] F. Vacin, and F. W. Went, "some pH changes in nutrient solution." Botanic Gardens Conservation News., vol. 110, pp. 605-613, 1949.
- [17] T. Murashige and F. Skoog, "A revised medium for rapid growth and bioassay with tobacco tissue cultures," *Physiol Plant*, vol.15, no. 3, pp. 473 - 497, 1962.
- [18] Z. Molnár, E Virág, and V Ördög "Natural substances in tissue culture media of higher plants". Acta Biol Szeged, vol.55, no1, pp.123-127, 2011.
- [19] T.A. Thorpe, C. Stasolla, E.C. Yeung, G.J. de Klerk, A. Roberts and E.F. George, "The components of plant tissue culture media II: Organic addition, osmotic and pH effects, and support systems, *Springer-Verlag*, 3rd ed, vol.1, pp. 115-173, 2008.
- [20] M. A. Baque, Y.K. Shin, T. Elshmari, E.J. Lee and K.Y. Paek, "Effect of light quality, sucrose and coconut water concentration on the microporpagation of Calanthe hybrids 'Bukduseong' × 'Hyesung' and 'Chunkwang' × 'Hyesung')", Aust J Crop Sci vol. 5, no10, pp.1247-1254, 2011.
- [21] D. Puchooa, "Comparison of different culture media for the *in vitro* culture of *Dendrobium* (orchidaceae)". *Int J Agri Biol*, vol. 6, no.5, pp.884-888, 2004.
- [22] S. Dohling, S. Kumaria and P. Tandon, "Multiple shoot induction from axillary bud culture of the medicinal orchid, *Dendrobium longicornu*"., *AoB Plants*, pp.1-7, 2012.
- [23] JWH Yong, L. Ge, Y. F. Ng and S. N. Tan, "The chemical composition and biological properties of coconut (*Cocos nucifera* L.) Water". Molecules vol.14, no.5, pp.144-5164, 2009
- [24] V. Bartova and J. Barta, "Chemical composition and nutritional value of protein concentrates isolated from potato (Solanum tuberosum L.) fruit

# International Journal of Biological, Life and Agricultural Sciences ISSN: 2415-6612 Vol:13, No:4, 2019

- juice by precipitation with ethanol or ferric chloride". J Agric Food Chem, vol.57, pp.9028-9034, 2009.
  [25] S. Malmgren, "Large-scale asymbiotic propagation of Cypripedium calceolus plant physiology from a surgeons point of view". Botanical Garden Micropropagation News, vol.1, pp.59-63, 1992.
  [26] B. N. Hazarika, "Acclimatization of tissue-cultured plants,". Curr Sci vol, 85, no.12, pp.1704-1712, 2003.
  [27] L. M. Moraes, R. T. Faria and F. L Cuquel, "Activated charcoal for in vitro propagation of Brazilian orchids," Acta Hortic, vol. 683, pp. 383-390, 2005.
  [28] S Dohling, S Kumaria and P Tandon. "Multiple shoot induction from

- [28] S Dohling, S Kumaria and P Tandon, "Multiple shoot induction from axillary bud culture of the medicinal orchid, *Dendrobium longicornu*,". AoB Plants ,vol. 17, 2012.