Instability of a nonlinear differential equation of fifth order with variable delay

Cemil Tunç

Abstract—In this paper, we study the instability of the zero solution to a nonlinear differential equation with variable delay. By using the Lyapunov functional approach, some sufficient conditions for instability of the zero solution are obtained.

Keywords—Instability, Lyapunov-Krasovskii functional, delay differential equation, fifth order.

I. INTRODUCTION

Study on the instability of fifth order non-linear differential equations without delay is not a new topic. Significant results in this direction have been obtained by Ezeilo ([2]-[4]), Li and Duan [7], Li and Yu [8], Sadek [9], Sun and Hou [10], Tiryaki [11], Tunç ([12], [13]), Tunç and Erdogan [19], Tunç and Karta [20] and Tunç and Şevli [21] on instability for some fifth order nonlinear differential equations without delay. Throughout all of these papers, based on Krasovskii's properties (see Krasovskii [6]), the Lyapunov's second (or direct) method has been used as a basic tool to prove the results established therein. This method is one of the powerful and fruitful techniques that has over the years, gained increasing significance in studying qualitative behavior of solutions of differential equations. However, to the best of our knowledge from the literature, an author has considered instability of the solutions of fifth order non-linear differential equations with varying time delays (see Tunç [14]-[18])). Thus, it is worthwhile to continue to investigate the instability of the solutions of fifth order non-linear differential equations with varying time delays in this case.

It is worth mentioning that in 1979, Ezeilo [3] proved an instability theorem to the fifth order nonlinear differential equation without delay

$$x^{(5)} + a_1 x^{(4)} + a_2 x^{\prime\prime\prime} + g(x^{\prime}) x^{\prime\prime} + h(x, x^{\prime}, x^{\prime\prime}, x^{\prime\prime}, x^{(4)}) + f(x) = 0$$

In this paper, instead of Eq. (1), we consider nonlinear differential equation of the fifth order with variable deviating argument $\tau(t)$:

$$x^{(5)} + a_1 x^{(4)} + a_2 x^{\prime\prime\prime} + g(x^{\prime}) x^{\prime\prime} + h(x(t - \tau(t)), x^{\prime}(t - \tau(t)), ..., x^{(4)}(t - \tau(t))) + f(x(t - \tau(t))) = 0.$$
(2)

C. Tunç is with the Department of Mathematics, Faculty of Sciences, Yüzüncü Yıl University, 65080, Van-Turkey, e-mail: cemtunc@yahoo.com Manuscript received xxx; revised xxx. We write Eq. (2) in system form as follows

$$\begin{aligned} x' &= y, \\ y' &= z, \\ z' &= w, \\ w' &= u, \\ u' &= -a_1 u - a_2 w - g(y) z - h(x(t - \tau(t)), ..., u(t - \tau(t))) \\ -f(x) &+ \int_{t - \tau(t)}^{t} f'(x(s)) y(s) ds, \end{aligned}$$
(3)

where a_1 and a_2 are some positive constants, $\tau(t)$ is variable delay, the primes in Eq. (2) denote differentiation with respect to $t, t \in \Re^+ = [0, \infty)$; g, h and f are continuous functions in their arguments on \Re , \Re^5 and \Re , respectively, and with f(0) = 0. The continuity of these functions is a sufficient condition for the existence of the solution of Eq. (2) (see [1], pp.14). It is also assumed that g, h and f satisfy a Lipschitz condition in their respective arguments so that the uniqueness of solutions of Eq. (2) is guaranteed (see [1], pp.15). We assume in what follows that f is also differentiable, and x(t), y(t), z(t), w(t) and u(t) are abbreviated as x, y, z, w and u, respectively.

The motivation for the current paper comes from the works of Ezeilo [3] and Tunç ([14]-[18]). Our results extend and improve the results obtained by Ezeilo [3] for the instability of the zero solution of Eq. (2). Furthermore our result complements existing results on qualitative behavior of solutions of fifth order nonlinear differential equations.

In the following theorem, we give basic idea of the method about the instability of solutions of ordinary differential equations. The following theorem, due to the Russian mathematician N. G. Cetaev's (see LaSalle and Lefschetz [5]).

Theorem A (Instability Theorem of Cetaev's). Let Ω be a neighborhood of the origin. Let there be given a function V(x) and region Ω_1 in Ω with the following properties:

(i) V(x) has continuous first partial derivatives in Ω_1 .

(ii) V(x) and $\dot{V}(x)$ are positive in Ω_1 .

(iii) At the boundary points of Ω_1 inside Ω , V(x) = 0.

(iv) The origin is a boundary point of Ω_1 .

Under these conditions the origin is unstable.

Let $r \ge 0$ be given, and let $C = C([-r, 0], \Re^n)$ with

$$\|\phi\| = \max_{-r \le s \le 0} |\phi(s)|, \phi \in C$$

For H > 0 define $C_H \subset C$ by

$$C_H = \{ \phi \in C : \|\phi\| < H \}.$$

If $x : [-r, A) \to \Re^n$ is continuous, $0 < A \le \infty$, then, for each t in [0, A), x_t in C is defined by

$$x_t(s) = x(t+s), -r \le s \le 0, t \ge 0.$$

Let G be an open subset of C and consider the general autonomous delay differential system with finite delay

$$\dot{x} = F(x_t), x_t = x(t+\theta), -r \le \theta \le 0, t \ge 0$$

where $F: G \to \Re^n$ is a continuous and maps closed and bounded sets into bounded sets. It follows from these conditions on F that each initial value problem

$$\dot{x} = F(x_t), x_0 = \phi \in G$$

has a unique solution defined on some interval [0, A), $0 < A \le \infty$. This solution will be denoted by $x(\phi)(.)$ so that $x_0(\phi) = \phi$.

Definition. The zero solution, x = 0, of $\dot{x} = F(x_t)$ is stable if for each $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon) > 0$ such that $\|\phi\| < \delta$ implies that $|x(\phi)(t)| < \varepsilon$ for all $t \ge 0$. The zero solution is said to be unstable if it is not stable.

II. MAIN RESULTS

Our main result is the following theorem.

Theorem. In addition to the basic assumptions imposed on the functions g, h and f that appear in Eq. (2), we assume that there exist non-zero constants a_2 , a_4 , a_5 , M_0 and M such that the following conditions hold:

$$\begin{aligned} a_4 > \frac{1}{4}a_2^2, f(0) &= 0, f(x) \neq 0 \quad \text{when} \quad x \neq 0, |f'(x)| \le |a_5|, \\ h(x(t - \tau(t)), 0, ..., u(t - \tau(t))) &= 0, \\ yh(x(t - \tau(t)), ..., u(t - \tau(t))) \ge a_4y^2. \end{aligned}$$

If

$$\sup_{0 \le t < \infty} \tau(t) < \frac{1}{4M_0 M |a_5|},$$

then the zero solution of Eq. (2) is unstable for all arbitrary a_1 .

Proof. We define a Lyapunov functional $V = V(x_t, y_t, z_t, w_t, u_t)$ by

$$V = \frac{1}{2}M[a_{1}z^{2} - 2\int_{0}^{y}g(\eta)\eta d\eta] + [yz - M\{a_{1}yw + yu - zw + \int_{0}^{x}f(s)ds\}] -\lambda \int_{-\tau(t)}^{0}\int_{t+s}^{t}y^{2}(\theta)d\theta ds,$$
(4)

where s is a real variable such that the integral $\int_{-\tau(t)}^{0} \int_{t+s}^{t} y^2(\theta) d\theta ds$ is non-negative, and λ is a positive constant which will be determined later in the proof.

It is clear that

$$V(0,0,\varepsilon^2,\varepsilon,0) = M(\frac{1}{2}a_1\varepsilon^4 + \varepsilon^3) > 0$$

for all sufficiently small $\varepsilon > 0$, which verifies the property (K_1) of Krasovskii [6].

Using the Lyapunov functional V and (3), the time derivative of V yields that

$$\dot{V} = Myh(x(t - \tau(t)), ..., u(t - \tau(t))) + Mw^{2}$$

$$+ (a_{2}M + 1)yw + z^{2} - My \int_{t - \tau(t)}^{t} f'(x(s))y(s)ds$$

$$-\lambda\tau(t)y^{2} + \lambda(1 - \tau'(t)) \int_{t - \tau(t)}^{t} y^{2}(s)ds.$$
(5)

Using the assumptions of the theorem and applying the estimate $2|mn| \le m^2 + n^2$, we get the following estimates for some terms included in (5):

$$yh(x(t - \tau(t)), ..., u(t - \tau(t))) \ge a_4 y^2,$$

$$-My \int_{t-\tau(t)}^t f'(x(s))y(s)ds$$

$$\ge -M|y| \int_{t-\tau(t)}^t |f'(x(s))| |y(s)| ds$$

$$\ge -\frac{1}{2}M|a_5|\tau(t)y^2 - \frac{1}{2}M|a_5| \int_{t-\tau(t)}^t y^2(s)ds.$$

Then, we have

$$\begin{split} \dot{V} &\geq Ma_2y^2 + Mw^2 + (a_2M + 1)yw + z^2 \\ &- \frac{1}{2}M \left| a_5 \right| \tau(t)y^2 - \lambda \tau(t)y^2 + \left\{ \lambda (1 - \tau'(t)) - \frac{1}{2}M \left| a_5 \right| \right\} \\ &\times \int_{t-\tau(t)}^t y^2(s) ds \\ &= M \left[w + \frac{1}{2}M^{-1}(Ma_2 + 1)y \right]^2 \\ &+ \frac{1}{4}M^{-1} \{ (4a_4 - a_2^2)M^2 - 2Ma_2 - 1 \} y^2 + z^2 \\ &- \frac{1}{2}M \left| a_5 \right| \tau(t)y^2 - \lambda \tau(t)y^2 + \{ \lambda (1 - \tau'(t)) - \frac{1}{2}M \left| a_5 \right| \} \\ &\times \int_{t-\tau(t)}^t y^2(s) ds. \end{split}$$
et 1 $- \tau'(t) \geq \frac{1}{2}$ and $\lambda = \frac{1}{2} |a_5| M$. Hence

Let $1 - \tau'(t) \ge \frac{1}{2}$ and $\lambda = \frac{1}{2} |a_5| M$. Hence

$$\dot{V} \geq M \left[w + \frac{1}{2} M^{-1} (Ma_2 + 1)y \right]^2$$
$$\frac{1}{4} M^{-1} \{ (4a_4 - a_2^2) M^2 - 2Ma_2 - 1 \} y^2$$
$$+ z^2 - M |a_5| \tau(t) y^2.$$

It follows that the coefficient $(4a_4 - a_2^2)$ of M^2 is positive. We can also assert that

$$(4a_4 - a_2^2)M^2 - 2Ma_2 - 1 \ge 1$$

when $M > M_0$ for sufficiently large $M_0 \equiv M_0(a_2, a_4) > 0$ so that

$$\dot{V} \geq M_0 \left[w + \frac{1}{2} M_0^{-1} (M_0 a_2 + 1) y \right]^2 + \left\{ \frac{1}{4} M_0^{-1} - M |a_5| \tau(t) \right\} y^2 + z^2.$$

If

$$\sup_{0 \le t < \infty} \tau(t) < \frac{1}{4M_0 M |a_5|},$$

then

$$\dot{V} \ge M_0 \left[w + \frac{1}{2} M_0^{-1} (M_0 a_2 + 1) y \right]^2 + \delta y^2 + z^2 > 0$$
 (6)

for some positive constant δ , which verifies the property (K_2) of Krasovskii [6].

Indeed, because of (6), $\dot{V} = 0$ necessarily implies that

$$y = 0 = z = w. \tag{7}$$

This fact in turn leads to

$$x(t) = \xi$$
 (constant) $u = w' = 0, u' = 0.$ (8)

The substitutions of (7) and (8) in (3) gives that $f(\xi) = 0$ which by the assumptions f(0) = 0, $f(x) \neq 0$ when $x \neq 0$ and $h(x(t - \tau(t)), 0, ..., u(t - \tau(t))) = 0$, implies that $\xi = 0$. Thus $\dot{V} = 0$ implies necessarily that

$$x = 0 = y = z = w = u,$$

which verifies the remaining property (K_3) of Krasovskii [6]. By this discussion, we conclude that the zero solution of Eq.

(2) is unstable. The theorem is established.

REFERENCES

- Èl'sgol'ts, L. È., Introduction to the theory of differential equations with deviating arguments. Translated from the Russian by Robert J. McLaughlin Holden-Day, Inc., San Francisco, Calif.-London-Amsterdam, 1966.
- [2] Ezeilo, J. O. C., Instability theorems for certain fifth-order differential equations. *Math. Proc. Cambridge Philos. Soc.*, no. 2, 84 (1978), 343-350.
- [3] Ezeilo, J. O. C., A further instability theorem for a certain fifth-order differential equation. *Math. Proc. Cambridge Philos. Soc.*, no. 3, 86 (1979), 491-49
- [4] Ezeilo, J. O. C., Extension of certain instability theorems for some fourth and fifth order differential equations. *Atti Accad. Naz. Lincei Rend.Cl. Sci. Fis. Mat. Natur.* (8) 66 (1979), no. 4, 239-242.
- [5] LaSalle, J.; Lefschetz, S., Stability by Liapunov's direct method, with applications. Mathematics in Science and Engineering, Vol. 4 Academic Press, New York-London 1961.
- [6] Krasovskii, N. N., Stability of motion. Applications of Lyapunov's second method to differential systems and equations with delay. Translated by J. L. Brenner Stanford University Press, Stanford, Calif. 1963.
- [7] Li, Wen-jian; Duan, Kui-chen, Instability theorems for some nonlinear differential systems of fifth order. J. Xinjiang Univ. Natur. Sci. 17 (2000), no. 3, 1-5.
- [8] Li, W. J.; Yu, Y. H., Instability theorems for some fourth-order and fifthorder differential equations, (Chinese) J. Xinjiang Univ. Natur. Sci., no. 2, 7(1990), 7-10.
- [9] Sadek, A. I., Instability results for certain systems of fourth and fifth order differential equations. *Appl. Math. Comput.* 145 (2003), no. 2-3, 541-549.
- [10] Sun, W. J.; Hou, X., New results about instability of some fourth and fifth order nonlinear systems. (Chinese) J. Xinjiang Univ. Natur. Sci. 16 (1999), no. 4, 14-17.

- [11] Tiryaki, A., Extension of an instability theorem for a certain fifth order differential equation. National Mathematics Symposium (Trabzon, 1987). J. Karadeniz Tech. Univ. Fac. Arts Sci. Ser. Math.-Phys. 11 (1988), 225-227 (1989).
- [12] Tunç, C., On the instability of solutions of certain nonlinear vector differential equations of fifth order. *Panamer. Math. J.* 14 (2004), no. 4, 25-30.
- [13] Tunç, C., An instability result for a certain non-autonomous vector differential equation of fifth order. *Panamer. Math. J.* 15 (2005), no. 3, 51-58.
- [14] Tunç, C., Further results on the instability of solutions of certain nonlinear vector differential equations of fifth order. *Appl. Math. Inf. Sci.* 2 (2008), no. 1, 51-60.
- [15] Tunç, C., Recent advances on instability of solutions of fourth and fifth order delay differential equations with some open problems. World Scientific Review, Vol. 9, World Scientific Series on Nonlinear Science Series B (Book Series), (2011), 105-116.
- [16] Tunç, C.,On the Instability of Solutions of Nonlinear Delay Differential Equations of Fourth and Fifth Order. *Sains Malaysiana* 40(12), (2011), 1455-1459.
- [17] Tunç, C., An instability theorem for a certain fifth-order delay differential equation, *Filomat*, 25 (2011), no. 3, 145-151.
 [18] Tunç, C., Instability result of a fifth order non-linear delay system.
- [18] Tunç, C., Instability result of a fifth order non-linear delay system. Optoelectronics and Advanced Materials-Rapid Communications. 5 (2011), no. 9, 891-894.
- [19] Tunç, C.; Erdogan, F., On the instability of solutions of certain nonautonomous vector differential equations of fifth order. SUT J. Math. 43 (2007), no. 1, 35-48.
- [20] Tunç, C.; Karta, M., A new instability result to nonlinear vector differential equations of fifth order. *Discrete Dyn. Nat. Soc.* 2008, Art. ID 971534, 6 pp.
- [21] Tunç, C.; Şevli, H., On the instability of solutions of certain fifth order nonlinear differential equations. *Mem. Differential Equations Math. Phys.* 35 (2005), 147-156.

Cemil Tunç Cemil Tunc was born in Yesiloz Koyu (Kalbulas), Horasan-Erzurum, Turkey, in 1958. He received the Ph. D. degree in Applied Mathematics from Erciyes University, Kayseri, in 1993. His research interests include qualitative behaviors of solutions of differential equations. At present he is Professor of Mathematics at Yuzuncu Yil University, Van-Turkey.