

# Rational Chebyshev Tau method for solving natural convection of Darcian fluid about a vertical full cone embedded in porous media with a prescribed wall temperature

Kourosh Parand, Zahra Delafkar, Fatemeh Baharifard

**Abstract**—The problem of natural convection about a cone embedded in a porous medium at local Rayleigh numbers based on the boundary layer approximation and the Darcy's law have been studied before. Similarity solutions for a full cone with the prescribed wall temperature or surface heat flux boundary conditions which is the power function of distance from the vertex of the inverted cone give us a third-order nonlinear differential equation. In this paper, an approximate method for solving higher-order ordinary differential equations is proposed. The approach is based on a rational Chebyshev Tau (RCT) method. The operational matrices of the derivative and product of rational Chebyshev (RC) functions are presented. These matrices together with the Tau method are utilized to reduce the solution of the higher-order ordinary differential equations to the solution of a system of algebraic equations. We also present the comparison of this work with others and show that the present method is applicable.

**Keywords**—Tau method, Semi-infinite, Nonlinear ODE, Rational Chebyshev, Porous media

## I. INTRODUCTION

SPECTRAL methods have been successfully applied in the approximation of differential boundary value problems defined in unbounded domains. For problems whose solutions are sufficiently smooth, they exhibit exponential rates of convergence/spectral accuracy. We can apply different spectral methods that are used to solve problems in semi-infinite domains. The most common method is the use of polynomials that are orthogonal over unbounded domains, such as the Hermite and Laguerre spectral methods [1], [2], [3], [4], [5], [6], [7], [8]. Guo [9], [10], [11] proposed a method that proceeds by mapping the original problem in an unbounded domain to a problem in a bounded domain, and then using suitable Jacobi polynomials to approximate the resulting problems. Another approach is replacing infinite domain with  $[-L, L]$  and semi-infinite interval with  $[0, L]$  by choosing  $L$ , sufficiently large. This method is named domain truncation [12]. Another effective direct approach for solving such problems is based on rational approximations. Christov [13] and Boyd [14], [15] developed some spectral

methods on unbounded intervals by using mutually orthogonal systems of rational functions. Boyd [15] defined a new spectral basis, named rational Chebyshev functions on the semi-infinite interval, by mapping to the Chebyshev polynomials. Guo et al. [16] introduced a new set of rational Legendre functions which are mutually orthogonal in  $L^2(0, +\infty)$ . They applied a spectral scheme using the rational Legendre functions for solving the Korteweg-de Vries equation on the half line. Boyd et al. [17] applied pseudospectral methods on a semi-infinite interval and compared rational Chebyshev, Laguerre and mapped Fourier sine. Parand et al. [18], [19], [20], [21], [22], [23], [24] and Tajvidi et al. [25] applied spectral method to solve nonlinear ordinary differential equations on semi-infinite intervals. Their approach was based on rational Tau and collocation method.

A porous medium means a material consisting of a solid matrix with an interconnected void. The solid matrix is either rigid or it undergoes small deformation. The interconnectedness of the pores allows the flow of one or more fluids through the material [26].

Natural convective heat transfer in porous media has received considerable attention during the past few decades. This interest can be attributed due to its wide range of applications in ceramic processing, nuclear reactor cooling system, crude oil drilling, chemical reactor design, ground water pollution and filtration processes. External natural convection in a porous medium adjacent to heated bodies was analyzed by Nield and Bejan [26], Merkin [27], [28], Minkowycz and Cheng [29], [30], [31], Pop and Cheng [32], [33], Ingham and Pop [34] and vafai [35]. In all of these analysis, it is assumed that boundary layer approximations are applicable and the coupled set of governing equations are solved by numerical methods.

In this paper, we introduce a new computational method to solve the problem of natural convection about an inverted heated cone embedded in a porous medium of infinite extent. No similarity solution exists for the truncated cone, but for the case of full cone similarity solutions exist if the prescribed wall temperature or surface heat flux is a power function of distance from the vertex of the inverted cone [26], [32], [36].

Bejan and Khair [37] used Darcy's law to study the vertical natural convective flows driven by temperature and concentration gradients. Nakayama and Hossain [38] applied the integral method to obtain the heat and mass transfer by free convection from a vertical surface with constant wall temperature and

K. Parand is with Department of Computer Sciences, Shahid Beheshti University, Tehran, Iran (Phone:+98 21 22431653; Fax:+98 21 22431650; e-mail: k.parand@sbu.ac.ir).

Z. Delafkar is with Department of Computer Sciences, Shahid Beheshti University, Tehran, Iran (e-mail: zahra.delafkar@gmail.com).

F. Baharifard is with Department of Computer Sciences, Shahid Beheshti University, Tehran, Iran (e-mail: fateme.baharifard@gmail.com).

concentration. Yih [39] examined the coupled heat and mass transfer by free convection over a truncated cone in porous media for variable wall temperature and variable heat and mass fluxes, Also he [40] applied the uniform transpiration effect on coupled heat and mass transfer in mixed convection about inclined surfaces in porous media for the entire regime. Cheng [41] used an integral approach to study the heat and mass transfer by natural convection from truncated cones in porous media with variable wall temperature and [42] studied the Soret and Dufour effects on the boundary layer flow due to natural convection heat and mass transfer over a vertical cone in a porous medium saturated with Newtonian fluids with constant wall temperature. Natural convective mass transfer from upward-pointing vertical cones, embedded in saturated porous media, has been studied using the limiting diffusion [43]. The natural convection along an isothermal wavy cone embedded in a fluid-saturated porous medium are presented in [44], [45]. The problem of steady laminar hydromagnetic heat transfer over a vertical plate embedded in a uniform porous medium is studied in [46], [47], [48]. In [36] fluid flow and heat transfer of vertical full cone embedded in porous media have been solved by Homotopy analysis method.

Also in this work, we have applied rational Chebyshev Tau approach to find the numerical solutions of nonlinear ordinary differential equations arising from similarity solution of natural convection of Darcian fluid about a vertical full cone embedded in porous media. The operational matrices of the derivative and the product of rational Chebyshev functions are derived. These matrices together with the Tau method are then utilized to evaluate the solution to this equation. The Tau method was invented by Lanczos [49]. The method is based on expanding the required approximate solution as the elements of a complete set of orthogonal functions. In the Tau method, unlike the Galerkin approximation, the expansion functions are not required to satisfy the boundary constraint individually [50], [51]. we have made a comparison with [32], [36] and other solutions based on Runge-Kutta method. The similarity solution which was applied by [32] is based on thermal boundary conditions as wall temperature power function of distance from the vertex of the inverted cone. The transformed ordinary differential equation, with the corresponding boundary conditions, was solved by Runge-Kutta method for the systematically guessing of the missing initial conditions [36]. In Spectral methods an important question is: What sets of "basis functions" will work? It is obvious that we would like our basis sets to have a number of properties: (i) easy computation (ii) rapid convergence and (iii) completeness, which means that any solution can be represented to arbitrarily high accuracy by taking the truncation  $N$  to be sufficiently large[12]. In this area we find that the orthogonal rational Chebyshev functions have these properties. As a sample, in this paper we show that rational Chebyshev functions with these properties can be applied to approximate third order nonlinear differential equations. Numerical results indicate the convergence and effectiveness of the present approach

## II. PROBLEM FORMULATION

Consider an inverted cone with semi-angle  $\gamma$  and take axes in the manner indicated in Fig. 1(a). The stream function  $\psi$  defined as

$$u = \frac{1}{r} \frac{\partial \psi}{\partial y}, \quad v = -\frac{1}{r} \frac{\partial \psi}{\partial x}. \quad (1)$$

In terms of the heated frustum  $x = x_0$  is considered [26], [36].

The boundary layer equations are:

$$\frac{1}{r} \frac{\partial^2 \psi}{\partial y^2} = \frac{g\beta K}{v} \frac{\partial T}{\partial y}, \quad (2)$$

$$\frac{1}{r} \left( \frac{\partial \psi}{\partial y} \frac{\partial T}{\partial x} - \frac{\partial \psi}{\partial x} \frac{\partial T}{\partial y} \right) = \alpha \frac{\partial^2 T}{\partial y^2}.$$

We have approximately  $r = x \sin(\gamma)$  and suppose the temperature is power function of distance from the vertex of the inverted cone. Accordingly, the boundary conditions are:

$$\begin{aligned} u = 0, \quad T = T_\infty & \quad y \rightarrow \infty, & (3) \\ u = 0 & \quad y = 0, x_0 \leq x < \infty, \\ T = T_w = T_\infty + A(x - x_0)^\lambda & \quad y = 0, x_0 \leq x < \infty. \end{aligned}$$

For the case of a full cone ( $x_0 = 0$ , Fig.1(b)) a similarity solution exists. In the case of prescribed wall temperature, we let:

$$\psi = \alpha r Ra_x^{1/2} f(\eta), \quad (4)$$

$$T - T_\infty = (T_w - T_\infty) \theta(\eta),$$

$$\eta = \frac{y}{x} Ra_x^{1/2},$$

where the Rayleigh number is

$$Ra_x = \frac{g\beta K \cos(\gamma)(T_w - T_\infty)x}{\nu \alpha}. \quad (5)$$

The governing equations are, Ref. [26], [32], [36]:

$$f' = \theta, \quad (6)$$

$$\theta'' + \left( \frac{\lambda + 3}{2} \right) f \theta' - \lambda f' \theta = 0,$$

with boundary conditions,

$$f(0) = 0, \quad \theta(0) = 1, \quad \theta(\infty) = 0 \quad (7)$$

Finally, we have:

$$\begin{cases} ODE. & f''' + \left( \frac{\lambda + 3}{2} \right) f f'' - \lambda (f')^2 = 0, \\ B.C. & f(0) = 0, \quad f'(0) = 1, \quad f'(\infty) = 0. \end{cases} \quad (8)$$

It is of interest to obtain the value of the local Nusselt number which is defined as:

$$Nu_x = \frac{q_w x}{k(T_w - T_\infty)}. \quad (9)$$

Where  $q_w$  for the case of prescribed wall temperature can be computed from:

$$q_w = -k \left( \frac{\partial T}{\partial y} \right)_{y=0}. \quad (10)$$

From Eqs. (4), (5), (9) and (10), it follows that the local Nusselt number is given by [26], [36]:

$$Nu_x = Ra_x^{1/2} [-\theta'(0)]. \quad (11)$$

**Nomenclature**

$A$	prescribed constant
$f$	similarity function for stream function
$g$	acceleration due to gravity
$K$	permeability of the fluid-saturated porous medium
$Nu$	local Nusselt number
$q_w$	surface heat flux
$r$	local radius of the cone
$Ra_x$	local Raleigh number
$T$	temperature
$T_\infty$	ambient temperature
$T_w$	wall temperature
$u, v$	velocity vector along x, y axis
$x, y$	Cartesian coordinate system
$x_0$	distance of start point of cone from the vertex

**Greek symbols**

$\theta$	similarity function for temperature
$\eta$	independent dimensionless parameter
$\lambda$	prescribed constants
$\beta$	expansion coefficient of the fluid
$\alpha$	thermal diffusivity the fluid-saturated porous medium
$\nu$	kinematics viscosity of the fluid
$\psi$	stream function




Fig. 1. (a) Coordinate system for the boundary layer on a heated frustum of a cone, (b) full cone,  $x_0 = 0$

**III. PROPERTIES OF RATIONAL CHEBYSHEV FUNCTIONS**

**A. Rational Chebyshev functions**

The well-known Chebyshev polynomials are orthogonal in the interval  $[-1, 1]$  with respect to the weight function  $w(y) = \frac{1}{\sqrt{1-y^2}}$  and can be determined with the aid of the following recurrence formula:

$$T_0(y) = 1, \quad T_1(y) = y, \\ T_{n+1}(y) = 2yT_n(y) - T_{n-1}(y), \quad n \geq 1.$$

The RC functions are defined in [12] as

$$R_n(x) = T_n\left(\frac{x-L}{x+L}\right).$$

Thus RC functions satisfy:

$$R_0(x) = 1, \quad R_1(x) = \frac{x-L}{x+L}, \quad (12) \\ R_{n+1}(x) = 2\left(\frac{x-L}{x+L}\right)R_n(x) - R_{n-1}(x), \quad n \geq 1.$$

Rational Chebyshev functions are orthogonal with respect

to the weight function  $w(x) = 1/(\sqrt{x}(x+L))$  in the interval  $[0, +\infty)$ , with the orthogonality property:

$$R_n(x) = \int_0^\infty R_n(x)R_m(x)w(x)dx = \frac{c_m\pi}{2} \delta_{nm}, \quad (13)$$

$$c_m = \begin{cases} 2, & m = 0, \\ 1, & m \neq 1, \end{cases} \quad (14)$$

and  $\delta_{nm}$  is the Kronecker function.

**B. Function approximation**

A function  $f(x)$  defined over the interval  $[0, +\infty)$  may be expanded as

$$f(x) = \sum_{i=0}^\infty a_i R_i(x), \quad (15)$$

where

$$a_i = \frac{2}{c_i\pi} \int_0^\infty R_i(x)f(x)w(x)dx. \quad (16)$$

If  $f(x)$  in Eq. (15) is truncated up to the  $N$ th terms, then it can be written as

$$f_N(x) = \sum_{i=0}^{N-1} a_i R_i(x) = A^T R(x), \quad (17)$$

with

$$A = [a_0, a_1, \dots, a_{N-1}]^T, \quad (18)$$

$$R(x) = [R_0(x), R_1(x), \dots, R_{N-1}(x)]^T. \quad (19)$$

**C. Operational Matrix of Derivative**

The derivative of the vector  $R(x)$  defined in Eq. (19) can be expressed as

$$R'(x) = \frac{dR(x)}{dx} \simeq DR(x), \quad (20)$$

where  $D$  is the  $N \times N$  operational matrix for the derivative. Differentiating Eq. (12) we get:

$$R'_{n+1}(x) = \frac{4L}{(x+L)^2} R_n(x) + 2\left(\frac{x-L}{x+L}\right)R'_n(x) - R'_{n-1}(x), \quad n \geq 1.$$

By using

$$\frac{2L}{(x+L)^2} = \frac{1}{L} \left[ \frac{3}{4}R_0(x) - R_1(x) + \frac{1}{4}R_2(x) \right] = R'_1(x), \quad (21)$$

the elements  $d_{ij}$  of the matrix  $D$  can be obtained from

$$\begin{cases} R'_{n+1}(x) = 2(R_1(x)R_n(x))' - R'_{n-1}(x), & n \geq 1, \\ R'_1(x) = \frac{1}{L} \left[ \frac{3}{4}R_0(x) - R_1(x) + \frac{1}{4}R_2(x) \right], \\ R'_0(x) = 0. \end{cases} \quad (22)$$

The general form of the matrix  $D$  is a lower-Heisenberg matrix. The matrix  $D$  can be expressed as  $D = \frac{1}{L}(D_1 + D_2)$ , where  $D_1$  is a tridiagonal matrix which is obtained from

$$D_1 = \text{diag}\left(\frac{7}{4}i, -i, \frac{i}{4}\right), \quad i = 0, \dots, N-1, \quad (23)$$

and the  $d_{ij}$  elements of matrix  $D_2$  are obtained from  $d_{10} = -1$  and

$$d_{ij} = \begin{cases} 2, & j \geq i - 1, \\ kic_j, & j < i - 1, \end{cases} \quad (24)$$

where  $k = (-1)^{i+j+1}$ ,  $c_0 = 1$  and  $c_j = 2$ , for  $j > 1$ . for  $N = 7$  we have

$$D = \frac{1}{L} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 3/4 & -1 & 1/4 & 0 & 0 & 0 & 0 \\ -2 & 7/2 & -2 & 1/2 & 0 & 0 & 0 \\ 3 & -6 & 21/4 & -3 & 3/4 & 0 & 0 \\ -4 & 8 & -8 & 7 & -4 & 1 & 0 \\ 5 & -10 & 10 & -10 & 35/4 & -5 & 5/4 \\ -6 & 12 & -12 & 12 & -12 & 21/2 & -6 \end{pmatrix}$$

D. The product operational matrix

The following property of the product of two rational Chebyshev function vectors will also be used. Since

$$R_m(x)R_n(x) = \frac{1}{2}[R_{m+n}(x) + R_{|n-m|}(x)], \quad (25)$$

if we truncate the terms higher than  $R_{N-1}$ , then we have

$$R(x)R^T(x)A \simeq \tilde{A}R(x), \quad (26)$$

where  $\tilde{A}$  is an  $N \times N$  product operational matrix for the vector  $A$  and the elements  $\tilde{a}_{ij}$  of the matrix  $\tilde{A}$  can be obtained from

$$\tilde{a}_{ij} = \begin{cases} a_j, & i = 0, \\ \frac{1}{2}a_i, & j = 0, i \neq 0, \\ a_{i-j} + \frac{1}{2}a_{i+j}, & i = j, i, j \neq 0, \\ \frac{1}{2}(a_{|i-j|} + a_{i+j}), & i \neq j, i, j \neq 0, i + j \leq N, \\ \frac{1}{2}(a_{|i-j|}), & i \neq j, i, j \neq 0, i + j > N, \end{cases}$$

and the matrix  $\tilde{A}$  is in the form

$$\tilde{A} = \begin{pmatrix} a_0 & a_1 & a_2 & a_3 & \dots & a_{N-1} \\ \frac{1}{2}a_1 & a_0 + \frac{1}{2}a_2 & \frac{1}{2}(a_1 + a_3) & \frac{1}{2}(a_2 + a_4) & \dots & \frac{1}{2}a_{N-2} \\ \frac{1}{2}a_2 & \frac{1}{2}(a_1 + a_3) & a_0 + \frac{1}{2}a_4 & \frac{1}{2}(a_1 + a_5) & \dots & \frac{1}{2}a_{N-3} \\ \frac{1}{2}a_3 & \frac{1}{2}(a_2 + a_4) & \frac{1}{2}(a_1 + a_5) & a_0 + \frac{1}{2}a_6 & \dots & \frac{1}{2}a_{N-4} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{1}{2}a_{N-1} & \frac{1}{2}a_{N-2} & \frac{1}{2}a_{N-3} & \frac{1}{2}a_{N-4} & \dots & a_0 \end{pmatrix}$$

IV. SOLVING THE PROBLEM

To solve Eq. (8) with initial conditions, we set

$$f(x) \simeq f_N(x), \quad (27)$$

$$f_N(x) = \sum_{i=0}^{N-1} a_i R_i(x) = A^T R(x), \quad (28)$$

$$f_N^{(j)}(x) = \sum_{i=0}^{N-1} a_i R_i^{(j)}(x) \simeq A^T D^j R(x) \quad j = 1, 2, 3. \quad (29)$$

where  $R_i(x)$  defined in Eq. (12) in a way that  $L = 1$  and  $D^j$  is the  $j$ th power of the matrix  $D$  given in Eq. (20).

TABLE I  
A COMPARISON OF METHODS IN [32], [36] AND THE PRESENT METHOD WITH THE VALUES FOR  $f''(0)$

$\lambda$	Runge-Kutta[36]	RCT method(N=26 and L=1)		Other methods	
		RCT	HAM[36]	Ref. [32]	
0	-0.76854	-0.76600	-0.77363	-0.769	
1/4	-0.88498	-0.88330	-0.88890	-	
1/3	-0.92101	-0.92100	-0.92433	-0.921	
1/2	-0.98956	-0.98581	-0.99382	-0.992	
3/4	-1.08518	-1.08598	-1.08840	-	
1	-1.17372	-1.17040	-1.17686	-	

Using Eqs. (28) and (29) we get

$$f_N''(x) \simeq A^T DR(x)R^T(x)D^T A = \quad (30)$$

$$A^T DR(x)R^T(x)E \simeq A^T D\tilde{E}R(x),$$

$$f_N(x)f_N''(x) \simeq A^T R(x)R^T(x)(D^2)^T A = \quad (31)$$

$$A^T R(x)R^T(x)F \simeq A^T \tilde{F}R(x),$$

Where  $E = D^T A$  and  $F = (D^2)^T A$  and the matrices  $\tilde{A}$ ,  $\tilde{E}$  and  $\tilde{F}$  can be calculated similarly to Eq. (26). Using Eqs. (28-31) the residual function ( $Res(x)$ ) for Eq. (8) can be written as

$$Res(x) = [A^T D^3 + (\frac{\lambda + 3}{2})A^T \tilde{F} - \lambda A^T D\tilde{E}]R(x). \quad (32)$$

As in typical Tau method [51], [19], [25] we generate  $N - 2$  algebraic equations by applying

$$\langle Res(x), R_k(x) \rangle = \int_0^{+\infty} Res(x)R_k(x)w(x)dx = 0, \quad k = 0, 1, \dots, N - 3. \quad (33)$$

using Eqs. (28-29), for the initial conditions we get

$$f_N(0) = A^T R(0) = 0, \quad f_N'(0) = A^T DR(0) = 1. \quad (34)$$

Take into account

$$\lim_{x \rightarrow \infty} R'(x) = 0,$$

and Eq. (29) the boundary condition  $f'(\infty) = 0$  already is satisfied.

Eq. (33-34) generates a set of  $N$  nonlinear algebraic equations. Consequently, the unknown coefficients  $a_i$  of the vector  $A$  in Eq. (28) can be calculated.

Table I shows good agreement between RCT method and Homotopy analysis and Runge-Kutta method [36] for  $f''(0)$  or  $\theta'(0)$  with various  $\lambda$  and  $N = 26$ .

The results for  $f'(\eta)$  have been shown in Table II with two selected  $\lambda = 0$  and  $\lambda = 1/2$  and comparison have been made between the Runge-Kutta's solution [36] and the presented numerical solution. Absolute errors show that RCT give us approximate solution with a high degree of accuracy with a small  $N$ .

The resulting graph of Eq. (8) for  $N = 26$  and  $L = 1$  is shown in Fig. 2.

V. CONCLUSION

In this study, we have applied rational Chebyshev Tau method to solve three order nonlinear differential equations




Fig. 2. RCT approximation of  $f'(\eta)$  for different values  $\lambda = 0, 1/4, 1/3, 1/2, 3/4$  and  $1$

TABLE II  
COMPARISON BETWEEN RCT SOLUTION AND RUNGE-KUTTA SOLUTION FOR  $f'(0)$  WITH  $\lambda = 0$  AND  $\lambda = 1/2$  WITH  $N = 26$  AND  $L = 1$

$\eta$	$\lambda = 0, f'(\eta)$			$\lambda = 1/2, f'(\eta)$		
	RCT Solution	Runge-Kutta Solution	Absolute Error	RCT Solution	Runge-Kutta Solution	Absolute Error
0	1.0000	1.0000	0.0000	0.9999	1.0000	0.0000
0.2	0.8477	0.8478	0.0000	0.8130	0.8130	0.0000
0.4	0.7036	0.7036	0.0000	0.6500	0.6500	0.0000
0.6	0.5732	0.5733	0.0001	0.5125	0.5124	0.0001
0.8	0.4598	0.4599	0.0001	0.3994	0.3994	0.0001
1	0.3641	0.3643	0.0001	0.3084	0.3084	0.0001
1.2	0.2853	0.2855	0.0001	0.2364	0.2364	0.0000
1.4	0.2218	0.2218	0.0000	0.1802	0.1802	0.0000
1.6	0.1713	0.1713	0.0001	0.1367	0.1367	0.0002
1.8	0.1316	0.1315	0.0003	0.1034	0.1034	0.0003
2	0.1007	0.1007	0.0003	0.0779	0.0780	0.0004

arising from similarity solution of natural convection of Darcian fluid about a vertical full cone embedded in porous media in whole domain, and compared the results with [36] and [32]. Local similarity solutions are obtained for a full cone with the prescribed wall temperature being a power function of distance from the vertex of the inverted cone. The obtained approximate solution by RCT provides us with calculating Nusselt number. The operational matrices of the derivative and product of RC functions together with the Tau method have been obtained to solve the higher-order ordinary differential equations numerically. The stability and convergence of the Tau approximations [50] are merits that make the approach very useful. Through the comparisons among the numerical solutions of Cheng et al. [32], Sohoulil et al. [36] and the current work, it has been shown that the present work has provided acceptable approach for this type equations.

VI. ACKNOWLEDGMENTS

The research of corresponding author (K. Parand) was supported by a grant from Shahid Beheshti University.

REFERENCES

- [1] O. Coulaud, D. Funaro, O. Kavian, Laguerre spectral approximation of elliptic problems in exterior domains, *Comput. Method. Appl. Mech. Eng.* 80 (1990) 451–458.
- [2] D. Funaro, O. Kavian, Approximation of some diffusion evolution equations in unbounded domains by hermite functions, *Math. Comput.* 57 (1991) 597–619.
- [3] D. Funaro, Computational aspects of pseudospectral laguerre approximations, *Appl. Numer. Math.* 6 (1990) 447–457.
- [4] B. Guo, Error estimation of hermite spectral method for nonlinear partial differential equations, *Math. Comput.* 68 (227) (1999) 1067–1078.
- [5] B. Guo, J. Shen, Laguerre-galerkin method for nonlinear partial differential equations on a semi-infinite interval, *Numer. Math.* 86 (2000) 635–654.
- [6] Y. Maday, B. Pernaud-Thomas, H. Vandeven, Reappraisal of laguerre type spectral methods, *La. Rech. Aerospaciale.* 6 (1985) 13–35.
- [7] J. Shen, Stable and efficient spectral methods in unbounded domains using laguerre functions, *SIAM J. Numer. Anal.* 38 (4) (2000) 1113–1133.
- [8] H. Siyyam, Laguerre tau methods for solving higher order ordinary differential equations, *J. Comput. Anal. Appl.* 3 (2) (2001) 173–182.
- [9] B. Guo, Gegenbauer approximation and its applications to differential equations on the whole line, *J. Comput. Anal. Appl.* 226 (1) (1998) 180–206.
- [10] B. Guo, Jacobi spectral approximation and its applications to differential equations on the half line, *J. Comput. Math.* 18 (1) (2000) 95–112.
- [11] B. Guo, Jacobi approximations in certain hilbert spaces and their applications to singular differential equations, *J. Comput. Anal. Appl.* 243 (2) (2000) 373–408.
- [12] J. Boyd, *Chebyshev and Fourier Spectral Methods*, 2nd Edition, Dover, New York, 2000.
- [13] C. Christov, A complete orthogonal system of functions in  $l^2(-\infty, \infty)$  space, *SIAM J. Appl. Math.* 42 (1982) 1337–1344.
- [14] J. Boyd, Spectral methods using rational basis functions on an infinite interval, *J. Comput. Phys.* 69 (1) (1987) 112–142.
- [15] J. Boyd, Orthogonal rational functions on a semi-infinite interval, *J. Comput. Phys.* 70 (1) (1987) 63–88.
- [16] B. Guo, J. Shen, Z. Wang, A rational approximation and its applications to differential equations on the half line, *J. Sci. Comput.* 15 (2) (2000) 117–147.
- [17] J. Boyd, C. Rangan, P. Bucksbaum, Pseudospectral methods on a semi-infinite interval with application to the hydrogen atom: a comparison of the mapped fourier-sine method with laguerre series and rational chebyshev expansions, *J. Comput. Phys.* 188 (1) (2003) 56–74.
- [18] K. Parand, M. Razzaghi, Rational chebyshev tau method for solving voltaerra’s population model, *Appl. Math. Comput.* 149 (2004) 893–900.
- [19] K. Parand, M. Razzaghi, Rational chebyshev tau method for solving higher-order ordinary differential equations, *Int. J. Comput. Math.* 81 (2004) 73–80.
- [20] K. Parand, M. Razzaghi, Rational legendre approximation for solving some physical problems on semi-infinite intervals, *Phys. Scr.* 69 (2004) 353–357.
- [21] K. Parand, M. Shahini, Rational chebyshev pseudospectral approach for solving thomas-fermi equation, *Phys. Lett. A.* 373 (2009) 210–213.
- [22] K. Parand, A. Taghavi, Rational scaled generalized laguerre function collocation method for solving the blasius equation, *J. Comput. Appl. Math.* 233 (4) (2009) 980–989.
- [23] K. Parand, M. Shahini, M. Dehghan, Rational legendre pseudospectral approach for solving nonlinear differential equations of lane-emden type, *J. Comput. Phys.* 228 (23) (2009) 8830–8840.
- [24] K. Parand, M. Dehghan, A. Pirkhedri, Sinc-collocation method for solving the blasius equation, *Phys. Lett. A.* 273 (44) (2009) 4060–4065.
- [25] T. Tajvidi, M. Razzaghi, M. Dehghan, Modified rational legendre approach to laminar viscous flow over a semi-infinite flat plate, *Chaos Soliton. Fract.* 35 (2008) 59–66.
- [26] D. Nield, A. Bejan, *Convection in Porous Media*, Springer-Verlag, New York, 2006.
- [27] J. Merkin, Free convection boundary layers in a saturated porous medium with lateral mass flux, *Int. J. Heat. Mass. Tran.* 21 (1978) 1499–1504.
- [28] J. Merkin, Free convection boundary layers on axi-symmetric and two dimensional bodies of arbitrary shape in a saturated porous medium, *Int. J. Heat. Mass. Tran.* 22 (1979) 1461–1462.
- [29] W. Minkowycz, P. Cheng, Free convection about a vertical cylinder embedded in a porous medium, *Int. J. Heat. Mass. Tran.* 19 (1976) 805–813.

- [30] W. Minkowycz, P. Cheng, Local non-similar solutions for free convective flow with uniform lateral mass flux in a porous medium, *Int. J. Heat. Mass. Tran.* 9 (1982) 159–168.
- [31] W. Minkowycz, P. Cheng, F. Moalem, The effect of surface mass transfer on buoyancy induced darcian flow adjacent to a horizontal heated surface, *Int. Commun. Heat. Mass.* 12 (1985) 55–65.
- [32] P. Cheng, T. Le, I. Pop, Natural convection of a darcian fluid about a cone, *Int. Commun. Heat. Mass.* 12 (1985) 705–717.
- [33] I. Pop, P. Cheng, An integral solution for free convection of a darcian fluid about a cone with curvature effects, *Int. Commun. Heat. Mass.* 13 (1986) 433–438.
- [34] D. Ingham, I. Pop, Natural convection about a heated horizontal cylinder in a porous medium, *J. Fluid. Mech.* 184 (1987) 157–181.
- [35] K. Vafai, *Handbook of porous media*, Marcel Dekker, New York, 2000.
- [36] A. Sohoulil, D. Domairry, M. Famouri, A. Mohsenzadeh, Analytical solution of natural convection of darcian fluid about a vertical full cone embedded in porous media prescribed wall temperature by means of ham, *Int. Commun. Heat. Mass.* 35 (2008) 1380–1384.
- [37] A. Bejan, K. Khair, Heat and mass transfer by natural convection in a porous medium, *Int. J. Heat. Mass. Tran.* 28 (1985) 909–918.
- [38] A. Nakayama, M. Hossain, An integral treatment for combined heat and mass transfer by natural convection in a porous medium, *Int. J. Heat. Mass. Tran.* 38 (1995) 761–765.
- [39] K. Yih, Coupled heat and mass transfer by free convection over a truncated cone in porous media:  $Vwt/vwc$  or  $vhf/vmf$ , *Acta. Mech.* 137 (1999) 83–97.
- [40] K. Yih, Uniform transpiration effect on coupled heat and mass transfer in mixed convection about inclined surfaces in porous media: the entire regime, *Acta. Mech.* 132 (1999) 229–240.
- [41] C. Cheng, An integral approach for heat and mass transfer by natural convection from truncated cones in porous media with variable wall temperature and concentration, *Int. Commun. Heat. Mass.* 27 (2000) 437–548.
- [42] C. Cheng, Soret and dufour effects on natural convection heat and mass transfer from a vertical cone in a porous medium, *Int. Commun. Heat. Mass.* 36 (2009) 1020–1024.
- [43] S. Rahman, K. Mahgoub, A. Nafees, Natural convective mass transfer from upward pointing conical surfaces in porous media, *Chem. Eng. Commun.* 194 (2007) 280–290.
- [44] I. Pop, T. Na, Natural convection of a darcian fluid about a wavy cone, *Int. Commun. Heat. Mass.* 21 (1994) 891–899.
- [45] I. Pop, T. Na, Natural convection over a frustum of a wavy cone in a porous medium, *Mech. Res. Commun.* 22 (1995) 181–190.
- [46] O. D. Makinde, On mhd boundary-layer flow and mass transfer past a vertical plate in a porous medium with constant heat flux, *Int. J. Numer. Meth. Heat. Fluid. Flow.* 19 (2009) 546–554.
- [47] O. D. M. A. Ogulu, Unsteady hydromagnetic free convection flow of a dissipative and radiating fluid past a vertical plate with constant heat flux, *Chem. Eng. Comm.* 196 (2009) 454–462.
- [48] P. S. O. D. Makinde, Magnetohydrodynamic mixed-convective flow and heat and mass transfer past a vertical plate in a porous medium with constant wall suction, *J. Heat. Transfer.* 130 (2008) 1–8.
- [49] C. Lanczos, *Applied Analysis*, Springer-Verlag, 1987.
- [50] D. Gottlieb, M. Hussaini, S. Orszg, In: R. Voigt and D. Gottlieb and M. Y. Hussaini (Eds.) *Theory and applications of spectral methods in spectral methods for partial differential equations*, Philadelphia: SIAM, 1984.
- [51] C. Canuto, M. Hussaini, A. Quarteroni, T. Zang, *Spectral methods in fluid dynamic*, Englewood Cliffs (NJ): Prentice-Hall, 1986.