

# Reflectance Imaging Spectroscopy Data (Hyperspectral) for Mineral Mapping in the Orientale Basin Region on the Moon Surface

V. Sivakumar, R. Neelakantan

**Abstract**—Mineral mapping on the Moon surface provides the clue to understand the origin, evolution, stratigraphy and geological history of the Moon. Recently, reflectance imaging spectroscopy plays a significant role in identifying minerals on the planetary surface in the Visible to NIR region of the electromagnetic spectrum. The Moon Mineralogy Mapper ( $M^3$ ) onboard Chandrayaan-1 provides unprecedented spectral data of lunar surface to study about the Moon surface. Here we used the  $M^3$  sensor data (hyperspectral imaging spectroscopy) for analysing mineralogy of Orientale basin region on the Moon surface. Reflectance spectrums were sampled from different locations of the basin and continuum was removed using ENvironment for Visualizing Images (ENVI) software. Reflectance spectra of unknown mineral composition were compared with known Reflectance Experiment Laboratory (RELAB) spectra for discriminating mineralogy. Minerals like olivine, Low-Ca Pyroxene (LCP), High-Ca Pyroxene (HCP) and plagioclase were identified. In addition to these minerals, an unusual type of spectral signature was identified, which indicates the probable Fe-Mg-spinel lithology in the basin region.

**Keywords**—Chandrayaan-1, moon mineralogy mapper, orientale basin, moon, spectroscopy, hyperspectral.

## I. INTRODUCTION

**H**YPERSPECTRAL imaging spectrometer (spectroscopy) acquire images in many narrow and contiguous spectral bands. Imaging spectrometry has been widely used in geological / mineral mapping in the Earth and planetary surface. Hyperspectral data facilitate the discriminating various minerals across the spectrum. Reflectance spectroscopic data is helpful for identifying minerals on the planetary surface due to analytical absorption bands as a result of transitions of electrons in a crystal field [1]. The spectral (spectra) characteristics such as shape and absorption centers are very much useful for identifying various minerals. Mapping and analysing mineralogy of the lunar surface provides insights into the origin, evolution of crust, geological history and stratigraphy of the Moon. The  $M^3$  instrument onboard Chandrayaan-1 provides reflectance spectral data of Moon surface. Many literatures are provides insight into detection of minerals on the lunar surface based on minerals absorption characteristics [2]-[7]. This present study aims to

identifying minerals on the lunar surface using Chandrayaan-1  $M^3$  data.

## II. STUDY AREA

The Orientale basin is located on the western limb of the Moon highland terrain and centered at 19° S, 95° W (Fig. 1). This basin is the youngest and most well-preserved large multi ring basin on the Moon and is sparsely filled by mare basalt [8]. The Orientale is a large impact structure basin and displays at least four concentric rings (multi-ring basin) [8], [9]. Scientists are curiosity to investigate on this basin due to its unique morphological setup. The study area elevation ranges from -4700 m to 9400 m (Fig. 2). Lower areas are shaded blue, with higher altitudes in red. The lowest areas are about -4700 m below the average height with the highest being about 9400 m above average (Fig. 2).

### A. Geology

Orientale basin partly filled by mare basalt, therefore its original floor configuration can be clearly seen over most of the basin interior (Fig. 3). Recently, this basin has been remapped and named two new units namely mare and massif material by [10], utilizing the stratigraphic nomenclature of [11], which has been largely unchanged except that some formations have been subdivided into members on the basis of surface texture. Several geological formations collectively make up the Orientale Group; in the interior of the basin, formations are the Maunder, Montes Rook, and Hevelius formations. The Orientale basin interior displays a number of small melt ponds. The innermost center of the basin covers thinly basalt [10]. The basalts of Mare Orientale appear to be moderate in titanium content (~2.3 wt), which is relatively low Ti compared to the Apollo samples but higher than the Ti content of other typical farside maria [12]. The other informal rock unit is made up of the massifs of the inner basin rings. Some parts of the Inner ring are composed of massifs made up of pure anorthosite [13]-[15]. In some cases, these anorthosites are shocked to levels of at least 20 GPa, but less than 30 GPa, as evidenced by the presence of the 1250 nm plagioclase absorption feature [15].

## III. MATERIALS AND METHODS

$M^3$  data is used for collecting reflectance spectra for identifying the minerals in the Orientale basin area. The Indian Space Research Organisation's (ISRO) first mission to the Moon Chandrayaan-1 has  $M^3$  instrument, as a guest instrument from NASA [16]. The  $M^3$  records the reflected

Sivakumar V. is with the Department of Geology, Periyar University, Salem - 636011, India; Centre for Development of Advanced Computing (C-DAC), Pune University Campus, Pune - 411007, India (e-mail: v.sivakumars@gmail.com).

Neelakantan R. is with the Department of Industries and Earth Sciences, Tamil University, Tanjore - 613010, India (e-mail: neels2004@gmail.com).

radiance from the Moon's surface in pushbroom mode between 0.46 and 2.97  $\mu\text{m}$  in 85 contiguous spectral bands,

between 20 – 40 nm spectral sampling with 140 m/pixel spatial resolution [17].




Fig. 1 Study area location ( $M^3$  coverage) is highlighted with black color box on Diviner Global Composition map [18]




Fig. 2 LRO's Wide-Angle Camera Digital Terrain Model of the Orientale Basin (1,100 km diameter). Lower areas are shaded blue, with higher altitudes in red. The lowest areas are about -4700 and highest are about 9400 m. Image credit: NASA/Goddard/Arizona State University

Photometrically and thermally corrected Level-2 was downloaded from PDS geosciences node (Fig. 4 (a)). The spectral profiles from different locations were collected (fresh/bright exposures are considered as representative pixel) using ENVI image processing software and continuum was removed for better understanding of absorption parameters (Fig. 4 (c)). The sample locations are shown on  $M^3$  image (Fig. 4 (a)). Reflectance spectral profiles (3\*3 pixel average) were collected from different locations of the basin area and

continuum has been removed for reflectance spectra for better understanding absorption spectra characteristic. Unknown sample reflectance spectra were matched with known RELAB spectra (Fig. 4 (b)) [19] for identifying minerals. The comparison was carried out based on pixel-by-pixel to express the similarity between the unknown and known spectra.

#### IV. RESULT

Fig. 4 (c) shows the continuum removed reflectance spectra

of olivine, LCP, HCP, plagioclase and Fe-Mg-spinel minerals. Olivine shows strong absorption band near 1000 nm and weak absorption band near 2000 nm [20]. Near 2000 nm weak absorption band of olivine may be presence of spinel in the olivine. LCP shows absorption bands near 1000 nm and 2000 nm [7], [21]. HCP shows absorption band near 1000 nm and also beyond >2000 nm, the band shift of longer wavelength (>2000 nm) may be due to Ca<sup>+</sup> contribution [7], [20]. Plagioclase shows absorption band near 1250 nm [22], [23]. An atypical strong absorption band have been identified near 2000 nm and relatively weaker band 1000 nm, probably this signature indicates Fe-Mg-spinel minerals present on the surface [e.g., 24, 25].







Fig. 3 Geological map of the Orientale Basin (Basin Interior) [10]



(a)



(b)



(c)

Fig. 4 (a) M<sup>3</sup> coverage of the study area. Reflectance sample locations are plotted on the M<sup>3</sup> image. 1-Plagioclase, 2-Low-Ca pyroxene (LCP), 3-Fe-Mg-Spinel, 4-Olivine and 5-High-Ca pyroxene (LCP), (b) RELAB reflectance spectral data, and (c) Continuum removed reflectance spectra of 1-Plagioclase, 2-Low-Ca pyroxene (LCP), 3-Fe-Mg-Spinel, 4-Olivine and 5-High-Ca pyroxene (LCP) minerals

## V. CONCLUSION

Reflectance imaging spectroscopy data has allowed mapping/identifying minerals in the Orientale basin region on the Moon surface. Olivine, Low-Ca pyroxene (LCP), High-Ca pyroxene (LCP), plagioclase and Fe-Mg-spinel minerals were identified. This study put forward that Reflectance imaging spectroscopy (Hyperspectral imaging) is suitable for the mineral mapping on the lunar surface.

## ACKNOWLEDGMENT

Authors wish to acknowledge M<sup>3</sup> and RELAB team for providing the data. First author thankfully acknowledges C-DAC for providing resource to carry out the study. The authors are grateful to the Editor and anonymous reviewers for their comments and suggestion for improving the article.



## REFERENCES

- [1] Burns, R. G., Mineralogical applications of crystal field theory. Cambridge University Press, 1993, vol. 5, pp.220-242.
- [2] Sivakumar, V. and Neelakantan R., Mineral mapping of lunar highland region using Moon Mineralogy Mapper (M<sup>3</sup>) hyperspectral data, Geological Society of India, 2015, vol. 86 (5), pp.513–518.
- [3] Sivakumar, V., Neelakantan, R. and Biju, C., An analysis of mineral composition on the Moon surface using Moon orbital data (hyperspectral imaging spectrometer data), Geological Society of India, 2015, (In Press).
- [4] Arivazhagan, S., Quantitative Characterization of Lunar Mare Orientale Basalts Detected by Moon Mineralogy Mapper on Chandrayaan-1. In Planetary Exploration and Science: Recent Results and Advances, Springer Berlin Heidelberg, 2015, pp.21-43.
- [5] Anbazhagan, S., & Arivazhagan, S., Reflectance spectra of analog anorthosites: Implications for lunar highland mapping. Planetary and Space Science, 2010, vol. 58(5), pp.752-760.
- [6] Klima, R. L., Pieters, C. M., Boardman, J. W., Green, R. O., Head, J. W., Isaacson, P. J., ... & Tompkins, S., New insights into lunar petrology: Distribution and composition of prominent low-Ca pyroxene exposures as observed by the Moon Mineralogy Mapper (M<sup>3</sup>). Journal of Geophysical Research: Planets (1991–2012), 2011, vol. 116(E6), pp.21-29.
- [7] Adams, J. B., Visible and near-infrared diffuse reflectance spectra of pyroxenes as applied to remote sensing of solid objects in the solar system, Journal of Geophysical Research, 1974, vol.79(32), pp.4829–4835.
- [8] Wilhelms, D. E., McCauley, J. F. and Trask, N. J., The geologic history of the Moon, Washington: USGPO; Denver, CO (Federal Center, Box 25425, Denver 80225): US Geological Survey, 1987, vol. 1348, pp.302.
- [9] Spudis, P. D., The Geology of Multi-ring Basins: The Moon and Other Planets, Cambridge University Press, New York and Cambridge, 1993, pp.263.
- [10] Spudis, P. D., and Ryder G., Apollo 17 impact melts and their relation to the Serenitatis Basin. In multi-ring basins, Proc. Lunar Planet. Sci. Conf., 1981, vol.12A, pp.133–148.
- [11] McCauley, J. F., Orientale and Caloris, Phys. Earth Planet. Inter., 1977, vol.15, pp.220–250.
- [12] Gillis, J. J., The composition and geologic setting of mare deposits on the far side of the Moon, PhD dissertation, Rice Univ., Houston TX, 1998, pp.257.
- [13] Spudis, P. D., B. R. Hawke, and Lucey P., Composition of Orientale Basin deposits and implications for the lunar basin-forming process. Proceedings, lunar and planetary science conference 15, J. Geophys. Res., 1984, vol.89, pp.C197–C210.
- [14] Bussey, D. B. J., and Spudis P. D., Compositional analysis of the Orientale Basin using full resolution Clementine data: Some preliminary results, Geophys. Res. Lett., 1997, vol.24, pp.445–448.
- [15] Cheek, L. C., Donaldson Hanna K. L., Pieters C. M., Head J. W., and Whitten J. L., Distribution and purity of anorthosite across the Orientale Basin: New perspectives from Moon Mineralogy Mapper data, J. Geophys. Res. Planets, 2013, vol.118, pp.1805–1820.
- [16] Goswami, J. N. and Annadurai, M., Chandrayaan-1: India's first planetary science mission to the Moon, Lunar Planetary Science (CDROM), Abstract, 2009, vol. 40, pp.2571.
- [17] Pieters, C., Boardman, J., Buratti, B., Chatterjee, A., Clark, R., Glavich, T., Green, R., Head III, J., Isaacson, P., Malaret, E., et al., The Moon Mineralogy Mapper (M<sup>3</sup>) on Chandrayaan-1, Current Science, 2009, vol. 96, pp.500–505.
- [18] Diviner Global Composition map by NASA/GSFC/UCLA/JPL, 2010. [http://www.nasa.gov/images/content/482363main\\_v\\_06\\_b\\_greenhagen\\_sml.jpg](http://www.nasa.gov/images/content/482363main_v_06_b_greenhagen_sml.jpg), accessed on October, 2015.
- [19] Pieters, C.M., Compositional diversity and stratigraphy of the lunar crust derived from Reflectance spectroscopy, Remote Geochemical Analysis, Cambridge University Press, 1993, vol.14, pp.309-339.
- [20] Borst, A. M., Foing, B. H., Davies, G. R., and Van Westrenen, W., (Surface mineralogy and stratigraphy of the lunar South Pole-Aitken basin determined from Clementine UV/VIS and NIR data, Planetary and Space Science, 2012, vol.68(1), pp.76-85.
- [21] Klima, R. L. and Pieters, C. M., MGM analysis of pyroxene mineral separates from Apollo 15 and 17, Lunar and Planetary Science Conference, Lunar and Planetary Institute Technical Report, 2008, vol.39 pp.1756.
- [22] Adams, J. B. and Goullaud, L. H., Plagioclase feldspars: Visible and near infrared diffuse reflectance spectra as applied to remote sensing, in Lunar Planetary Science Conference, 1978, vol.9, pp.1–3.
- [23] Conel, J.E. and Nash, D.B., Spectral reflectance and albedo of Apollo 11 lunar samples: Effects of irradiation and vitrification and comparison with telescopic observations, Proceedings of the Apollo 11 Lunar Science Conference, 1970, vol.3, pp.2013–2024.
- [24] Cloutis, E. A., Sunshine, J. M., and Morris, R. V. Spectral reflectance-compositional properties of spinels and chromites: Implications for planetary remote sensing and geothermometry, Meteoritics and Planetary Science, 2004, vol.39(4), pp.545-566.
- [25] Pieters, C. M., Hanna, K. D., Cheek, L., Dhingra, D., Prissel, T., Jackson, C., ... & Taylor, L. A., The distribution of Mg-spinel across the Moon and constraints on crustal origin, The Second Conference on the Lunar Highlands Crust and New Directions, American Mineralogist, 2014, vol.99(10), pp.1893-1910.



**Mr. Sivakumar** is a part time research scholar in Department of Geology, Periyar University, Salem, India and presently working as a Senior Technical Officer in Centre for Development of Advanced Computing (C-DAC). He received his Masters of Technology (M.Tech.) degree in Geological Remote Sensing & Geoinformatics at Bharathidasan University, Tamil Nadu, India and Masters of Science (M.Sc.) degree in Geology at University of Madras, India. Prior to C-DAC he was associated with Indian Space Research Organization – ISRO (NRSC) as a research fellow. Mr. Sivakumar's scientific accomplishment includes more than 25 journal publications, books and conference contributions. He is also associated with national/international professional bodies in various roles. He received many awards/honors such as Best Industrial Researcher, Best Technological Research, Best Paper Publication, Best Faculty, research fellowship, merit scholarship, etc. He is involved in many research & funded projects, business lead, project planning & designing, decision making and research project initiation. He is active member in various geoscience/geospatial/research forums in providing emerging solutions, serving as regional/associate editor, editorial board member, etc. His research areas include natural resource management, planetary science studies, remote sensing and GIS, computational sciences, high performance computing (HPC) - satellite data processing & information extraction, natural disaster management, etc. He teaches & visits various academic & research institutes across the India. He has mentored various postgraduate and undergraduate students from various fields.



**Prof. Dr. R. Neelakantan** is Professor of the Department of Industries and Earth Sciences at Tamil University, Thanjavur, Tamil Nadu, India, where he has been since 2009. He also currently serves as Assistant Professor (on-lien) in Department of Geology at Periyar University, Salem, India. He received a M.Sc. in Geology from Annamalai University, Chidambaram and M.Tech in Remote Sensing from Bharathidasan University, Tiruchirappalli. He received his Ph.D. in Earth Sciences from Bharathidasan University. From 1991 to 2003, he was associated with various institutes in various roles such as Research Scientist, Research Associate, CSIR Senior Research Fellow, etc. He has mentored various Postdoctoral, postgraduate and undergraduate students from various fields. He has coauthored numerous (more than 70) highly cited journal publications, conference articles and book chapters in the multidisciplinary topics, and has received several awards and grants from various funding agencies including the DST young Scientist Fellowship 2006-2009, C.S.I.R Senior Research Fellowship and Consultant Geologist for the State Government of Tamil Nadu. He is active member in various academic activities in different institutes. He has strong administrative experience in high level decision making bodies for academic purpose. He has organized numerous conference and workshops and delivered numerous expert talks in various institutes across the country. His research interest includes natural disaster management (landslide studies), environmental geomorphology, water resources management and planetary science.