

Regular Generalized Star Star closed sets in Bitopological Spaces

K. Kannan, D. Narasimhan, K. Chandrasekhara Rao and R. Ravikumar

Abstract—The aim of this paper is to introduce the concepts of $\tau_1\tau_2$ -regular generalized star star closed sets, $\tau_1\tau_2$ -regular generalized star star open sets and study their basic properties in bitopological spaces.

Keywords— $\tau_1\tau_2$ -regular closed sets; $\tau_1\tau_2$ -regular open sets; $\tau_1\tau_2$ -regular generalized closed sets; $\tau_1\tau_2$ -regular generalized star closed sets; $\tau_1\tau_2$ -regular generalized star star closed sets.

I. INTRODUCTION

IN 1963, J.C. Kelly [11] initiated the study of bitopological spaces as a natural structure by studying quasi metrics and its conjugate. This structure is a richer structure than that of a topological space. Considerable effort had been expended in obtaining appropriate generalizations of standard topological properties to bitopological category by various authors. Most of them deal with the theory itself but very few with applications.

K. Chandrasekhara Rao and N. Palaniappan [6] introduced the concepts of regular generalized star closed sets and regular generalized star open sets in a topological space and they are extended to bitopological settings by K. Chandrasekhara Rao and K. Kannan [1]. On the other hand Chandrasekhara Rao and N. Palaniappan [7] introduced the concept of regular generalized star star closed sets and regular generalized star star open sets in topological spaces and study their properties. In this sequel, we introduce the concepts of $\tau_1\tau_2$ -regular generalized star star closed sets ($\tau_1\tau_2$ -rg* closed sets) and $\tau_1\tau_2$ -regular generalized star star open sets ($\tau_1\tau_2$ -rg* open sets) and study their basic properties in bitopological spaces.

Throughout this paper, (X, τ_1, τ_2) or simply X denote a bitopological space. The intersection (resp. union) of all τ_i -closed sets containing A (resp. τ_i -open sets contained in A) is called the τ_i -closure (resp. τ_i -interior) of A , denoted by $\tau_i\text{-cl}(A)$ {resp. $\tau_i\text{-int}(A)$ }. The closure and interior of B relative to A with respect to the topology τ_i are written as $\tau_i\text{-cl}_A(B)$ and $\tau_i\text{-int}_A(B)$ respectively.

For any subset $A \subseteq X$, $\tau_i\text{-rint}(A)$ and $\tau_i\text{-rcl}(A)$ denote the regular interior and regular closure of a set A with respect to the topology τ_i respectively. The regular closure and regular interior of B relative to A with respect to the topology τ_i are written as $\tau_i\text{-rcl}_A(B)$ and $\tau_i\text{-rint}_A(B)$ respectively. The set of all τ_2 -regular closed sets in X is denoted by $\tau_2\text{-R.C}(X, \tau_1, \tau_2)$. The set of all $\tau_1\tau_2$ -regular open sets in X is denoted by

$\tau_1\tau_2\text{-R.O}(X, \tau_1, \tau_2)$. A^C denotes the complement of A in X unless explicitly stated.

We shall require the following known definitions and results:

Definition 1.1: [1] A set A of a bitopological space (X, τ_1, τ_2) is called

- $\tau_1\tau_2$ -regular closed if $\tau_1\text{-cl}[\tau_2\text{-int}(A)] = A$.
- $\tau_1\tau_2$ -regular open if $\tau_1\text{-int}[\tau_2\text{-cl}(A)] = A$.
- $\tau_1\tau_2$ -regular generalized closed ($\tau_1\tau_2$ -rg closed) in X if $\tau_2\text{-cl}(A) \subseteq U$ whenever $A \subseteq U$ and U is $\tau_1\tau_2$ -regular open in X .
- $\tau_1\tau_2$ -regular generalized open ($\tau_1\tau_2$ -rg open) in X if $F \subseteq \tau_2\text{-int}(A)$ whenever $F \subseteq A$ and F is $\tau_1\tau_2$ -regular closed in X .
- $\tau_1\tau_2$ -regular generalized star closed ($\tau_1\tau_2$ -rg* closed) in X if and only if $\tau_2\text{-rcl}(A) \subseteq U$ whenever $A \subseteq U$ and U is $\tau_1\tau_2$ -regular open in X .
- $\tau_1\tau_2$ -regular generalized star open ($\tau_1\tau_2$ -rg* open) in X if and only if its complement is $\tau_1\tau_2$ -regular generalized star closed ($\tau_1\tau_2$ -rg* closed) in X .

Lemma 1.2: [1] Let A be a τ_1 -open set in (X, τ_1, τ_2) and let U be $\tau_1\tau_2$ -regular open in A . Then $U = A \cap W$ for some $\tau_1\tau_2$ -regular open set W in X .

Lemma 1.3: [1] If A is $\tau_1\tau_2$ -open and U is $\tau_1\tau_2$ -regular open in X then $U \cap A$ is $\tau_1\tau_2$ -regular open in A .

II. $\tau_1\tau_2$ -REGULAR GENERALIZED STAR STAR CLOSED SETS

Definition 2.1: A subset A of a bitopological space (X, τ_1, τ_2) is called $\tau_1\tau_2$ -regular generalized star star closed ($\tau_1\tau_2$ -rg** closed) in X if and only if $\tau_2\text{-cl}[\tau_1\text{-int}(A)] \subseteq U$ whenever $A \subseteq U$ and U is $\tau_1\tau_2$ -regular open in X .

Example 2.2: Let $X = \{a, b, c\}$, $\tau_1 = \{\phi, X, \{a\}\}$, $\tau_2 = \{\phi, X, \{a\}, \{b, c\}\}$. Then the set of all subsets $P(X)$ of X are $\tau_1\tau_2$ -rg** closed sets in (X, τ_1, τ_2) .

Theorem 2.3: Let A be a subset of a bitopological space (X, τ_1, τ_2) . If A is $\tau_1\tau_2$ -rg** closed then $\tau_2\text{-cl}[\tau_1\text{-int}(A)] - A$ does not contain non empty $\tau_1\tau_2$ -regular closed sets.

Proof: Suppose that A is $\tau_1\tau_2$ -rg** closed. Let F be a $\tau_1\tau_2$ -regular closed set such that $F \subseteq \tau_2\text{-cl}[\tau_1\text{-int}(A)] - A$. Then $F \subseteq \tau_2\text{-cl}[\tau_1\text{-int}(A)] \cap A^C$. Since $F \subseteq A^C$, we have $A \subseteq F^C$. Since F is $\tau_1\tau_2$ -regular closed set, we have F^C is $\tau_1\tau_2$ -regular open. Since A is $\tau_1\tau_2$ -rg** closed, we have $\tau_2\text{-cl}[\tau_1\text{-int}(A)] \subseteq F^C$. Therefore, $F \subseteq [\tau_2\text{-cl}[\tau_1\text{-int}(A)]]^C$. Also $F \subseteq \tau_2\text{-cl}[\tau_1\text{-int}(A)]$. Hence $F \subseteq \phi$. Therefore, $F = \phi$. ■

Theorem 2.4: If A is $\tau_1\tau_2$ -rg** closed and B is $\tau_1\tau_2$ -g closed, then $A \cup B$ is $\tau_1\tau_2$ -rg** closed.

Proof: Let $A \cup B \subseteq U$ and U is $\tau_1\tau_2$ -regular open in X . Since $A \subseteq U$ and A is $\tau_1\tau_2$ -rg** closed, we have

K. Kannan, D. Narasimhan, K. Chandrasekhara Rao are with Department of Mathematics, Srinivasa Ramanujan Centre, SASTRA University, Kumbakonam, India, E-Mail: anbukannan@rediffmail.com, dnsastra@rediffmail.com and k.chandrasekhara@rediffmail.com

R. Ravikumar is with Department of Mathematics, A.M.A. College of Engineering, Vadavandam, India, E-Mail: dr.ravikumar@rediffmail.com

$\tau_2\text{-cl}[\tau_1\text{-int}(A)] \subseteq U$. Since $B \subseteq U$ and B is $\tau_1\tau_2\text{-}g$ closed, we have $\tau_2\text{-cl}(B) \subseteq U$. Now, $\tau_2\text{-cl}[\tau_1\text{-int}(A \cup B)] \subseteq \tau_2\text{-cl}[\tau_1\text{-int}[A \cup \tau_2\text{-cl}(B)]] \subseteq U$. Therefore, $A \cup B$ is $\tau_1\tau_2\text{-}rg^{**}$ closed. ■

Theorem 2.5: If a subset A is $\tau_1\tau_2\text{-}rg$ closed then A is $\tau_1\tau_2\text{-}rg^{**}$ closed.

Proof: Let $A \subseteq U$ and U is $\tau_1\tau_2\text{-}rg$ open. Since A is $\tau_1\tau_2\text{-}rg$ closed, we have $\tau_2\text{-cl}(A) \subseteq U$. Hence $\tau_2\text{-cl}[\tau_1\text{-int}(A)] \subseteq U$. Therefore, A is $\tau_1\tau_2\text{-}rg^{**}$ closed. ■

Definition 2.6: Let $B \subseteq Y \subseteq X$. A subset B of Y is said to be $\tau_1\tau_2\text{-}rg^{**}$ closed relative to Y if B is $\tau_1\tau_2\text{-}rg^{**}$ closed in the subspace Y .

Theorem 2.7: Let $(Y, \tau_{1/Y}, \tau_{2/Y})$ be a subspace of (X, τ_1, τ_2) . Suppose that a subset B of Y is $\tau_1\tau_2\text{-}rg^{**}$ closed relative to $(Y, \tau_{1/Y}, \tau_{2/Y})$ and Y is $\tau_1\tau_2\text{-}open$ and $\tau_1\tau_2\text{-}g$ closed in (X, τ_1, τ_2) then B is $\tau_1\tau_2\text{-}rg^{**}$ closed in (X, τ_1, τ_2) .

Proof: Let $B \subseteq U$ and U is $\tau_1\tau_2\text{-}rg$ open in X . Since Y is $\tau_1\tau_2\text{-}open$ in X , we have $U \cap Y$ is $\tau_1\tau_2\text{-}rg$ open in Y {by Lemma 1.3} and $B \subseteq U \cap Y$. Since B is $\tau_1\tau_2\text{-}rg^{**}$ closed relative to $(Y, \tau_{1/Y}, \tau_{2/Y})$, we have $\tau_2\text{-cl}_Y[\tau_1\text{-int}_Y(B)] \subseteq U \cap Y$. Hence $\tau_2\text{-cl}[\tau_1\text{-int}(B)] \cap Y \subseteq U \cap Y$. Let $G = U \cup \{X - \tau_2\text{-cl}[\tau_1\text{-int}(B)]\}$. Then G is $\tau_1\text{-open}$ and $Y \subseteq G$. Since Y is $\tau_1\tau_2\text{-}g$ closed, we have $\tau_2\text{-cl}(Y) \subseteq G$. Now, $\tau_2\text{-cl}[\tau_1\text{-int}(B)] = \tau_2\text{-cl}(Y) \cap \tau_2\text{-cl}[\tau_1\text{-int}(B)] \subseteq G \cap \tau_2\text{-cl}[\tau_1\text{-int}(B)] = U \cap \tau_2\text{-cl}[\tau_1\text{-int}(B)] \cup \phi \subseteq U$. Therefore, B is $\tau_1\tau_2\text{-}rg^{**}$ closed in (X, τ_1, τ_2) . ■

Theorem 2.8: Suppose that a subset B of Y is $\tau_1\tau_2\text{-}rg^{**}$ closed in (X, τ_1, τ_2) and Y is $\tau_1\tau_2\text{-}open$ in (X, τ_1, τ_2) then B is $\tau_1\tau_2\text{-}rg^{**}$ closed relative to $(Y, \tau_{1/Y}, \tau_{2/Y})$.

Proof: Let $B \subseteq U$ and U is $\tau_1\tau_2\text{-}rg$ open in Y . Since Y is $\tau_1\text{-open}$, we have $U = Y \cap W$ for some $\tau_1\tau_2\text{-}rg$ open set W in (X, τ_1, τ_2) {by Lemma 1.2}. Since $B \subseteq Y \cap W \subseteq W$ and B is $\tau_1\tau_2\text{-}rg^{**}$ closed in (X, τ_1, τ_2) , we have $\tau_2\text{-cl}[\tau_1\text{-int}(B)] \subseteq W$. Therefore, $\tau_2\text{-cl}_Y[\tau_1\text{-int}_Y(B)] = \tau_2\text{-cl}[\tau_1\text{-int}(B)] \cap Y \subseteq W \cap Y = U$. Hence B is $\tau_1\tau_2\text{-}rg^{**}$ closed relative to Y . ■

Theorem 2.9: Let A and B be subsets such that $A \subseteq B \subseteq \tau_2\text{-cl}[\tau_1\text{-int}(A)]$. If A is $\tau_1\tau_2\text{-}rg^{**}$ closed, then B is $\tau_1\tau_2\text{-}rg^{**}$ closed.

Proof: Let $B \subseteq U$ and U is $\tau_1\tau_2\text{-}rg$ open in X . Since $A \subseteq B$, we have $A \subseteq U$. Since A is $\tau_1\tau_2\text{-}rg^{**}$ closed, we have $\tau_2\text{-cl}[\tau_1\text{-int}(A)] \subseteq U$. Since $B \subseteq \tau_2\text{-cl}[\tau_1\text{-int}(A)]$, we have $\tau_2\text{-cl}[\tau_1\text{-int}(B)] \subseteq \tau_2\text{-cl}(B) \subseteq \tau_2\text{-cl}[\tau_1\text{-int}(A)] \subseteq U$. Therefore, B is $\tau_1\tau_2\text{-}rg^{**}$ closed. ■

Theorem 2.10: Suppose that $\tau_1\tau_2\text{-}R.O(X, \tau_1, \tau_2) \subseteq \tau_2\text{-}C(X, \tau_1, \tau_2)$. Then every subset of X is $\tau_1\tau_2\text{-}rg^{**}$ closed.

Proof: Let A be a subset of X . Let $A \subseteq U$ and U is $\tau_1\tau_2\text{-}rg$ open in X . Since $\tau_1\tau_2\text{-}R.O(X, \tau_1, \tau_2) \subseteq \tau_2\text{-}C(X, \tau_1, \tau_2)$, we have U is $\tau_2\text{-closed}$ in X . Since $A \subseteq U$, we have $\tau_2\text{-cl}(A) \subseteq \tau_2\text{-cl}(U) = U$. Therefore, $\tau_2\text{-cl}[\tau_1\text{-int}(A)] \subseteq \tau_2\text{-cl}(A) \subseteq U$. Hence A is $\tau_1\tau_2\text{-}rg^{**}$ closed. ■

III. $\tau_1\tau_2\text{-REGULAR GENERALIZED STAR STAR OPEN SETS}$

Definition 3.1: A subset A of a bitopological space (X, τ_1, τ_2) is called $\tau_1\tau_2\text{-regular generalized star star open}$

$(\tau_1\tau_2\text{-}rg^{**} \text{ open})$ in X if and only if its complement is $\tau_1\tau_2\text{-regular generalized star star closed}$ ($\tau_1\tau_2\text{-}rg^{**} \text{ closed}$) in X .

Example 3.2: Let $X = \{a, b, c\}$, $\tau_1 = \{\phi, X, \{a\}\}$, $\tau_2 = \{\phi, X, \{a\}, \{b, c\}\}$. Then the set of all subsets $P(X)$ are $\tau_1\tau_2\text{-}rg^{**}$ open sets in (X, τ_1, τ_2) .

A necessary and sufficient condition for a set A to be a $\tau_1\tau_2\text{-}rg^{**}$ open set is obtained in the next theorem.

Theorem 3.3: A subset A of a bitopological space (X, τ_1, τ_2) is $\tau_1\tau_2\text{-}rg^{**}$ open if and only if $F \subseteq \tau_2\text{-int}[\tau_1\text{-cl}(A)]$ whenever $F \subseteq A$ and F is $\tau_1\tau_2\text{-regular closed}$ in X .

Proof: Necessity: Let $F \subseteq A$ and F is $\tau_1\tau_2\text{-regular closed}$ in X . Then $A^C \subseteq F^C$ and F^C is $\tau_1\tau_2\text{-regular open}$ in X . Since A is $\tau_1\tau_2\text{-}rg^{**}$ open, we have A^C is $\tau_1\tau_2\text{-}rg^{**}$ closed. Hence, $\tau_2\text{-cl}[\tau_1\text{-int}(A^C)] \subseteq F^C$. Consequently, $[\tau_2\text{-int}[\tau_1\text{-cl}(A)]]^C \subseteq F^C$. Therefore, $F \subseteq \tau_2\text{-int}[\tau_1\text{-cl}(A)]$.

Sufficiency: Let $A^C \subseteq U$ and U is $\tau_1\tau_2\text{-regular open}$ in X . Then $U^C \subseteq A$ and U^C is $\tau_1\tau_2\text{-regular closed}$ in X . By our assumption, we have $U^C \subseteq \tau_2\text{-int}[\tau_1\text{-cl}(A)]$. Hence $[\tau_2\text{-int}[\tau_1\text{-cl}(A)]]^C \subseteq U$. Therefore, $\tau_2\text{-cl}[\tau_1\text{-int}(A^C)] \subseteq U$. Consequently A^C is $\tau_1\tau_2\text{-}rg^{**}$ closed. Hence A is $\tau_1\tau_2\text{-}rg^{**}$ open. ■

Theorem 3.4: Let A and B be subsets such that $\tau_2\text{-int}[\tau_1\text{-cl}(A)] \subseteq B \subseteq A$. If A is $\tau_1\tau_2\text{-}rg^{**}$ open, then B is $\tau_1\tau_2\text{-}rg^{**}$ open.

Proof: Let $F \subseteq B$ and F is $\tau_1\tau_2\text{-regular closed}$ in X . Since $B \subseteq A$, we have $F \subseteq A$. Since A is $\tau_1\tau_2\text{-}rg^{**}$ open, we have, $F \subseteq \tau_2\text{-int}[\tau_1\text{-cl}(A)]$ {by Theorem 3.3}. Since $\tau_2\text{-int}[\tau_1\text{-cl}(A)] \subseteq B$, we have $\tau_2\text{-int}[\tau_2\text{-int}[\tau_1\text{-cl}(A)]] \subseteq \tau_2\text{-int}(B) \subseteq \tau_2\text{-int}[\tau_1\text{-cl}(B)]$. Hence $F \subseteq \tau_2\text{-int}[\tau_1\text{-cl}(A)] \subseteq \tau_2\text{-int}[\tau_1\text{-cl}(B)]$. Therefore, B is $\tau_1\tau_2\text{-}rg^{**}$ open. ■

Theorem 3.5: If a subset A is $\tau_1\tau_2\text{-}rg^{**}$ closed, then $\tau_2\text{-cl}[\tau_1\text{-int}(A)] - A$ is $\tau_1\tau_2\text{-}rg^{**}$ open.

Proof: Let $F \subseteq \tau_2\text{-cl}[\tau_1\text{-int}(A)] - A$ and F is $\tau_1\tau_2\text{-regular closed}$. Since A is $\tau_1\tau_2\text{-}rg^{**}$ closed, we have $\tau_2\text{-cl}[\tau_1\text{-int}(A)] - A$ does not contain nonempty $\tau_1\tau_2\text{-regular closed}$ {by Theorem 2.3}. Therefore, $F = \phi$. Hence $\tau_2\text{-cl}[\tau_1\text{-int}(A)] - A$ is $\tau_1\tau_2\text{-}rg^{**}$ open. ■

REFERENCES

- [1] K. Chandrasekhara Rao and K. Kannan, *Regular generalized star closed sets in bitopological spaces*, Thai Journal of Mathematics, Vol 4, (2) (2006), 341–349.
- [2] K. Chandrasekhara Rao and K. Kannan, *Generalized star regular closed sets in bitopological spaces*, Antarctica Math. J., 4, (2)(2007), 139–146.
- [3] K. Chandrasekhara Rao, K. Kannan and D. Narasimhan, *Generalized star star regular closed sets*, Natl. Acad. Sci. Lett., Vol.31, No. 1 & 2, (2008), 31–33.
- [4] K. Chandrasekhara Rao and M. Mariasingam, *On Bitopological spaces*, Acta Ciencia Indica, Vol. XXVI M (4) (2000), 283–288.
- [5] K. Chandrasekhara Rao and N. Palaniappan, *Regular generalized closed sets*, Kyungpook Mathematical Journal, 33(2)(1993) 211–219.
- [6] K. Chandrasekhara Rao and N. Palaniappan, *Regular generalized star closed sets*, Bulletin of Pure and Applied Sciences, 19(2)(2000), 291–306.
- [7] K. Chandrasekhara Rao and N. Palaniappan, *Ph. D. Thesis*, Alagappa University, (2000).
- [8] T. Fukutake, *On generalized closed sets in bitopological spaces*, Bull. Fukuoka Univ. Ed. Part III, 35 (1986), 19–28.
- [9] T. Fukutake, *Semi open sets on bitopological spaces*, Bull. Fukuoka Uni. Education, 38(3)(1989), 1–7.

- [10] T. Fukutake, P. Sundaram and M. Sheik John, *w-closed sets, w-open sets and w-continuity in bitopological spaces*, Bull. Fukuoka Univ. Ed. Part III, 51 (2002), 1–9.
- [11] J. C. Kelly, *Bitopological spaces*, Proc. London Math. Society, 13 (1963), 71–89.
- [12] M. Rajamani and K. Vishwanathan, *On α -generalized semi closed sets in bitopological spaces*, Bulletin of Pure and Applied Sciences, Vol. 24E (No.1) (2005), 39–53.
- [13] M. Sheik John and P. Sundaram, *g^* -closed sets in Bitopological Spaces*, Indian J. Pure Appl. Math., 35(1) (2004), 71–80.