

Rice cDNA Encoding PROLM is Capable of Rescuing Salt Sensitive Yeast Phenotypes G19 and Axt3K from Salt Stress

Prasad Senadheera , Younousse Saidi , Frans JM Maathuis

Abstract—Rice seed expression (cDNA) library in the Lambda Zap 11® phage constructed from the developing grain 10-20 days after flowering was transformed into yeast for functional complementation assays in three salt sensitive yeast mutants *S. cerevisiae* strain CY162, G19 and Axt3K. Transformed cells of G19 and Axt3K with pYES vector with cDNA inserts showed enhance tolerance than those with empty pYes vector. Sequencing of the cDNA inserts revealed that they encode for the putative proteins with the sequence homologous to rice putative protein *PROLM24* (Os06g31070), a prolamin precursor. Expression of this cDNA did not affect yeast growth in absence of salt. Axt3k and G19 strains expressing the *PROLM24* were able to grow upto 400 mM and 600 mM of NaCl respectively. Similarly, Axt3k mutant with *PROLM24* expression showed comparatively higher growth rate in the medium with excess LiCl (50 mM). The observation that expression of *PROLM24* rescued the salt sensitive phenotypes of G19 and Axt3k indicates the existence of a regulatory system that ameliorates the effect of salt stress in the transformed yeast mutants. However, the exact function of the cDNA sequence, which shows partial sequence homology to yeast *UTR1* is not clear. Although *UTR1* involved in ferrous uptake and iron homeostasis in yeast cells, there is no evidence to prove its role in Na⁺ homeostasis in yeast cells. Absence of transmembrane regions in Os06g31070 protein indicates that salt tolerance is achieved not through the direct functional complementation of the mutant genes but through an alternative mechanism.

Keywords—Rice seed expression, salt stress, prolamin, salinity tolerance, *Oryza sativa*

I. INTRODUCTION

SALINITY tolerance in rice is a multigenic trait involving a variety of physiological and biochemical pathways. Salinity tolerance in rice varies not only across the cultivars but also across the growth stages. It is relatively salt tolerant during germination, active tillering, and towards maturity, but sensitive during early seedling stage and reproductive stage [1,2]. Partitioning of Na⁺ ions alters the concentration of Na⁺ ions in the different tissues of the rice plant in growth stage dependant manner. The higher accumulation rates of Na⁺ than K⁺ in the developing floral parts were observed and may be one of the underlying causes for the sensitivity of the rice to salt stress at panicle induction stage [3]. However, rice grain

development was found to be less sensitive to salinity at milking stage [4]. These observations where in the floral parts show higher sensitivity to Na⁺ but becomes relatively insensitive during subsequent seed development indicate that developing grain gradually acquire the capability to tolerate salt stress. Hence, studies on the expression of genes and proteins in developing rice seed in response to salt stress could pave the way to identify new salt responsive genes along with novel mechanisms of salt tolerance.

Heterologous expression of plant genes in yeast is a commonly used strategy for functional validation of uncharacterized sequences. *Saccharomyces cerevisiae* L. is a good model system for identifying genes involved in the salt tolerance of plants as both organisms have similar transport systems [5]. Besides, yeast has been extensively studied as a host of heterologous membrane proteins of plants. The mechanisms of ion transport at the plasma membrane and vacuolar membrane are relatively well characterized at the molecular level but the overall homeostatic mechanisms which maintain intracellular ion concentrations are still being elucidated in plants and fungi [6, 7, 8].

Sodium is not essential for most fungi and plants and they are unable to tolerate high NaCl concentrations in soils and water [9, 10]. The major deleterious effects of high salinity in yeast are caused by Na⁺ accumulation in the cytoplasm, where many metabolic activities are sensitive to Na⁺ inhibition [11]. Influx of Na⁺ (and Li⁺) in yeast occurs through the K⁺ uptake pathway, of which two component transporters were identified as TRK1 and TRK2 [12, 13]. Exposure to high Na⁺ content makes this TRK1 and TRK2 systems shifted to a high affinity K⁺ uptake system to reduce the Na⁺ influx [14]. The efflux of Na⁺ is mainly mediated by a putative P-type ATPase encoded by the tandem array of four closely related genes ENA1-4 [15, 16 17]. ENA2, ENA3, and ENA4 present a very weak constitutive expression, while ENA1, although poorly expressed on basal conditions, is induced upon exposure to high concentrations of Li⁺ or Na⁺, as well as to alkaline pH. Therefore, deletion of ENA1 results in cells hypersensitive to Na⁺ or Li⁺ [18, 19]. Plasma membrane localized Na⁺/H⁺ antiport protein, encoded by NHA1, seems to play a role in sodium efflux [20, 21]. This protein though has a little effect on Na⁺ efflux, may be important in survival under acidic external pH. In addition to the Na influx and efflux systems, the product of *ScNHX1*, which is homologous to plant *NHX*, has been shown to mediate sequestration of Na⁺ within an intracellular compartments, particularly in the vacuole [22].

1. Department of Botany, The Open University of Sri Lanka, Nawala, Sri Lanka. * (TP: 0094712279486 E-Mail: spsen@ou.ac.lk)

2. Biology Department, Area 9, University of York, York, YO10 5DD, UK

3. School of Biosciences, University of Birmingham, Birmingham, B15 2TT, UK

Several proteins that are not directly involved with Na⁺ and Li⁺ transport but modulate the function of monovalent cation transporters in yeast have been identified. HAL4/SAT4 and HAL5, that encode homologous protein kinases are implicated in the regulation of cation uptake. Overexpression of these genes increases the tolerance of yeast cells to sodium and lithium, whereas gene disruptions result in greater cation sensitivity. Hal4 and Hal5 protein kinases are found to be involved in the regulation of TRK1-TRK2 potassium transporters and the stabilization of the transporters in the plasma membrane [23, 24]. The genes of trehalose pathway, phosphoglucomutase 2 (PGM2) and hexokinase 2 (HXK2) was shown to affect the TRK activity through the modulation of glucose phosphates (glc-1-P and glc-6-P) [25]. A protein phosphatase, calcineurin, has also been reported to be required for the activation of TRK1 and TRK2 in presence as well as absence of sodium stress [14, 26]. Other proteins involved in the regulation of potassium transport include the protein kinase SKY1 [27]; the osmo-induced protein HAL1 [28]; Casen Kinase (CK-2) and G protein of the RAS superfamily, ARL1 [29]. In addition, ENA1 expression is modulated by the HAL3-PPZ1 regulatory subunit-protein phosphatase pair [30, 31]

Considerable progress in understanding plant salt tolerance at a cellular level has been made through heterologous expression studies in yeast. Functional complementation screening of wheat cDNA library led to identify *HKT1* transporter [32]. Screening of cDNA library prepared from 14-day-old seedlings of *Oryza sativa* cv. Akibare identified 1-pyrroline-5-carboxylate (P5C) synthetase (cOsP5CS), an enzyme involved in the biosynthesis of proline [33]. A rice cation transporter gene *OsKAT* involved with salt stress tolerance was identified through functional complementation of full-length cDNA expression library in salt sensitive G19 phenotype of yeast (*Saccharomyces cerevisiae*) [34]. Salt sensitive yeast mutant for vacuolar antiporter was used in characterising the *OsNHX* phenotype which is found to play an important role in the compartmentation of Na⁺ and K⁺ into the vacuoles and confers salt tolerance [35].

The objective of this study is to screen rice cDNA library constructed from developing rice grain (*Oryza sativa* L. variety 'Kinamaze') in heterologous expression system of yeast mutants to identify novel gene candidates involved with salt tolerance. Expression of Rice prolamin precursor protein encoded by Os06g0507200 was found to be capable of conferring salt tolerance in two yeast mutants hypersensitive to salt.

II. MATERIAL AND METHOD

A. Amplification of rice seed cDNA library in lambda phage

Rice seed expression (cDNA) library in the Lambda Zap 11® phage constructed from the developing grain 10-20 days after flowering was received from Prof Toshihiro Kumamaru of the Kyushu University, Fukuoka, 812-8581 of Japan.

This rice seed expression library in the Lambda Zap 11® phage was amplified in the *E. coli* strain XL Blue MRF⁺ cells

(e14-(McrA-) Δ(mcrCB-hsdSMR-mrr)171 sbcC recB recJ uvrC umuC::Tn5 (Kan^r) lac gyrA96 relA1 thi-1 endA1 λR [F' proAB lacIqZΔM15] Su-) and the titre was estimated of both the cDNA library and ExAssist® helper subsequently. Efficacy of the cDNA library was determined by blue white colony selection using X-gal and IPTG before carrying out the mass excision. Co-infection of XL-Blue MRF⁺ cells with Lambda phage and ExAssist Helper phage resulted in the in vivo excision of phagemid pBluescript® which is then transformed into the *E. coli* strain SOLR (Δ(mcrA)183 Δ(mcrCB-hsdSMR-mrr)173 endA1 supE44 thi-1recA1 gyrA96 relA1 lac [F' proAB lacIqZΔM15 Tn10 (Tetr)]) for amplification. All these steps were carried out following the protocols given in the Lambda ZAP® II Undigested Vector Kit (Cat no 236201; revision A) from the (Stratagene Europe: www.stratagene.com).

B. Functional complementation assays using *Saccharomyces cerevisiae*

If Rice cDNA inserts in the pBluescript were subcloned into the XhoI and EcoRI restriction sites of pYES2 rice expression vector (Invitrogen, California, USA) under control of the inducible Gal1-promoter [36]. Prior to subcloning, pBluescript vector (3 Kb) backbone was removed by gel purification after digestion with XhoI and EcoRI enzymes. cDNA inserts resulted from the gel purification was ligated to pYes using the T4 ligation kit (Fermentas, Maryland, USA) after the dephosphorylation with Antarctic Phosphatase (New England BioLabs, UK). Thus, transformed pYES2 vector was used for yeast functional complementation assays. *S. cerevisiae* strain CY162 (MATa, trk1, trk2::pCK64, his3, ura3), G19 (MATa, his3, ura3, trp1, ade2, and ena1::HIS3::ena4), Axt3K strain (Den1THIS3Tena4, Dnha1TLEU2, Dnhx1TKanMX4) and the parent strain of G19, W303 (MATa, his3, leu2, ura3, trp1, and ade2), were transformed with the pYES vector with cDNA inserts for complementation assay. Yeast transformations were performed using LiCl method as described by Chen et al. (37). Transformed yeast was plated in Synthetic defined (SD) medium (Galactose 20 g/L, Ammonium Sulphate 5 g/L, Agar bacteriological, 0.67% yeast nitrogen base, 20 g/L adjusted to pH 6) with all amino acids except pyrimidine base uracil with different concentrations of K⁺ and Na⁺. Histidine was also removed in addition to uracil in the medium that was used for G19 and Axt3K. CY 162 was screened in media containing 0.1 mM KCl upto 50 mM KCl and the control was 100 mM KCl. Axt3K was grown in series dilution of NaCl upto 500 mM, 100 mM LiCl while G19 was grown up to 700 mM NaCl.

C. PCR amplification of the cDNA inserts in the positively complemented clones and sequencing

pYES vector of the yeast clones that showed positive complementation was isolated from yeast using Qiagen plasmid miniprep (Qiagen, West Sussex, UK) protocol after pre-treating the yeast cultures in Zymolase (AMS Biotechnology, Abingdon, U.K.). The isolated plasmids were retransformed into the *E. coli* strain top 10 and cultivated in

the LB- Ampicilline Agar broth overnight in the 27 °C. Qiagen miniprep™ kit was used to extract plasmid DNA. The cDNA inserts in each vector were amplified with forward (TTCTAATTCGTAGTTTTTCAAGTTCT) and reverse primer(GAAAAAACCCTGGATCGGACTACTAGCAGCTG) flanking the cloning site (94 °C for 30 seconds, 58 °C for 30 seconds and 72 °C for 3 min). The cDNA inserts that showed reliable results after repeated screening in yeast was sequenced for the functional characterization of the cDNA insert at the Biology Technology Facility, University of York, UK.

D. Computer analyses and annotation

To understand and assign functional classes to the sequence information determined in this screen, computational searches were performed against sequence databases at NCBI (<http://www.ncbi.nlm.nih.gov/blast>) TIGR plant transcript assemblies (http://tigrblast.tigr.org/euk-blast/plantta_blast.cgi) and yeast genome database (<http://www.yeastgenome.org>) using the BLAST algorithm (Altschul et al., 1990).

III. RESULTS AND DISCUSSION

A. Amplification and mass excision of cDNA library from Lambda Zap 11

Rice cDNA library in the Lambda Zap 11 vector system (Stratagene, La Jolla, USA) was constructed from the mRNA of the developing grain of *Oryza sativa* L. variety 'Kinamazé'




Fig. 1 Estimating the titre and efficacy of the rice seed cDNA library. The *E. coli* strain, XL Blue MRF' ($OD_{600} = 0.5$) was mixed with Lambda Zap library in 1:1 and 1:10 dilution and incubated at 37 °C for 12 hours in NZY agar plates. IPTG and X-gal were added to the media for determining the efficacy. Plaque-forming units per milliliter (pfu/ml) was counted after 12 hours of incubation

10-20 days after flowering. Titre and the efficacy of the cDNA library were estimated following the protocols provided in the Stratagene Lambda ZAP 11 manual. Titre of the Lambda phage was 418000 pfu/ml and nearly 99% of efficacy in terms of insertions as was seen with blue white colony selection (Fig 1). These parameters indicated high quality of the cDNA library which used for the screening.

B. Screening rice cDNA library in yeast mutant sensitive to salt stress

Yeast mutants transformed with empty pYES vector and pYES with rice cDNA clones were cultured in the SD – Uracil-Histidine plates with different Na^+ and K^+ concentrations under inducible conditions. Testing for uracil and histidine prototrophy minimize the chance for spontaneous reversion of yeast mutants for tolerance. Of the three mutant strains

screened (G19, Axt3K and CY162), growth complementation was seen only in G19 and Axt3K for salt (Na^+) tolerance. Transformed cells of G19 and Axt3K with pYES vector with cDNA inserts showed enhanced tolerance than those with empty pYes vector (Fig 2). G19 and Axt3K are found to be sensitive to external Na^+ concentration due to the mutation in genes that encode major Na^+ and Li^+ efflux system. ENA1-4 genes that encode p-type Na^+ pumps are being disrupted in the G19 leading to Na^+ accumulation [38, 39]. Besides the ENA pumps, NHA1 (an antiporter localized in the plasma membrane) and NHX1 (the antiporter that localized in tonoplast and sequester Na^+ in the vacuole) are disrupted in Axt3K clone [40, 41, 42]. Hence, Axt3K clone tends to accumulate more Na^+ than the G19 rendering it more salt sensitive than the G19. There were several positive clones of both G19 and Axt3k with enhanced tolerance to salt stress (Fig 3). Positive clones that showed growth complementation at high Na^+ concentrations were retransformed into the same yeast mutant background and rescreened in the higher Na^+ concentrations to confirm the growth complementation. This screening led to identify a consistent salt tolerant clone in Axt3k backgrounds. Sequencing of the cDNA inserts revealed that they encode for the putative proteins with sequence homologous to rice putative protein PROLM24, a prolamin precursor (BLASTN at MSU Osa1 Release 6.1).

C. Rice cDNA encoding for prolamin precursor PROLM24 confers salt tolerance in salt sensitive yeast mutants

Expression of cDNA clone of PROLM24 conferred tolerance to salt stress induced by NaCl in both Axt3K and G19 yeast mutants compared with the control strains transformed with the empty pYES2 vector.




Fig. 2 Screening yeast clone Axt3k (A and B) and G19 (C and D) transformed with rice cDNA (pYES+cDNA) and empty pYES vector for the salt stress tolerance. There were few positive clones of Axt3K appeared at 300 mM Na^+ (upper B) and several positive clones of G19 at 500 mM Na^+ (right D). Growth of the colonies carrying empty vector of both Axt3K and G19 had been severely affected at these Na^+ concentrations

However, expression of this cDNA did not affect yeast growth in absence of salt. Axt3k and G19 strains expressing the PROLM24 were able to grow upto 400 mM and 600 mM of NaCl respectively (Fig 3). These concentrations are well beyond the maximal salt concentration tolerated by wild-type yeast cells in SD medium (data not shown). Similarly, Axt3k mutant with PROLM24 expression showed higher growth rate in the medium with excess LiCl (50 mM) (Fig 4). Transport of Li^+ across the plasma membrane is mediated by the same transport system for Na^+ in yeast [18, 43]. However Li is more toxic than the Na^+ , hence cells survived only upto 50 mM LiCl.




Fig. 3 Complementation of yeast mutants by expression of PROLM24: Growth of the salt-sensitive yeast strain (Axt3K), Axt3K transformed with either the empty plasmid pYES (EV) or PROLM24 expression plasmid (PROLM). B: Growth of the salt-sensitive yeast strain (G19), G19 transformed with either the empty plasmid pYES (EV) or PROLM expression plasmid (PROLM). Drop assay of tenfold serial dilution of yeast cell suspension cultures in the SC-Uracil-Histidine with different NaCl concentrations. Growth was followed for 5 days

The observation that expression of PROLM24 rescued the salt sensitive phenotypes of G19 and Axt3k indicates the existence of a regulatory system that ameliorates the effect of salt stress in the transformed yeast mutants. As mentioned above, salt stress tolerance is dependent on the monovalent ion homeostasis which is maintained by Na^+ uptake and efflux systems in yeast which are represented by TRK1-2 and ENA1-4 genes [12, 13, 15, 18] mainly. In addition, the function of proteins CK-2, HAL3 and PPZ1 [31, 44] and the calcineurin [14] showed to have been involved in salt tolerance in yeast. Study with the expression of CK-2 revealed that the salt sensitivity of the ckb1 mutant (CK-2) is independent of Na^+ fluxes and hypothesized to be involved with protein dephosphorylation [44].

In absence of any evidence for probable function of PROLM24 in the yeast, a sequence similarity search of the cDNA insert (PROLM24) was carried out with BLAST tool against the yeast nucleotide and protein database (www.yeastgenome.org) to identify if there were any functionally similar proteins in yeast.




Fig. 4 Complementation of yeast mutants by expression of PROLM24. Growth of the salt-sensitive yeast strain (Axt3k) transformed with either the empty plasmid pYES (EV) or PROLM24 expression plasmid (PROLM) in LiCl. Drop assay of tenfold serial dilution of yeast cell suspension cultures in the SC-Uracil-Histidine with different LiCl concentrations. Growth was followed for 5 days

The result showed that PROLAM24 is partially homologous to yeast protein YJR049C, which encode for ATP-NAD kinases (EC 2.7.1.23) called UTR1 [45] (Fig. 5). Expression of cDNA sequence of heterologous protein PROLAM24 which has partial homology to nucleotide sequence of YJR049C may modify the expression of YJR049C and its protein UTR1 [46].


However, there is no evidence to establish any direct relationship between UTR1 and monovalent cation transport as it has no predictable transmembrane domains [47]. Nevertheless, UTR1 hardly have any effect on modulating the expression or activity of the ENA 1-4 or TRK transporters in G19 and Axt3K mutant backgrounds. Given that there is no significant difference in the intra-cellular Na^+ / K^+ ion ratio between wild type and transformed mutant cells (data not shown), probably the rice PROLM24 (Os06g31070) in yeast may have a function similar to that of Casein Kinase 2 which renders proteins insensitive to Na^+ and Li^+ toxicity [44].

D. Differential regulation of PROLM24 gene in rice

Prolamins are principal storage proteins of rice and so far a total of 34 gene copies located on chromosomes 3, 5, 6, 7, 11 and 12 have been identified in the rice cultivar Nipponbare [48]. Prolamin proteins are not located inside starch granules but primarily located on the exterior of starch granule. Prolamin protein makes a vitreous hydrophobic matrix by cross-linking with starch.

Blasting with NCBI transcript reference sequence database revealed that sequence of cDNA insert is homologous to Os06g31070 (Os06g0507200) of *Oryza sativa* L. [Japonica Group].

S. cerevisiae WU-BLAST2 Search



- [7] Serrano, R., and Villalba J. M. (1995). Expression and localization of plant membrane proteins in Saccharomyces. *Methods in Cell Biology*, 50, 481–496.
- [8] Ramos, J. (1999). Contrasting salt tolerance mechanisms in *Saccharomyces cerevisiae* and *Debaryomyces hansenii*. *Recent Research Development Microbiology*, 3, 377–390.
- [9] Greenway, H., & Munns R. (1980). Mechanisms of salt tolerance in non-halophytes. *Annual Review of Plant Physiology*, 31, 149–190.
- [10] Marschner, H. (1995). Mineral nutrition of higher plants. Springer, Berlin, Germany.
- [11] Serrano, R. 1996. Salt tolerance in plants and microorganisms: toxicity targets and defense responses. *Int. Rev. Cytol.* 165:1–51.
- [12] Gaber, R. F., Styles, C. A. & Fink, G. R. (1988) TRK1 encodes a plasma membrane protein required for high-affinity potassium transport in *Saccharomyces cerevisiae*. *Molecular and Cellular Biology*, 8, 2848–2859.
- [13] Ko, C. H. & Gaber R. F. (1991). TRK1 and TRK2 encode structurally related K⁺ transporters in *Saccharomyces cerevisiae*. *Molecular and Cellular Biology*, 11, 4266–4273.
- [14] Mendoza, I., Rubio, F., Rodríguez-Navarro, A. & Pardo, J. M. (1994). The protein phosphatase calcineurin is essential for NaCl tolerance of *Saccharomyces cerevisiae*. *Journal of Biological Chemistry*, 269:8792–8796.
- [15] Rudolph, H.K., Antebi, A., Fink, G.R., Buckley, C.M., Dorman, T.E., LeVitre, J., Davidow, L.S., Mao, J.I. & Moir, D.T. (1989). The yeast secretory pathway is perturbed by mutations in PMR1, a member of a Ca²⁺ ATPase family. *Cell*, 58(1), 133–145.
- [16] Martínez, R., Latreille, M.T. & Mirande, M. (1991). A PMR2 tandem repeat with a modified C-terminus is located downstream from the KRS1 gene encoding lysyl-tRNA synthetase in *Saccharomyces cerevisiae*. *Molecular and General Genetics*, 227(1), 149–154.
- [17] Rodríguez-Navarro, A., Quintero, F.J. & Garcíadeblás, B. (1994). Na⁺-ATPases and Na⁺/H⁺ antiporters in fungi. *Biochimica et Biophysica Acta*, 1187(2), 203–205.
- [18] Haro, R., Bañuelos, M. A., Quintero, F. J., Rubio, F. & Rodríguez-Navarro, A. (1993) Genetic basis of Sodium exclusion and Sodium tolerance in yeast. A model for plants. *Plant Physiology*, 89, 868–874.
- [19] Garcíadeblás, B., Rubio, F., Quintero, F.J., Bañuelos, M.A. & Haro, R. (1993). Differential expression of two genes encoding isoforms of the ATPase involved in sodium efflux in *Saccharomyces cerevisiae*. *Molecular & General Genetics*, 236, 363–368.
- [20] Andre, B. (1995). An overview of membrane transport proteins in *Saccharomyces cerevisiae*. *Yeast*, 11, 1575–1611.
- [21] Prior, C., Potier, S., Souciet J et al. Characterization of the NHA1 gene encoding a Na⁺/H⁺-antiporter of the yeast *Saccharomyces cerevisiae*. *FEBS Letter*, 1996; 387:89–93.
- [22] Nass, R. & Rao, R. (1998). Novel localization of a Na⁺/H⁺ exchanger in a late endosomal compartment of yeast. *Journal of Biological Chemistry*, 273: 21054–21060.
- [23] Mulet, J.M., Leube, M.P., Kron, S.J., Rios, G., Fink, G.R. & Serrano, R. (1999). A Novel Mechanism of Ion Homeostasis and Salt Tolerance in Yeast: the Hal4 and Hal5 Protein Kinases Modulate the Trk1-Trk2 Potassium Transporter. *Molecular and cellular biology*, 19(5), 3328–3337.
- [24] Pérez-Valle J., Jenkins H., Merchan S., Montiel V., Ramos J., Sharma S., Serrano R. & Yenush L. (2007). Key role for intracellular K⁺ and protein kinases Sat4/Hal4 and Hal5 in the plasma membrane stabilization of yeast nutrient transporters. *Molecular and Cellular Biology*, 27(16):5725–36.
- [25] Mulet, J.M., Alejandro, S., Romero, C., Serrano R., Munson, A.M., Haydon, D.H., Love, S.L., Fell, G.L., Palanivel, V.R. & Rosenwald, A.G. (2004). Yeast ARL1 encodes a regulator of K⁺ influx. *Journal of Cell Sciences*, 117(11):2309–20.
- [26] Casado, C., Yenush, L., Melero, C., Ruiz Mdel, C., Serrano, R., Pérez-Valle, J., Ariño, J. & Ramos J. (2010). Regulation of Trk-dependent potassium transport by the calcineurin pathway involves the Hal5 kinase. *FEBS Letters*, 584(11), 2415–2420.
- [27] Forment, J., Mulet, J.M., Vicente, O. & Serrano, R. (2002). The yeast SR protein kinase Sky1p modulates salt tolerance, membrane potential and the Trk1,2 potassium transporter. *Biochimica et Biophysica Acta*, 1565(1):36–40.
- [28] Rios, G., Ferrando, A. & Serrano, R. (1997). Mechanisms of salt tolerance conferred by overexpression of the HAL1 gene in *Saccharomyces cerevisiae*. *Yeast*, 13(6), 515–28.
- [29] Munson, A.M., Love, S.L., Shu, J., Palanivel, V.R. & Rosenwald, A.G. (2004). ARL1 participates with ATC1/LIC4 to regulate responses of yeast cells to ions. *Biochemical and Biophysical Research Communications*, 315(3), 617–623.
- [30] Ferrando, A., Kron, S. J., Rios, G., Fink, G. R. & Serrano, R. (1995). Regulation of cation transport in *Saccharomyces cerevisiae* by the salt tolerance gene HAL3. *Molecular and Cellular Biology*, 15, 5470–5481.
- [31] de Nadal, E., Clotet, J., Posas, F., Serrano, R., Gómez, N. & Ariño, J. (1998). The yeast halotolerance determinant Hal3p is an inhibitory subunit of the Ppz1p Ser/Thr protein phosphatase. *Proceedings of National Academy of Science, USA*, 95, 7357–7362.
- [32] Schachtman, D.P. & Schroeder, J.I. (1994). Structure and transport mechanism of a high-affinity potassium uptake transporter from higher plants. *Nature*, 370, 655–658.
- [33] Igarashi, Y., Yoshida, Y., Sanada, Y., Yamaguchi-Shinozaki, K., Wada, K. & Shinozaki, K. (1997) Characterization of the gene for $\Delta 1$ -pyrroline-5-carboxylate synthetase and correlation between the expression of the gene and salt tolerance in *Oryza sativa* L. *Plant molecular Biology*, 33(5), 857–865.
- [34] Obata, T., Kitamoto, H.K., Nakamura, A., Fukuda, A., Tanaka, Y., (2007) Rice Shaker Potassium Channel OsKAT1 Confers Tolerance to Salinity Stress on Yeast and Rice Cells. *Plant Physiology*, 144, 1978–1985.
- [35] Fukuda, A., Nakamura, A., Tagiri, A., Tanaka, H., Miyao, A., Hirochika, H. & Tanaka, Y. (2004) Function, intracellular Localization and the Importance in Salt Tolerance of a Vacuolar Na⁺/H⁺ Antiporter from Rice. *Plant Cell Physiology*, 45(2), 146–159.
- [36] Gietz, D., St Jean, A., Woods, R.A. & Schiestl, R.H. (1992) Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Research*, 20, 1425.
- [37] Chen, D., Yang, B., Kuo, T. (1992). One-step transformation of yeast in stationary phase. *Current Genetic* 21: 83–84.
- [38] Quintero, F. J., Garcíadeblás, B. & Rodríguez-Navarro, A. (1996). The SAL1 gene of *Arabidopsis*, encoding an enzyme with 3 (2), 5 bisphosphate nucleotidase and inositol 1-phosphatase activities, increases salt tolerance in yeast. *Plant Cell*, 8, 529–537.
- [39] Gobert, A., Park, G., Amtmann, A., Sanders, D. & Maathuis, F.J.M. (2006). *Arabidopsis thaliana* cyclic nucleotide gated channel 3 forms a nonselective ion transporter involved in germination and cation transport. *Journal of Experimental Botany*, 57: 791–800.
- [40] Wieland, J., Nitsche, A.M., Strayle, J., Steiner, H. & Rudolph, H.K. (1995). The PMR2 gene cluster encodes functionally distinct isoforms of a putative Na⁺ pump in the yeast plasma membrane. *EMBO J* 14, 3870–3882. O. Young, “Synthetic structure of industrial plastics (Book style with paper title and editor),” in *Plastics*, 2nd ed. vol. 3, J. Peters, Ed. New York: McGraw-Hill, 1964, pp. 15–64.
- [41] Bañuelos, M.A., Sychrova, H., Bleykasten-Grosshans, C., Souciet, J.L. & Potier, S. (1998). The Nha1 antiporter of *Saccharomyces cerevisiae* mediates sodium and potassium efflux. *Microbiology*, 144, 2749–2758.
- [42] Darley, C.P., Wuytswinkel, O.C.M., Woude, K., Mager, W.H. & De Boer, A.H. (2000). *Arabidopsis thaliana* and *Saccharomyces cerevisiae* NHX1 genes encode amiloride sensitive electroneutral Na⁺/H⁺ exchangers. *Biochemical Journal*, 351, 241–249.
- [43] Borst-Pauwels, G. W. F. H. (1981). Ion transport in yeast. *Biochimica et Biophysica Acta*, 650, 88–127.
- [44] de Nadal, E., Calero, F., Ramos, J. & Ariño, J. (1999) Biochemical and Genetic Analyses of the Role of Yeast Casein Kinase 2 in Salt Tolerance. *Journal of Bacteriology*, 181(20), 6456–6462.
- [45] Kawai, S., Suzuki, S., Mori, S. & Murata K. (2001). Molecular cloning and identification of UTR1 of a yeast *Saccharomyces cerevisiae* as a gene encoding an NAD kinase. *FEMS Microbiology Letters*, 200(2), 181–184.
- [46] Batard Y, Hehn A, Nedelkina S, Schalk M, Pallett K, Schaller H, Werck-Reichhart D (2000) Increasing expression of P450 and P450-reductase proteins from monocots in heterologous systems. *Arch Biochem Biophys* 379:161–169.
- [47] Tusnady, G.E. & Simon, I. (2001). The HMMTOP transmembrane topology prediction server. *Bioinformatics*, 17, 849–850.
- [48] Sundaram R. M., Sakthivel K., Hariprasad A. S., Ramesha M. S., Viraktamath B. C., Neeraja C. N., Balachandran S. M., Shobha Rani N., Revathi P. Sandhya P., et al. (2010) Molecular Breeding Development and validation of a PCR-based functional marker system for the major wide-compatible gene locus S5 in rice 26, 719–727.

- [49] Marchler-Bauer A et al. (2011), "CDD: a Conserved Domain Database for the functional annotation of proteins.", *Nucleic Acids Res.*39(D)225-9.
- [50] Hruz, T., Laule, O., Szabo, G., Wessendorp, F., Bleuler, S., Oertle, L., Widmayer, P., Gruissem, W. & Zimmermann, P (2008). Genevestigator V3: a reference expression database for the meta-analysis of transcriptomes. *Advances in Bioinformatics*, 2008, 420747.
- [51] Pandit, A., Rai, V., Bal, S., Kumar, V., Chauhan, M., Gautam, R.K., Singh, R., Sharma, P.C. & Singh, K., (2010). Combining QTL mapping and transcriptome profiling of bulked RILs for identification of functional polymorphism for salt tolerance genes in rice (*Oryza sativa* L.), *Molecular Genetics and Genomics*, 284(2), 121-136.
- [52] Walia, H., Wilson, C., Condamine, P., Liu, X., Ismail, A.M., Zeng, L., Wanamaker, S.I., Mandal, J., Xu, J., Cui, X. & Close T.J., (2005) Comparative transcriptional profiling of two contrasting rice genotypes under salinity stress during the vegetative growth stage. *Plant Physiology* 139:822–835.
- [53] Cotsaftis, O., Plett, D., Johnson, A.A., Walia, H., Wilson, C., Ismail, A.M., Close, T.J., Tester, M. & Baumann U. (2011) Root-specific transcript profiling of contrasting rice genotypes in response to salinity stress. *Molecular Plant*, 4(1),25-41.
- [54] Senadheera, P., Singh, R. K. & Maathuis, F.J. M., (2009). Differentially expressed membrane transporters in rice roots may contribute to cultivar dependent salt tolerance, *Journal of Experimental Botany*, 60(9): 2553–2563.
- [55] Charoenlappanit S, Roytrakul S , Teerakathiti T, Juntawong N (2010) Proteome analysis of salt tolerant and salt sensitive rice suspension cells in response to NaCl stress. 36th Congress on Science and Technology of Thailand.