

# The Existence and Uniqueness of Positive Solution for Nonlinear Fractional Differential Equation Boundary Value Problem

Chuanyun Gu, Shouming Zhong

**Abstract**—In this paper, the existence and uniqueness of positive solutions for nonlinear fractional differential equation boundary value problem is concerned by a fixed point theorem of a sum operator. Our results can not only guarantee the existence and uniqueness of positive solution, but also be applied to construct an iterative scheme for approximating it. Finally, the example is given to illustrate the main result.

**Keywords**—Fractional differential equation, Boundary value problem, Positive solution, Existence and uniqueness, Fixed point theorem of a sum operator.

## I. INTRODUCTION

**F**RACtIONAL differential equations are used in various fields, such as mechanics, physics, chemistry, engineering, economics and biological sciences, etc.; see [1–9] and the references therein. In recent years, there are many papers discuss the existence and multiplicity of positive solutions for nonlinear fractional differential equation boundary value problem by the use of Leray-Schauder theory, fixed-point theorems, etc., see [10–14]. However, there are few papers consider the existence of a unique positive solution for nonlinear fractional differential equation boundary value problem, see [15, 16].

In particular, by means of mixed monotone method, Xu, Jiang and Yuan [16] obtained the uniqueness of solution to the singular boundary value problem for fractional differential equation

$$\begin{cases} D_{0+}^{\alpha} u(t) = f(t, u(t)), & 0 < t < 1, \quad 3 < \alpha \leq 4, \\ u(0) = u(1) = u'(0) = u'(1) = 0. \end{cases} \quad (1)$$

They need

$$f(t, u) = q(t)[\varphi(u) + \psi(u)], \varphi : [0, +\infty) \rightarrow [0, +\infty)$$

is continuous and nondecreasing,  $\psi : (0, +\infty) \rightarrow (0, +\infty)$  is continuous and nondecreasing and  $q \in C((0, 1), (0, +\infty))$  satisfies

$$\int_0^1 s^{2-\eta(2-\alpha)}(1-s)^{\alpha-2-2\eta} q(s) ds < +\infty, \eta \in (0, 1).$$

Naturally, when  $f(t, u)$  can not be denoted by  $f(t, u) = q(t)[\varphi(u) + \psi(u)]$ , how to consider the unique positive solution of it? In this paper, by means of a fixed

Chuanyun Gu is with the School of Mathematics and Finance-Economics, Sichuan University of Arts and Science, Dazhou 635000, PR China.

Shouming Zhong is with Key Laboratory for NeuroInformation of Ministry of Education, University of Electronic Science and Technology of China, Chengdu 611731, PR China.

Email address: guchaunyun@163.com.

point theorem for a sum operator, we obtain the existence and uniqueness of positive solution to the nonsingular boundary value problem for fractional differential equation (1) with the assumption  $f(t, u) = g(t, u) + h(t, u)$ . Moreover, we can construct an iterative scheme to approximate the unique solution, which is important for evaluation and application.

## II. PRELIMINARIES AND PREVIOUS RESULTS

In this section, we present some definitions, lemmas and basic results that will be used in the proof of our main result.

**Definition 1** [3] The integral

$$I_{0+}^{\alpha} f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x \frac{f(t)}{(x-t)^{1-\alpha}} dt, \quad x > 0$$

is called the Riemann-Liouville fractional integral of order  $\alpha$ , where  $\alpha > 0$  and  $\Gamma(\alpha)$  denotes the gamma function.

**Definition 2** [3] For a function  $f(x)$  given in the interval  $[0, \infty)$ , the expression

$$D_{0+}^{\alpha} f(x) = \frac{1}{\Gamma(n-\alpha)} \left( \frac{d}{dx} \right)^n \int_0^x \frac{f(t)}{(x-t)^{\alpha-n+1}} dt,$$

is called the Riemann-Liouville fractional derivative of order  $\alpha$ , where  $n = [\alpha] + 1$ ,  $[\alpha]$  denotes the integer part of number  $\alpha$ .

**Lemma 1** [16] Given  $y \in C[0, 1]$  and  $3 < \alpha \leq 4$ , the unique solution of the fractional differential equation

$$\begin{cases} D_{0+}^{\alpha} u(t) = y(t), & 0 < t < 1, \\ u(0) = u(1) = u'(0) = u'(1) = 0. \end{cases} \quad (2)$$

is

$$u(t) = \int_0^1 G(t, s) y(s) ds, \quad t \in [0, 1]$$

where

$$G(t, s) = \begin{cases} \frac{(t-s)^{\alpha-1} + (1-s)^{\alpha-2} t^{\alpha-2} [(s-t) + (\alpha-2)(1-t)s]}{\Gamma(\alpha)}, & 0 \leq s \leq t \leq 1, \\ \frac{t^{\alpha-2} (1-s)^{\alpha-2} [(s-t) + (\alpha-2)(1-t)s]}{\Gamma(\alpha)}, & 0 \leq t \leq s \leq 1. \end{cases} \quad (3)$$

Here  $G(t, s)$  is called the Green function of the fractional differential equation (2).

**Lemma 2** <sup>[16]</sup> The Green function  $G(t, s)$  defined by (3) has the following property:

$$\frac{\alpha-2}{\Gamma(\alpha)}(1-s)^{\alpha-2}s^2t^{\alpha-2}(1-t)^2 \leq G(t, s) \leq \frac{1}{\Gamma(\alpha)}M_0t^{\alpha-2}(1-t)^2 \quad (4)$$

for  $t, s \in (0, 1)$ . where  $M_0 = \max\{\alpha - 1, (\alpha - 2)^2\}$ .

In the sequel, we present some basic concepts in ordered Banach spaces for completeness and a fixed point theorem which will be used later.

Suppose  $(E, \|\cdot\|)$  is a real Banach space which is partially ordered by a cone  $P \subset E$ , i.e.  $x \leq y$  if and only if  $y - x \in P$ . If  $x \leq y$  and  $x \neq y$ , then we denote  $x < y$ . We denote the zero element of  $E$  by  $\theta$ . Recall that a non-empty closed convex set  $P \subset E$  is a cone if it satisfies (i)  $x \in P, \lambda \geq 0 \Rightarrow \lambda x \in P$ ; (ii)  $x \in P, -x \in P \Rightarrow x = \theta$ .

Putting  $P^0 = \{x \in P \mid x \text{ is an interior point of } P\}$ , a cone  $P$  is said to be solid if  $P^0$  is non-empty. Moreover,  $P$  is called normal if there exists a constant  $N > 0$  such that, for all  $x, y \in E, \theta \leq x \leq y$  implies  $\|x\| \leq N\|y\|$ ; in this case  $N$  is called the normality constant of  $P$ . We say that an operator  $A : E \rightarrow E$  is increasing if  $x \leq y$  implies  $Ax \leq Ay$ .

For all  $x, y \in E$ , the notation  $x \sim y$  means that there exist  $\lambda > 0$  and  $\mu > 0$  such that  $\lambda x \leq y \leq \mu x$ . Clearly  $\sim$  is an equivalence relation. Given  $w > \theta$  (i.e.  $w \geq \theta$  and  $w \neq \theta$ ), we denote the set  $P_w = \{x \in E \mid x \sim w\}$  by  $P_w$ . It is easy to see that  $P_w \subset P$  for  $w \in P$ .

**Theorem 1** <sup>[17]</sup> Let  $P$  be a normal cone in a real Banach space  $E$ ,  $A : P \rightarrow P$  be an increasing  $\gamma$ -concave operator and  $B : P \rightarrow P$  be an increasing sub-homogeneous operator. Assume that

- (i) there is  $w > \theta$  such that  $Aw \in P_w$  and  $Bw \in P_w$ ;
- (ii) there exists a constant  $\delta_0 > 0$  such that  $Ax \geq \delta_0 Bx, \forall x \in P$ .

Then operator equation  $Ax + Bx = x$  has a unique solution  $x^*$  in  $P_w$ . Moreover, constructing successively the sequence  $y_n = Ay_{n-1} + By_{n-1}, n = 1, 2, \dots$  for any initial value  $y_0 \in P_w$ , we have  $y_n \rightarrow x^*$  as  $n \rightarrow \infty$ .

### III. MAIN RESULTS

In this section, we apply Theorem 1 to investigate the fractional differential equation (1) with the assumption  $f(t, u) = g(t, u) + h(t, u)$ , and we obtain the new result on the existence and uniqueness of positive solution.

In this paper, we will work in the Banach space  $C[0, 1] = \{x : [0, 1] \rightarrow R \text{ is continuous}\}$  with the standard norm  $\|x\| = \sup\{|x(t)| : t \in [0, 1]\}$ . Notice that this space can be endowed with a partial order given by  $x, y \in C[0, 1], x \leq y \Leftrightarrow x(t) \leq y(t) \text{ for } t \in [0, 1]$ .

Let  $P = \{x \in C[0, 1] \mid x(t) \geq 0, t \in [0, 1]\}$  be the standard cone. Evidently,  $P$  is a normal cone in  $C[0, 1]$  and the normality constant is 1.

**Theorem 2** Assume that

- (H1)  $g, h : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$  are continuous and increasing with respect to the second argument,  $h(t, 0) \neq 0$ ;
- (H2) there exists a constant  $\gamma \in (0, 1)$  such that  $g(t, \lambda x) \geq \lambda^\gamma g(t, x), \forall t \in [0, 1], \lambda \in (0, 1), x \in [0, \infty)$ , and  $h(t, \mu x) \geq \mu h(t, x)$  for  $\mu \in (0, 1), t \in [0, 1], x \in [0, \infty)$ ;

(H3) there exists a constant  $\delta_0 > 0$  such that  $g(t, x) \geq \delta_0 h(t, x), t \in [0, 1], x \geq 0$ . Then the fractional differential equation (1) has a unique positive solution  $u^*$  in  $P_w$ , where  $w(t) = t^{\alpha-2}(1-t)^2, t \in [0, 1]$ . Moreover, for any initial value  $u_0 \in P_w$ , constructing successively the iterative scheme

$$u_{n+1}(t) = \int_0^1 G(t, s)f(s, u_n(s))ds, \quad n = 0, 1, 2, \dots,$$

we have  $u_n(t) \rightarrow u^*(t)$  as  $n \rightarrow \infty$ , where  $G(t, s)$  is given as (3).

**Proof:** To begin with, from Lemma 1, the fractional differential equation (1) has an integral formulation given by

$$\begin{aligned} u(t) &= \int_0^1 G(t, s)f(s, u(s))ds \\ &= \int_0^1 G(t, s)[g(s, u(s)) + h(s, u(s))]ds \end{aligned}$$

where  $G(t, s)$  is given as in Lemma 1.

Define two operators  $A : P \rightarrow E$  and  $B : P \rightarrow E$  by

$$\begin{aligned} Au(t) &= \int_0^1 G(t, s)g(s, u(s))ds, \\ Bu(t) &= \int_0^1 G(t, s)h(s, u(s))ds. \end{aligned}$$

It is easy to prove that  $u$  is the solution of the fractional differential equation (1) if and only if  $u = Au + Bu$ . By assumption (H1) and Lemma 2, we know that  $A : P \rightarrow P$  and  $B : P \rightarrow P$ . In the sequel we check that  $A, B$  satisfy all assumptions of Theorem 1.

Firstly, we prove that are two increasing operators.

In fact, from assumption (H1) and Lemma 2, for  $u, v \in P$  with  $u \geq v$ , we know that  $u(t) \geq v(t), t \in [0, 1]$  and obtain

$$\begin{aligned} Au(t) &= \int_0^1 G(t, s)g(s, u(s))ds \\ &\geq \int_0^1 G(t, s)g(s, v(s))ds \\ &= Av(t) \end{aligned}$$

That is  $Au \geq Av$ . Similarly,  $Bu \geq Bv$ .

Next we show that  $A$  is a  $\gamma$ -concave operator and  $B$  is a sub-homogeneous operator.

In fact, for any  $\lambda \in (0, 1)$  and  $u \in P$ , from (H2) we know that

$$\begin{aligned} A(\lambda u)(t) &= \int_0^1 G(t, s)g(s, \lambda u(s))ds \\ &\geq \lambda^\gamma \int_0^1 G(t, s)g(s, u(s))ds \\ &= \lambda^\gamma Au(t) \end{aligned}$$

That is,  $A(\lambda u) \geq \lambda^\gamma Au$  for  $\lambda \in (0, 1), u \in P$ . So the operator  $A$  is a  $\gamma$ -concave operator. Also, for any  $\mu \in (0, 1)$  and  $u \in P$ , by (H2) we obtain

$$\begin{aligned} B(\mu u)(t) &= \int_0^1 G(t, s)h(s, \mu u(s))ds \\ &\geq \mu \int_0^1 G(t, s)h(s, u(s))ds \\ &= \mu Bu(t) \end{aligned}$$

That is,  $B(\mu u) \geq \mu Bu$  for  $\mu \in (0, 1), u \in P$ . So the operator  $B$  is a sub-homogeneous operator.

Now we show that  $Aw \in P_w$  and  $Bw \in P_w$ , where  $w(t) = t^{\alpha-2}(1-t)^2$ .

By (H1) and Lemma 2,

$$\begin{aligned} \frac{\alpha-2}{\Gamma(\alpha)}w(t) \int_0^1 s^2(1-s)^{\alpha-2}g(s, 0)ds &\leq Aw(t) \\ &= \int_0^1 G(t, s)g(s, w(s))ds \\ &\leq \frac{1}{\Gamma(\alpha)}w(t) \int_0^1 M_0g(s, 1)ds \end{aligned}$$

From (H1) and (H3), we have

$$g(s, 1) \geq g(s, 0) \geq \delta_0 h(s, 0) \geq 0$$

Since  $h(t, 0) \neq 0$ , we can get

$$\int_0^1 g(s, 1)ds \geq \int_0^1 g(s, 0)ds \geq \delta_0 \int_0^1 h(s, 0)ds > 0,$$

and in consequence,

$$l_1 := \frac{\alpha-2}{\Gamma(\alpha)} \int_0^1 s^2(1-s)^{\alpha-2} g(s, 0)ds > 0,$$

$$l_2 := \frac{1}{\Gamma(\alpha)} \int_0^1 M_0 g(s, 1)ds > 0.$$

So  $l_1 w(t) \leq Aw(t) \leq l_2 w(t), t \in [0, 1]$ ; and hence we have  $Aw \in P_w$ . Similarly,

$$\frac{\alpha-2}{\Gamma(\alpha)} w(t) \int_0^1 s^2(1-s)^{\alpha-2} h(s, 0)ds \leq Bw(t)$$

$$\leq \frac{1}{\Gamma(\alpha)} w(t) \int_0^1 M_0 h(s, 1)ds$$

from  $h(t, 0) \neq 0$ , we easily prove  $Bw \in P_w$ . Hence the condition (i) of Theorem 1 is satisfied.

In the following we show that the condition (ii) of Theorem 1 is satisfied.

For  $u \in P$ , by (H3),

$$Au(t) = \int_0^1 G(t, s)g(s, u(s))ds$$

$$\geq \delta_0 \int_0^1 G(t, s)h(s, u(s))ds$$

$$= \delta_0 Bu(t)$$

Then we get  $Au \geq \delta_0 Bu, u \in P$ .

Finally, by means of Theorem 1, the operator equation  $Au + Bu = u$  has a unique positive solution  $u^*$  in  $P_w$ . Moreover, constructing successively the iterative scheme

$$u_n = Au_{n-1} + Bu_{n-1}, n = 1, 2, \dots$$

for any initial value  $u_0 \in P_w$ , we have  $u_n \rightarrow u^*$  as  $n \rightarrow \infty$ . That is, the fractional differential equation (1) has a unique positive solution  $u^*$  in  $P_w$ . For any initial value  $u_0 \in P_w$ , constructing successively the iterative scheme

$$u_{n+1}(t) = \int_0^1 G(t, s)f(s, u_n(s))ds, \quad n = 0, 1, 2, \dots,$$

we have  $u_n \rightarrow u^*$  as  $n \rightarrow \infty$ .

**Corollary 1** When  $f(t, u) = q(t)[\varphi(u) + \psi(u)]$  satisfy the conditions of theorem 2, the nonsingular boundary value problem for fractional differential equation (1) has a unique positive solution  $u^*$  in  $P_w$ , where  $w(t) = t^{\alpha-2}(1-t)^2, t \in [0, 1]$ . Moreover, for any initial value  $u_0 \in P_w$ , constructing successively the iterative scheme

$$u_{n+1}(t) = \int_0^1 G(t, s)f(s, u_n(s))ds, \quad n = 0, 1, 2, \dots,$$

we have  $u_n(t) \rightarrow u^*(t)$  as  $n \rightarrow \infty$ , where  $G(t, s)$  is given as (3).

**Remark 1** By Theorem 2, Corollary 1 is obvious. Moreover, the unique positive solution  $u^*$  we obtain satisfies:

(i) there exist  $\lambda > \mu > 0$  such that  $\mu t^{\alpha-2}(1-t)^2 \leq u^* \leq \lambda t^{\alpha-2}(1-t)^2, t \in [0, 1]$ ,

(ii) we can take any initial value in  $P_w$  and then construct an iterative scheme which can approximate the unique solution.

#### IV. EXAMPLE

We present one example to illustrate Theorem 2.

**Example 1** Consider the following fractional differential equation:

$$\begin{cases} D_{0+}^{\frac{10}{3}} u(t) = u^{\frac{1}{5}}(t) + \arctan u(t) + t^3 + t^2 + \frac{\pi}{2}, \\ u(0) = u(1) = u'(0) = u'(1) = 0. \end{cases} \quad (5)$$

In this example, we have  $\alpha = \frac{10}{3}$ . Let

$$g(t, u) = u^{\frac{1}{5}}(t) + t^2 + \frac{\pi}{2},$$

$$h(t, u) = \arctan u(t) + t^3,$$

$$\gamma = \frac{1}{5}.$$

Obviously,  $g, h : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$  are continuous and increasing with respect to the second argument,  $h(t, 0) = t^3 \neq 0$ . Besides, for  $t \in [0, 1], \lambda \in (0, 1), x \in [0, \infty)$ , we have

$$g(t, \lambda u) = \lambda^{\frac{1}{5}} u^{\frac{1}{5}}(t) + t^2 + \frac{\pi}{2}$$

$$\geq \lambda^{\frac{1}{5}} u^{\frac{1}{5}}(t) + \lambda^{\frac{1}{5}} (t^2 + \frac{\pi}{2})$$

$$= \lambda^{\frac{1}{5}} (u^{\frac{1}{5}}(t) + t^2 + \frac{\pi}{2})$$

$$= \lambda^{\gamma} g(t, u)$$

and for  $t \in [0, 1], \mu \in (0, 1), x \in [0, \infty)$ , we have  $\arctan(\mu u) \geq \mu \arctan u$ , and thus  $h(t, \mu u) \geq \mu h(t, u)$ .

Moreover, if we take  $\delta_0 \in (0, 1]$ , then we obtain

$$g(t, u) = u^{\frac{1}{5}}(t) + t^2 + \frac{\pi}{2}$$

$$\geq t^2 + \frac{\pi}{2}$$

$$\geq t^3 + \arctan u$$

$$\geq \delta_0 (t^3 + \arctan u)$$

$$= \delta_0 h(t, u)$$

Hence all the conditions of Theorem 2 are satisfied. An application of Theorem 2 implies that the fractional differential equation (5) has a unique positive solution in  $P_w$ , where  $w(t) = t^{\frac{10}{3}}(1-t)^2, t \in [0, 1]$ .

**Remark 2** The nonlinearity  $f$  in example 1 can not be denoted by  $f(t, u) = q(t)[\varphi(u) + \psi(u)]$ , so the positive solution of example 1 can not be obtained by virtue of [16]. Theorem 2 generalizes the results in [16].

#### ACKNOWLEDGMENT

This work was supported by the National Basic Research Program of China (2010CB732501) and the Program for New Century Excellent Talents in University (NCET-10-0097).

#### REFERENCES

- [1] KB. Oldham, J. Spanier, *The fractional calculus*, Williams and Wilkins, New York: Academic Press, 1974.
- [2] KS. Miller, B. Ross, *An introduction to the fractional calculus and fractional differential equations*, John Wiley, New York, 1993.
- [3] SG. Samko, AA. Kilbas, OI Marichev, *Fractional integral and derivatives: theory and applications*, Gordon and Breach, Switzerland, 1993.
- [4] I. Podlubny, *Fractional differential equations, mathematics in science and engineering*, New York: Academic Press, 1999.
- [5] AA. Kilbas, HM. Srivastava, JJ. Trujillo, *Theory and applications of fractional differential equations*, North-Holland mathematics studies, 2006, 204.
- [6] WG. Glockle, TF. Nonnenmacher, A fractional calculus approach of self-similar protein dynamics, *Biophys J*, 1995, 68, 46-53.

- [7] F. Mainardi, Fractional calculus: some basic problems in continuum and statistical mechanics, *Fractals and fractional calculus in continuum mechanics*, 1997,291-348.
- [8] K. Diethelm, AD. Freed, On the solutions of nonlinear fractional order differential equations used in the modelling of viscoplasticity, *Scientific computing in chemical engineering II C computational fluid dynamics, reaction engineering and molecular properties*, 1999,217-224.
- [9] EM. Rabei, KI. Nawaeh, RS. Hijawi, SI. Muslih, D. Baleanu, The Hamilton formalism with fractional derivatives, *J Math Anal Appl*, 2007, 327,891-897.
- [10] Y. Zhao, S. Sun, Z. Han, The existence of multiple positive solutions for boundary value problems of nonlinear fractional differential equations, *Commun Nonlinear Sci Numer Simulat*, 2011, 16,2086-2097.
- [11] D. Jiang, C. Yuan, The positive properties of the Green function for Dirichlet-type boundary value problems of nonlinear fractional differential equations and its application, *Nonlinear Analysis*, 2010, 72,710-719.
- [12] S. Liang, J. Zhang, Existence and uniqueness of strictly nondecreasing and positive solution for a fractional three-point boundary value problem, *Comput Math Appl*, 2011, 62,1333-1340.
- [13] X. Yang, Z. Wei, W. Dong, Existence of positive solutions for the boundary value problem of nonlinear fractional differential equations, *Commun Nonlinear Sci Numer Simulat*, 2012, 17,85-92.
- [14] X. Ding, Y. Feng, R. Bu, Existence, nonexistence and multiplicity of positive solutions for nonlinear fractional differential equations, *J Appl Math Comput*, 2012, 40,371-381.
- [15] C. Zhai, W. Yan, C. Yang, A sum operator method for the existence and uniqueness of positive solutions to Riemann-Liouville fractional differential equation boundary value problems, *Commun Nonlinear Sci Numer Simulat*, 2013, 18,858-866.
- [16] X. Xu, D. Jiang, C. Yuan, Multiple positive solutions for the boundary value problem of a nonlinear fractional differential equation, *Nonlinear Analysis*, 2009, 71,4676-4688.
- [17] C. Zhai, DR. Anderson, A sum operator equation and applications to nonlinear elastic beam equations and Lane-Emden-Fowler equations, *J. Math. Anal. Appl*, 2011, 375,388-400.

**Chuanyun Gu** was born in Henan Province, China, in 1982. he received the B.S. degree from Nanyang Normal University, Nanyang, in 2008. he received the M.S. degree in applied mathematics from Xihua University, Chengdu, in 2012. His research interests include differential equations and functional analysis.

**Shouming Zhong** was born in 1955 in Sichuan, China. He received B.S. degree in applied mathematics from UESTC, Chengdu, China, in 1982. From 1984 to 1986, he studied at the Department of Mathematics in Sun Yat-sen University, Guangzhou, China. From 2005 to 2006, he was a visiting research associate with the Department of Mathematics in University of Waterloo, Waterloo, Canada. He is currently as a full professor with School of Applied Mathematics, UESTC. His current research interests include differential equations, neural networks, biomathematics and robust control. He has authored more than 80 papers in reputed journals such as the International Journal of Systems Science, Applied Mathematics and Computation, Chaos, Solitons and Fractals, Dynamics of Continuous, Discrete and Impulsive Systems, Acta Automatica Sinica, Journal of Control Theory and Applications, Acta Electronica Sinica, Control and Decision, and Journal of Engineering Mathematics.