

The symmetric solutions for boundary value problems of second-order singular differential equation

Li Xiguang

Abstract—In this paper, by constructing a special operator and using fixed point index theorem of cone, we get the sufficient conditions for symmetric positive solution of a class of nonlinear singular boundary value problems with p-Laplace operator, which improved and generalized the result of related paper.

Keywords—Banach space, cone, fixed point index, singular differential equation, p-Laplace operator, symmetric solutions.

I. INTRODUCTION

THE boundary value problems with p-Laplace operator arises in a variety of applied mathematics and physics, and they are widely applied in studying for non-newtonian fluid mechanics, cosmological physics, plasma physics, and theory of elasticity, etc. In recent years, some important results have been obtained by a variety of method(see[1-4]). On the other hand, the study for the symmetric and multiple solutions to this problem is more and more active (see[5-6]). In paper [5], Sun study for the problem

$$\begin{cases} (u)'' + a(t)f(t, u(t)) = 0, t \in (0, 1) \\ u(0) = \alpha u(\eta) = u(1), \end{cases}$$

where $\alpha \in (0, 1), \eta \in (0, \frac{1}{2}]$, by using spectrum theory, Sun get the existence of symmetric and multiple solution. But when $p \neq 2$, $\phi_p(u)$ is nonlinear, so the method of the paper [5] is not suitable to p-laplace operator. In paper [6], Tian and Liu study for the problem

$$\begin{cases} (\phi_p(u'))' + a(t)f(t, u(t)) = 0, t \in (0, 1) \\ u(0) = \alpha u(\eta) = u(1), \end{cases}$$

where $\phi(s)$ is p-Laplace operator. Motivated by paper [5,6], we consider the existence of solution for the following problems:

$$\begin{cases} (\phi_p(u'))' + h_1(t)f(u, v) = 0, \\ (\phi_p(v'))' + h_2(t)g(u) = 0, \\ u(0) = \gamma u(\eta) = u(1), \\ v(0) = \gamma v(\eta) = v(1), \end{cases} \quad (1)$$

where $t \in (0, 1), \gamma \in (0, 1), \eta \in (0, \frac{1}{2}]$, $\phi(s)$ is a p-Laplace operator, i.e. $\phi_p(s) = |s|^{p-2}s, p > 1$. Obviously, if $\frac{1}{p} + \frac{1}{q} = 1$, then $(\phi_p)^{-1} = \phi_q$.

Compare with above paper, our method is different. By constructing a new operator, and using fixed point index theorem, we get the sufficient condition of the existence of

symmetric solution, which improved and generalized the result of paper [5,6,7].

In this paper, we always suppose that the following conditions hold:

(H₁) $f \in C([0, +\infty) \times [0, +\infty), [0, +\infty)), g \in C([0, +\infty), [0, +\infty))$.

(H₂) $h_i \in C((0, 1), [0, +\infty)), h_i(t) = h_i(1-t), t \in (0, 1)$, for any subinterval of $(0, 1)$, $h_i(t) \not\equiv 0$, and $\int_0^1 h_i(t)dt < +\infty (i = 1, 2)$.

(H₃) There exists $\alpha \in (0, 1]$, such that $\liminf_{u \rightarrow +\infty} \frac{g(u)}{u^{\frac{p-1}{\alpha}}} = +\infty$ and $\liminf_{v \rightarrow +\infty} \frac{f(u, v)}{v^{(p-1)\alpha}} > 0$ hold uniformly to $u \in R^+$.

(H₄) There exists $\beta \in (0, +\infty)$, such that $\limsup_{u \rightarrow 0^+} \frac{g(u)}{u^{\frac{p-1}{\beta}}} = 0$ and $\limsup_{v \rightarrow 0^+} \frac{f(u, v)}{v^{(p-1)\beta}} < +\infty$ hold uniformly to $u \in R^+$.

(H₅) There exists $n \in (0, 1]$, such that $\liminf_{u \rightarrow 0^+} \frac{g(u)}{u^{\frac{p-1}{n}}} = +\infty$ and $\liminf_{v \rightarrow 0^+} \frac{f(u, v)}{v^{(p-1)n}} > 0$ hold uniformly to $u \in R^+$.

(H₆) $f(u, v)$ and $g(u)$ are nondecreasing with respect to u and v , and there exists $R > 0$, such that $\frac{\gamma}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q(k_1(s))ds f(R, \frac{\gamma}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q(k_1(s))ds \times g(R)) < R$, where $k_i(s) = \int_s^{\frac{1}{2}} h_i(\tau)d\tau, i = 1, 2$.

For convenience, we list the following definitions and lemmas:

Definition 1.1 If $u(t) = u(1-t), t \in [0, 1]$, we call $u(t)$ is symmetric in $[0, 1]$.

Definition 1.2 If (u, v) is a positive solution of problem (1), and u, v is symmetric in $[0, 1]$, we call (u, v) is symmetric positive solution of problem (1).

Definition 1.3 If $u(\lambda t_1 + (1-\lambda)t_2) \geq \lambda u(t_1) + (1-\lambda)u(t_2)$, we call $u(t)$ is concave in $[0, 1]$.

Let $E = C[0, 1]$, define the norm $\|u\| = \max_{t \in [0, 1]} |u(t)|$, obviously $(E, \|\cdot\|)$ is a Banach space.

Let $K = \{u \in E | u(t) > 0, u(t) \text{ is a symmetric concave function, } t \in [0, 1]\}$, then K is a cone in E . By $(H_1), (H_2)$, the solution of problem (1) is equivalent to the solution of system of equation (2).

Li Xiguang is with the Department of Mathematics, Qingdao University of Science and Technology, Qingdao, 266061, China. e-mail: lxxg0417@tom.com.

$$\left. \begin{aligned}
 u(t) &= \begin{cases} \int_0^t \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds + \\ \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds, \\ 0 \leq t \leq \frac{1}{2}, \\ \int_t^1 \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds + \\ \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds, \\ \frac{1}{2} \leq t \leq 1, \end{cases} \\
 v(t) &= \begin{cases} \int_0^t \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds + \\ \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds, \\ 0 \leq t \leq \frac{1}{2}, \\ \int_t^1 \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds + \\ \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds, \\ \frac{1}{2} \leq t \leq 1. \end{cases}
 \end{aligned} \right\}$$

We define $T : K \rightarrow E$:

$$(Tu)(t) = \begin{cases} \int_0^t \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds + \\ \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds, \\ 0 \leq t \leq \frac{1}{2}, \\ \int_t^1 \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds + \\ \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds, \\ \frac{1}{2} \leq t \leq 1, \end{cases} \tag{3}$$

where

$$v(t) = \begin{cases} \int_0^t \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds + \\ \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds, 0 \leq t \leq \frac{1}{2}, \\ \int_t^1 \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds + \\ \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds, \frac{1}{2} \leq t \leq 1. \end{cases} \tag{4}$$

Obviously $Tu \in E$, it is easy to show if T has fixed point u , then by (4), problem (1) has a solution (u, v) .

Lemma 1.1 Let $(H_1), (H_2)$, then $T : K \rightarrow K$ is completely continuous.

Proof $\forall u \in K$, by $(H_1), (H_2)$, we can get $(Tu)(t) \geq 0, t \in [0, 1]$.

$$v'(t) = \begin{cases} \phi_q(\int_t^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau), 0 \leq t \leq \frac{1}{2}, \\ -\phi_q(\int_t^1 h_2(\tau) g(u(\tau)) d\tau), \frac{1}{2} \leq t \leq 1, \end{cases}$$

correspondingly $(\phi_p(v'))' = -h_2(t)g(u) \leq 0, 0 < t < 1$, so v is concave in $[0, 1]$.

Next we show v is symmetric in $[0, 1]$.

When $t \in [0, \frac{1}{2}], 1-t \in [\frac{1}{2}, 1]$, so

$$\begin{aligned}
 v(1-t) &= \int_{1-t}^1 \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds + \\ &\frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds \\ &= \int_0^t \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds + \\ &\frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds \\ &= v(t).
 \end{aligned}$$

Similarly, we have $v(1-t) = v(t), t \in [\frac{1}{2}, 1]$. So v is a symmetric concave function in $[0, 1]$.

$$(Tu)'(t) = \begin{cases} \phi_q(\int_t^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau), 0 \leq t \leq \frac{1}{2}, \\ -\phi_q(\int_t^1 h_1(\tau) f(u(\tau), v(\tau)) d\tau), \frac{1}{2} \leq t \leq 1, \end{cases}$$

(2) so $(\phi_p((Tu)'))' = -h_1(t)f(u, v) \leq 0, 0 < t < 1$, i.e. Tu is concave in $[0, 1]$.

Next we show Tu is symmetric in $[0, 1]$. when $t \in [0, \frac{1}{2}], 1-t \in [\frac{1}{2}, 1]$, so

$$\begin{aligned}
 (Tu)(1-t) &= \int_{1-t}^1 \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds + \\ &\frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds \\ &= \int_0^t \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds + \\ &\frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds \\ &= (Tu)(t).
 \end{aligned}$$

Similarly, we have $(Tu)(1-t) = (Tu)(t), t \in [\frac{1}{2}, 1]$. so Tu is concave in $[0, 1]$, so $TK \subset K$. On the other hand, let D is a arbitrary bounded set of K , then there exist constant $c > 0$, such that $D \subset \{u \in K \mid \|u\| \leq c\}$. Let $b = \max_{u \in [0, c]} g(u)$, so

$\forall u \in D$, we have

$$\begin{aligned}
 \|v\| &= \left| \int_0^{\frac{1}{2}} \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds + \right. \\ &\left. \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) g(u(\tau)) d\tau) ds \right| \\ &\leq \frac{b^{q-1}}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q(\int_s^{\frac{1}{2}} h_2(\tau) d\tau) ds = a.
 \end{aligned}$$

Let $L = \max_{u \in [0, c], v \in [0, a]} f(u, v)$, so $\forall u \in D$, we have

$$\begin{aligned}
 \|Tu\| &= \left| \int_0^{\frac{1}{2}} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds \right. \\ &\left. + \frac{\gamma}{1-\gamma} \int_0^\eta \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) f(u(\tau), v(\tau)) d\tau) ds, \right| \\ &\leq \frac{L^{q-1}}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau) d\tau) ds.
 \end{aligned}$$

$$\begin{aligned} \|(Tu)'\| &= \max\{|\phi_q(\int_0^{\frac{1}{2}} h_1(\tau)f(u(\tau), v(\tau))d\tau)|, \\ &|\phi_q(\int_{\frac{1}{2}}^1 h_1(\tau)f(u(\tau), v(\tau))d\tau)|\} \\ &\leq L^{q-1}\phi_q(\int_0^{\frac{1}{2}} h_1(\tau)d\tau). \end{aligned}$$

By Arzela-Ascoli theorem, we know TD is compact set. By Lebesgue dominated convergence theorem, it is easy to show T is continuous in K , so $T : K \rightarrow K$ is completely continuous.

Lemma 1.2 For any $0 < \epsilon < \frac{1}{2}, u \in K$, we have

- (1) $u(t) \geq \|u\|t(1-t), \forall t \in [0, 1];$
- (2) $u(t) \geq \epsilon^2\|u\|, t \in [\epsilon, 1-\epsilon].$ (the proof is elementary, we omit it.)

Lemma 1.3(see [8]) Let K is a cone of E in Banach space, Ω_1 and Ω_2 are open subsets in $E, \theta \in \Omega_1, \overline{\Omega_1} \subset \Omega_2$, and $T : K \cap (\overline{\Omega_2} \setminus \Omega_1) \rightarrow K$ is a completely continuous operator, and satisfy one of the following conditions:

(1) $\|Tx\| \leq \|x\|, \forall x \in K \cap \partial\Omega_1, \|Tx\| \geq x, \forall x \in K \cap \partial\Omega_2,$

(2) $\|Tx\| \geq \|x\|, \forall x \in K \cap \partial\Omega_1, \|Tx\| \leq x, \forall x \in K \cap \partial\Omega_2,$

then A has at least one fixed point in $K \cap (\Omega_2 \setminus \Omega_1).$

Lemma 1.4(see [9]) Let K is a cone of E in Banach space, $K_r = \{x \in K \mid \|x\| \leq r\}$, suppose $A : K_r \rightarrow K$ is a completely continuous, and satisfy $Tx \neq x, \forall x \in \partial K_r,$

- (1) If $\|Tx\| \leq x, \forall x \in \partial K_r,$ then $i(T, K_r, K) = 1,$
- (2) If $\|Tx\| \geq x, \forall x \in \partial K_r,$ then $i(T, K_r, K) = 0.$

II. CONCLUSION

Theorem 2.1 Suppose $(H_1) - (H_4)$ hold, then problem (1) has at least one positive solution.

Proof By (H_3) , there exist ν and a sufficient large number $M > 0$, such that

$$f(u, v) \geq \nu^{p-1}v^{(p-1)\alpha}, \forall u \in R^+, v > M, \tag{5}$$

$$g(u) \geq C_0^{p-1}u^{\frac{p-1}{\alpha}}, \forall u > M, \tag{6}$$

where $C_0 = \max\{(\frac{\gamma}{1-\gamma} \int_{\epsilon}^{\eta} \phi_q(k_2(s))ds)^{-1}, (\frac{2}{\nu\gamma^{\alpha}\epsilon^2(\frac{1}{1-\gamma} \int_0^{\eta} \phi_q(k_1(s))^{\alpha+1}})^{\frac{1}{\alpha}}\}$. Let $N = (M + 1)\epsilon^{-2},$ if $u \in K \cap \partial K_N,$ by Lemma 2, $\min_{\epsilon \leq t \leq 1-\epsilon} u(t) \geq \epsilon^2\|u\| = \epsilon^2N = M + 1,$ by (3)-(6) and the symmetric property, for any $t \in [\epsilon, 1-\epsilon]$

$$\begin{aligned} v(t) &= \int_0^t \phi_q(\int_{\frac{1}{2}}^s h_2(\tau)g(u(\tau))d\tau)ds + \\ &\frac{\gamma}{1-\gamma} \int_0^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_2(\tau)g(u(\tau))d\tau)ds \\ &\geq \frac{\gamma}{1-\gamma} \int_0^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_2(\tau)g(u(\tau))d\tau)ds \\ &\geq \frac{\gamma}{1-\gamma} \int_{\epsilon}^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_2(\tau)g(u(\tau))d\tau)ds \\ &\geq \frac{C_0\gamma}{1-\gamma} \int_{\epsilon}^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_2(\tau)(u(\tau)^{\frac{p-1}{\alpha}})d\tau)ds \\ &\geq \frac{C_0\gamma}{1-\gamma} \int_{\epsilon}^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_2(\tau))d\tau)ds(\epsilon^2\|u\|)^{\frac{1}{\alpha}} \\ &\geq \frac{C_0\gamma}{1-\gamma} \int_{\epsilon}^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_2(\tau))d\tau)ds(M + 1)^{\frac{1}{\alpha}} \\ &\geq M + 1. \end{aligned}$$

$$\begin{aligned} \|Tu\| &= |\int_0^{\frac{1}{2}} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau)f(u(\tau), v(\tau))d\tau)ds + \\ &\frac{\gamma}{1-\gamma} \int_0^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau)f(u(\tau), v(\tau))d\tau)ds, | \\ &\geq \frac{1}{1-\gamma} \int_0^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau)f(u(\tau), v(\tau))d\tau)ds, \\ &\geq \frac{\nu}{1-\gamma} \int_0^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau)v(\tau)^{(p-1)\alpha}d\tau)ds, \\ &\geq \frac{\nu}{1-\gamma} \int_{\epsilon}^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau)d\tau)ds \times \\ &(\frac{C_0\gamma}{1-\gamma} \int_{\epsilon}^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau)d\tau)ds)^{\alpha} \epsilon^2\|u\| \\ &= \nu C_0^{\alpha}\gamma^{\alpha}\epsilon^2(\frac{1}{1-\gamma} \int_0^{\eta} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau)d\tau)ds)^{\alpha+1}\|u\| \\ &\geq 2\|u\|, \end{aligned}$$

so $\|Tu\| > \|u\|, \forall u \in K \cap K_N,$ by lemma 1.4, we can get

$$i(T, K \cap K_N, K) = 0. \tag{7}$$

On the other hand, by the second limit of H_4 , there exists a sufficient small number $r_1 \in (0, 1)$ such that

$$C_1^{p-1} = \sup\{\frac{f(u, v)}{v^{(p-1)\beta}} \mid u \in R^+, v \in (0, r_1]\} < +\infty. \tag{8}$$

Let $\epsilon = \min\{\frac{r_1(1-\gamma)}{\int_0^{\frac{1}{2}} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau)d\tau)ds}, (\frac{C_1}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q(\int_s^{\frac{1}{2}} h_1(\tau)d\tau)ds)^{\frac{-\beta-1}{\beta}}\}$, by the first limit of H_4 , there exist a sufficient small number $r_2 \in (0, 1)$ such that

$$g(u) \leq \epsilon^{p-1}u^{\frac{p-1}{\beta}}, \forall u \in [0, r_2]. \tag{9}$$

Take $r = \min\{r_1, r_2\}$, by (9), we can get

$$\begin{aligned} v(t) &= \int_0^{\frac{1}{2}} \phi_q \left(\int_s^t h_2(\tau)g(u(\tau))d\tau \right) ds + \\ &\frac{\gamma}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_2(\tau)g(u(\tau))d\tau \right) ds \\ &\leq \frac{1}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_2(\tau)g(u(\tau))d\tau \right) ds \\ &\leq \frac{\epsilon}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_2(\tau)d\tau \right) ds \|u\|^{\frac{1}{\beta}} \\ &\leq r_1^{1+\frac{1}{\beta}} < r_1, \forall u \in K \cap \partial K_r, s \in [0, 1]. \end{aligned}$$

By (8), we can get

$$\begin{aligned} \|Tu\| &\leq \left| \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)f(u(\tau), v(\tau))d\tau \right) ds + \right. \\ &\frac{\gamma}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)f(u(\tau), v(\tau))d\tau \right) ds, \left. \right| \\ &\leq \frac{C_1}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)d\tau \right) ds \times \\ &\left(\frac{\epsilon}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)d\tau \right) ds \right)^\beta \|u\| \\ &= C_1 \epsilon^\beta \left(\frac{1}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)d\tau \right) ds \right)^{\beta+1} \|u\| \\ &\leq \|u\|, \forall u \in K \cap \partial K_r, t \in [0, 1]. \end{aligned}$$

So $\|Tu\| \leq \|u\|, \forall u \in K \cap \partial K_r$, by lemma 1.4, we get

$$i(T, K \cap K_r, K) = 1. \tag{10}$$

By lemma 1.5, T has at least one fixed point in $K \cap (\overline{K_N} \setminus K_r)$, so problem (1) has at least a system positive solution.

Theorem 2.2 Suppose $(H_1), (H_2), (H_3), (H_5), (H_6)$ hold, then problem (1) has at least two systems positive solutions.

Proof By (H_5) , there exists $\mu > 0$ and a sufficient small number $\xi \in (0, 1)$, such that

$$f(u, v) \geq \mu^{p-1} v^{n(p-1)}, \forall u \in R^+, 0 \leq v \leq \xi, \tag{11}$$

$$g(u) \geq (C_2 u)^{\frac{p-1}{n}}, \forall 0 \leq u \leq \xi, \tag{12}$$

where

$C_2 = 2 \left(\frac{\mu \epsilon^2}{1-\gamma} \left(\frac{\gamma}{1-\gamma} \right)^n \int_\epsilon^\eta \phi_q(k_1(s)) ds \int_\epsilon^\eta (\phi_q(k_2(s)))^n ds \right)^{-1}$
 since $g \in C(R^+, R^+)$, $g(0) \equiv 0$, so there exists $\sigma \in (0, \xi)$ such that $\forall u \in [0, \sigma]$, we have

$$g(u) \leq \left(\frac{1}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)d\tau \right) ds \right)^{-1},$$

this imply

$$v(t) \leq \frac{1}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_2(\tau)g(u(\tau))d\tau \right) ds \leq \xi, \forall u \in K \cap \partial K_\sigma. \tag{13}$$

By using Jensen inequality, $0 < q \leq 1$, and (11)-(13), we can get

$$\begin{aligned} (Tu)_{(\frac{1}{2})} &\geq \frac{\mu}{1-\gamma} \int_\epsilon^\eta \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)d\tau \right) ds \times \\ &\left(\frac{\gamma}{1-\gamma} \int_\epsilon^\eta \phi_q \left(\int_s^{\frac{1}{2}} h_2(\tau)g(u(\tau))d\tau \right) ds \right)^n \\ &\geq \frac{\mu}{1-\gamma} \int_\epsilon^\eta \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)d\tau \right) ds \times \\ &\left(\frac{\gamma}{1-\gamma} \right)^n \int_\epsilon^\eta (\phi_q \left(\int_s^{\frac{1}{2}} h_2(\tau)g(u(\tau))d\tau \right))^n ds \\ &\geq \frac{\mu C_2 \epsilon^2}{1-\gamma} \left(\frac{\gamma}{1-\gamma} \right)^n \int_\epsilon^\eta \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)d\tau \right) ds \times \\ &\int_\epsilon^\eta (\phi_q \left(\int_s^{\frac{1}{2}} h_2(\tau)d\tau \right))^n ds \|u\| \\ &= 2 \|u\|, \forall u \in K \cap \partial K_\sigma. \end{aligned}$$

So $\|Tu\| > \|u\|, \forall u \in K \cap \partial K_\sigma$, by lemma 1.4, we can get

$$i(T, K \cap K_\sigma, K) = 0. \tag{14}$$

We can choose $N > R > \sigma$, such that (7),(14) hold together.

On the other hand by (3),(4) and H_6 we can get

$$\begin{aligned} (Tu)(t) &< \frac{1}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)f(u(\tau), v(\tau))d\tau \right) ds \\ &\leq \frac{\gamma}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)d\tau \right) ds \times \\ &f(R, \frac{\gamma}{1-\gamma} \int_0^{\frac{1}{2}} \phi_q \left(\int_s^{\frac{1}{2}} h_1(\tau)d\tau \right) ds g(R)) \\ &< R, \forall u \in K \cap K_R, \forall t \in [0, 1]. \end{aligned}$$

So for any $u \in K \cap K_R$, by lemma 1.4, we can get

$$i(T, K \cap K_R, K) = 1. \tag{15}$$

By (7),(14),(15), we have

$$\begin{aligned} &i(T, K \cap (K_N \setminus \overline{K_R}), K) \\ &= i(T, K \cap K_N, K) - i(T, K \cap K_R, K) \\ &= -1. \\ &i(T, K \cap (K_R \setminus \overline{K_\sigma}), K) \\ &= i(T, K \cap K_R, K) - i(T, K \cap K_\sigma, K) \\ &= 1 \end{aligned}$$

So T have at least two fixed points in $K \cap (K_N \setminus \overline{K_R})$ and $K \cap (K_R \setminus \overline{K_\sigma})$, by (4), problem (1) has at least two system solutions.

REFERENCES

- [1] Su H, Wei Z L, Wang B H. The existence of positive solutions four a nonlinear four-point singular boundary value problem with a p-Laplace operator. *Nonlinear anal*,2007,66:2204-2217.
- [2] Ma D x, Han J X,Chen X G. Positive solution of boundary value problem for one-dimensional p-Laplacian with singularities. *J Math Anal Appl*,2006,324:118-133.
- [3] Liu Y J, Ge W G. Multiple positive solutions to a three-point boundary value problems with p-Laplacian. *J Math Anal Appl*,2003,277:293-302
- [4] Jin J X, Yin C H. Positive solutions for the boundary value problems of one-dimensional p-Laplacian with delay. *J Math Anal Appl*, 2007, 330:1238-1248.

- [5] Sun Y P. Optimal existence criteria for symmetric positive solutions a three-point boundary differential equations. *Nonlinear anal*,2007,66:1051-1063
- [6] Tian Yuansheng Liu Chungen. The existence of symmetric positive solutions for a three-point singular boundary value problem with a p-Laplace operator. *Acta Mathematica scientia* 2010,30A(3):784-792.
- [7] Xie Shengli. Positive solutions of multiple-point boundary value problems for systems of nonlinear second order differential equations .*Acta Mathematica scientia* 2010,30(A):258-266.
- [8] Guo Dajun *Nonlinear functional analysis*. Jinan. Shandong science and technology publishing house,2001 .
- [9] Guo D,Lakshmikantham V. *Nonlinear Problems in Abstract Cones*.New YorkAcademic Press,1988