

# Uniformly Strong Persistence for a Predator-Prey Model with Modified Leslie-Gower and Holling-Type II Schemes

Changjin Xu, Yusen Wu

**Abstract**—In this paper, a asymptotically periodic predator-prey model with Modified Leslie-Gower and Holling-Type II schemes is investigated. Some sufficient conditions for the uniformly strong persistence of the system are established. Our result is an important complementarity to the earlier results.

**Keywords**—Predator-prey model, uniformly strong persistence, asymptotically periodic, Holling-type II.

## I. INTRODUCTION

IT is well known that the dynamical behavior of predator-prey systems is a form of very common biological interaction in the natural world. This topic has attracted a lot of attention and many good results have already been reported. For example, Chen and Chen [1] studied the linear stability of trivial periodic solution and semi-trivial periodic solutions of a periodic predator-prey system with distributed time delays and impulsive effect. Mukherjee [2] made a discussion on the uniform persistence in a generalized prey-predator system with parasitic infection. Chen [3] gave a theoretical study on the almost periodic solution of the non-autonomous two-species competitive model with stage structure. Sen et al. [4] analyzed the bifurcation behavior of a ratio-dependent prey-predator model with the Allee effect. Agiza et al. [5] investigated the chaotic phenomena of a discrete prey-predator model with Holling type II. Aggelis et al. [6] considered the coexistence of both prey and predator populations of a prey-predator model. Nindjin and Aziz-Alaoui [7] focused on the persistence and global stability in a delayed Leslie-Gower type three species food chain. Ko and Ryu [8] discussed the coexistence states of a nonlinear Lotka-Volterra type predator-prey model with cross-diffusion. Fazly and Hesaaraki [9] dealt with periodic solutions of a predator-prey system with monotone functional responses. One can see [10-52] etc. For more related studies. However, the research work on asymptotically periodic predator-prey model is very few at present.

The so-called asymptotically periodic function is the function  $\bar{a}(t)$  which can be expressed by the form  $\bar{a}(t) = a(t) + \tilde{a}(t)$ , where  $a(t)$  is a periodic function and  $\tilde{a}(t)$  satisfies  $\lim_{t \rightarrow +\infty} \tilde{a}(t) = 0$ .

In 2003, Aziz-Alaoui and Okiye [53] investigated the stability and bifurcation of the following predator-prey model with

C. Xu is with the Guizhou Key Laboratory of Economics System Simulation, School of Mathematics and Statistics, Guizhou University of Finance and Economics, Guiyang 550004, PR China e-mail: xcj403@126.com.

P. Li is with the School of Mathematics and Statistics, Henan University of Science and Technology, Luoyang 471023, PR China e-mail: lplplpl@163.com.

time delay

$$\begin{cases} \frac{dx}{dt} = x(t) \left[ a - bx(t) - \frac{cy}{x(t)+k_1} \right], \\ \frac{dy}{dt} = y(t) \left[ d - \frac{ey(t)}{x(t)+k_2} \right], \end{cases} \quad (1)$$

with initial conditions  $x(0) \geq 0, y(0) \geq 0$ , where  $x(t)$  denotes the densities of prey, at time  $t$ ;  $y(t)$  denotes the densities of the predator at time  $t$ ;  $a, b, c, e, k_1, k_2$  are all positive constants. In details, one can see [53].

We must point out that all biological and environment parameters in model (1) are constants in time. However, any biological or environmental parameters are naturally subject to fluctuation in time. Thus the effect of a periodically varying environment is important for evolutionary theory as the selective forces on systems in a fluctuating environment differ from those in a stable environment. Therefore, the assumptions of periodicity of the parameters are a way of incorporating the periodicity the environment (such as seasonal effects of weather, food supplies, mating habits and so on). Stimulated by above discussion and considering the asymptotically periodic function, in this paper, we will modify system (1) as the form

$$\begin{cases} \frac{dx}{dt} = x(t) \left[ a(t) + \tilde{a}(t) - (b(t) + \tilde{b}(t))x(t) - \frac{(c(t) + \tilde{c}(t))y}{x(t) + k_1(t) + \tilde{k}_1(t)} \right], \\ \frac{dy}{dt} = y(t) \left[ d(t) + \tilde{d}(t) - \frac{(e(t) + \tilde{e}(t))y(t)}{x(t) + k_2(t) + \tilde{k}_2(t)} \right], \end{cases} \quad (2)$$

with initial conditions  $x(0) \geq 0, y(0) \geq 0$ .

The principle object of this article is to investigate the uniformly strong persistence of system (2). Only very few papers which deal with this topic, see [10,54].

In this paper, we always assume that system (2) satisfies

(H)  $a(t), b(t), c(t), d(t), k_1(t), k_2(t)$  are continuous, non-negative periodic functions;  $\tilde{a}(t), \tilde{b}(t), \tilde{c}(t), \tilde{d}(t), \tilde{k}_1(t), \tilde{k}_2(t)$  are continuous, nonnegative asymptotically items of asymptotically periodic functions.

## II. UNIFORMLY STRONG PERSISTENCE

In this section, we shall present some result about the uniformly strong persistence of system (2). For convenience and simplicity in the following discussion, we introduce the notations, definitions and Lemmas. Let

$$0 < f^l = \lim_{t \rightarrow +\infty} \inf f(t) \leq \lim_{t \rightarrow +\infty} \sup f(t) = f^u < +\infty.$$

In view of the definitions of lower limit and upper limit, it follows that for any  $\varepsilon > 0$ , there exists  $T > 0$  such that

$$f^l - \varepsilon \leq f(t) \leq f^u + \varepsilon, \text{ for } t \geq T. \quad (3)$$

**Definition 1.** The system (2) is said to be strong persistence, if every solution  $x(t)$  of system (2) satisfies

$$0 < \lim_{t \rightarrow +\infty} \inf x(t) \leq \lim_{t \rightarrow +\infty} \sup x(t) \leq \delta < +\infty.$$

**Lemma 1.** Both the positive and nonnegative cones of  $R^2$  are invariant with respect to system (2).

It follows from Lemma 1 that any solution of system (2) with a nonnegative initial condition remains nonnegative.

**Lemma 2.[10]** If  $a > 0, b > 0$  and  $\dot{x}(t) \geq (\leq)x(t)(b - ax^\alpha(t))$ , where  $\alpha$  is a positive constant, when  $t \geq 0$  and  $x(0) > 0$ , we have

$$x(t) \geq (\leq) \left(\frac{b}{a}\right)^{\frac{1}{\alpha}} \left[1 + \left(\frac{bx^{-\alpha}(0)}{a} - 1\right)e^{-b\alpha t}\right]^{-\frac{1}{\alpha}}.$$

In the following, we will ready to state our result.

**Theorem 1.** Let  $\theta_2$  be defined by (9). Assume that the condition (H) and  $a^l k_1^l > c^u \theta_2$  hold, then system (2) is uniformly strong persistence.

**Proof** It follows from (3) that for any  $\varepsilon > 0$ , there exists  $T_1 > 0$  such that for  $t \geq T_1$ ,

$$\begin{cases} a^l - \varepsilon \leq a(t) \leq a^u + \varepsilon, -\varepsilon < \tilde{a}(t) < \varepsilon, \\ b^l - \varepsilon \leq b(t) \leq b^u + \varepsilon, -\varepsilon < \tilde{b}(t) < \varepsilon, \\ c^l - \varepsilon \leq c(t) \leq c^u + \varepsilon, -\varepsilon < \tilde{c}(t) < \varepsilon, \\ d^l - \varepsilon \leq d(t) \leq d^u + \varepsilon, -\varepsilon < \tilde{d}(t) < \varepsilon, \\ k_1^l - \varepsilon \leq k_1(t) \leq k_1^u + \varepsilon, -\varepsilon < \tilde{k}_1(t) < \varepsilon, \\ k_2^l - \varepsilon \leq k_2(t) \leq k_2^u + \varepsilon, -\varepsilon < \tilde{k}_2(t) < \varepsilon. \end{cases} \quad (4)$$

Substituting (4) into the first equation of system (2), we have

$$\begin{aligned} \frac{dx}{dt} &= x(t) \left[ a(t) + \tilde{a}(t) - (b(t) + \tilde{b}(t))x(t) \right. \\ &\quad \left. - \frac{(c(t) + \tilde{c}(t))y}{x(t) + k_1(t) + \tilde{k}_1(t)} \right] \\ &\leq x(t) \left[ a(t) + \tilde{a}(t) - (b(t) + \tilde{b}(t))x(t) \right] \\ &\leq x(t) \left[ (a^u + 2\varepsilon) - (b^l - 2\varepsilon)x(t) \right]. \end{aligned} \quad (5)$$

By Lemma 2, we get

$$\lim_{t \rightarrow +\infty} \sup x(t) \leq \frac{a^u}{b^l} := \theta_1. \quad (6)$$

Then for any  $\varepsilon > 0$ , there exists  $T_2 > T_1 > 0$  such that

$$x(t) \leq \theta_1 + \varepsilon, \quad t \geq T_2. \quad (7)$$

Similarly, from (3) and the second equation of system (2), we obtain that for any  $\varepsilon > 0$ , there exists  $T_3 > T_2 > 0$  such that

$$\begin{aligned} \dot{y}(t) &= y(t) \left[ d(t) + \tilde{d}(t) - \frac{(e(t) + \tilde{e}(t))y(t)}{x(t) + k_2(t) + \tilde{k}_2(t)} \right] \\ &\leq y(t) \left[ (d^u - 2\varepsilon) - \frac{(e^l - 2\varepsilon)y(t)}{\theta_1 + \varepsilon + k_2^u + 2\varepsilon} \right]. \end{aligned} \quad (8)$$

In view of Lemma 2, we derive

$$\lim_{t \rightarrow +\infty} \sup y(t) \leq \frac{d^u(\theta_1 + k_2^u)}{e^l} := \theta_2. \quad (9)$$

Then for any  $\varepsilon > 0$ , there exists  $T_4 > T_3 > 0$  such that

$$y(t) \leq \theta_2 + \varepsilon, \quad t \geq T_4. \quad (10)$$

By (7), (10) and the first equation of system (2), we obtain that for any  $\varepsilon > 0$ , there exists  $T_5 > T_4 > 0$  such that

$$\begin{aligned} \frac{dx}{dt} &= x(t) \left[ a(t) + \tilde{a}(t) - (b(t) + \tilde{b}(t))x(t) \right. \\ &\quad \left. - \frac{(c(t) + \tilde{c}(t))y}{x(t) + k_1(t) + \tilde{k}_1(t)} \right] \\ &\geq x(t) \left[ (a^l - 2\varepsilon) - (b^u + 2\varepsilon)x(t) \right. \\ &\quad \left. - \frac{(c^u + 2\varepsilon)(\theta_2 + \varepsilon)}{k_1^l - 2\varepsilon} \right]. \end{aligned} \quad (11)$$

Using Lemma 2 again, we have

$$\lim_{t \rightarrow +\infty} \inf x(t) \geq \frac{a^l k_1^l - c^u \theta_2}{k_1^l b^u} := \delta_1. \quad (12)$$

Thus for any  $\varepsilon > 0$ , there exists  $T_6 > T_5 > 0$  such that

$$x(t) \geq \delta_1 - \varepsilon. \quad (13)$$

According (7), (10) and the second equation of system (2), we obtain that for any  $\varepsilon > 0$ , there exists  $T_7 > T_6 > 0$  such that

$$\begin{aligned} \dot{y}(t) &= y(t) \left[ d(t) + \tilde{d}(t) - \frac{(e(t) + \tilde{e}(t))y(t)}{x(t) + k_2(t) + \tilde{k}_2(t)} \right] \\ &\geq y(t) \left[ d^l + 2\varepsilon - \frac{e^u + 2\varepsilon}{k_2^l - 2\varepsilon} y(t) \right]. \end{aligned} \quad (14)$$

Using Lemma 2 again, we have

$$\lim_{t \rightarrow +\infty} \inf y(t) \geq \frac{d_1^l e^u}{k_2^l} := \delta_2. \quad (15)$$

Thus we complete the proof of Theorem 1.

### III. CONCLUSIONS

In this paper, we have investigated a asymptotically periodic predator-prey model with modified Leslie-gower and Holling-type II schemes. A set of sufficient conditions for the uniformly strong persistence of the system are derived. It is shown that under some suitable conditions, the asymptotically periodic predator-prey model with modified Leslie-Gower and Holling-type II schemes is uniformly strong persistence. Our results obtained in this paper complement the earlier results.

### ACKNOWLEDGMENT

This work is supported by National Natural Science Foundation of China(No.11261010; No.60902044; No.11161015), Natural Science and Technology Foundation of Guizhou Province(J[2012]2100), Governor Foundation of Guizhou Province([2012]53), 125 Science and Technology Grand Project of Department of Education of Guizhou Province ([2012]011) and Natural Science Innovation Team Project of Guizhou Province ([2013]14).

## REFERENCES

- [1] L.J. Chen, F.D. Chen Dynamic behaviors of the periodic predator-prey system with distributed time delays and impulsive effect, *Nonlinear Anal.: Real World Appl.* 12 (4) (2011) 2467-2473.
- [2] D. Mukherjee, Uniform persistence in a generalized prey-predator system with parasitic infection. *Biosystems* 47 (3) (1998) 149-155.
- [3] F.D. Chen, Almost periodic solution of the non-autonomous two-species competitive model with stage structure. *Appl. Math. Comput.* 181 (1) (2006) 685-693.
- [4] M. Sen, M. Banerjee, A. Morozov, Bifurcation analysis of a ratio-dependent prey-predator model with the Allee effect. *Ecological Complexity* 11 (2012) 12-27.
- [5] H.N. Agiza, E.M. Elabbasy, H. El-Metwally, A.A. Elsadany, Chaotic dynamics of a discrete prey-predator model with Holling type II. *Nonlinear Anal.: Real World Appl.* 10 (1) (2009) 116-129.
- [6] G. Aggelis, D.V. Vayenas, V. Tsagou, S. Pavlou, Prey-predator dynamics with predator switching regulated by a catabolic repression control mode, *Ecological Modelling* 183 (4) (2005) 451-462.
- [7] A.F. Nindjin, M.A. Aziz-Alaoui, Persistence and global stability in a delayed Leslie-Gower type three species food chain. *J. Math. Anal. Appl.* 340 (1) (2008) 340-357.
- [8] W. Ko, K. Ryu, Coexistence states of a nonlinear Lotka-Volterra type predator-prey model with cross-diffusion. *Nonlinear Anal.: TMA* 71 (12) (2009) 1109-1115.
- [9] M. Fazly, M. Hesaaraki, Periodic solutions for a discrete time predator-prey system with monotone functional responses. *Comptes Rendus de l'Académie des Sciences-Series I* 345 (4) (2007) 199-202.
- [10] F.D. Chen, On a nonlinear nonautonomous predator-prey model with diffusion and distributed delay. *J. Comput. Appl. Math.* 180 (1) (2005) 33-49.
- [11] Z.J. Liu, S.M. Zhong, X.Y. Liu, Permanence and periodic solutions for an impulsive reaction-diffusion food-chain system with holling type III functional response. *J. Franklin Inst.* 348 (2) (2011) 277-299.
- [12] A.F. Nindjin, M.A. Aziz-Alaoui, M. Cadivel, Analysis of a predator-prey model with modified Leslie-Gower and Holling-type II schemes with time delay. *Nonlinear Anal.: Real World Appl.* 7 (5) (2006) 1104-1118.
- [13] M. Scheffer, Fish and nutrients interplay determines algal biomass: a minimal model, *Oikos* 62 (1991) 271-282.
- [14] Y.K. Li, K.H. Zhao, Y. Ye, Multiple positive periodic solutions of species delay competition systems with harvesting terms. *Nonlinear Anal.: Real World Appl.* 12 (2)(2011) 1013-1022.
- [15] R. Xu, L.S. Chen, F.L. Hao, Periodic solution of a discrete time Lotka-Volterra type food-chain model with delays. *Appl. Math. Comput.* 171 (1) (2005) 91-103.
- [16] T.K. Kar, S. Misra, B. Mukhopadhyay, A bioeconomic model of a ratio-dependent predator-prey system and optimal harvesting. *J. Appl. Math. Comput.* 22 (1-2) (2006) 387-401.
- [17] R. Bhattacharyya, B. Mukhopadhyay, On an eco-epidemiological model with prey harvesting and predator switching: Local and global perspectives. *Nonlinear Anal.: Real World Appl.* 11 (5) (2010) 3824-3833.
- [18] K. Chakraborty, M. Chakraborty, T. K. Kar, Bifurcation and control of a bioeconomic model of a prey-predator system with a time delay. *Nonlinear Anal.: Hybrid Sys.* 5 (4) (2011) 613-625.
- [19] W.P. Zhang, D.M. Zhu, P. Bi, Multiple periodic positive solutions of a delayed discrete predator-prey system with type IV functional responses. *Appl. Math. Lett.* 20 (10) (2007) 1031-1038.
- [20] R.H. Wu, X.L. Zou, K. Wang, Asymptotic behavior of a stochastic non-autonomous predator-prey model with impulsive perturbations. *Comm. Nonlinear Sci. Numer. Simul.* 20 (3) (2015) 965-974.
- [21] A. Moussaoui, S. Bassaid, E.H.A. Dads, The impact of water level fluctuations on a delayed prey-predator model. *Nonlinear Anal.: Real World Appl.* 21 (2015) 170-184.
- [22] C. Wang, X.Y. Li, Further investigations into the stability and bifurcation of a discrete predator-prey model. *J. Math. Anal. Appl.* 422 (2) (2015) 920-939.
- [23] X.H. Wang, J.W. Jia, Dynamic of a delayed predator-prey model with birth pulse and impulsive harvesting in a polluted environment. *Phys. A: Statistical Mech. Appl.* 422(2015) 1-15.
- [24] M. Haque, M.S. Rahman, E. Venturino, B.L. Li, Effect of a functional response-dependent prey refuge in a predator-prey model. *Ecological Complexity* 20 (2014) 248-256.
- [25] Q.Y. Bie, Q.R. Wang, Z.A. Yao, Cross-diffusion induced instability and pattern formation for a Holling type-II predator-prey model, *Appl. Math. Comput.* 247 (2014) 1-12.
- [26] M. Sen, M. Banerjee, A. Morozov, Stage-structured ratio-dependent predator-prey models revisited: When should the maturation lag result in systems destabilization? *Ecological Complexity* 19 (2014) 23-34.
- [27] S.H. Xu, Dynamics of a general prey-predator model with prey-stage structure and diffusive effects. *Comput. Math. Appl.* 68 (3) (2014) 405-423.
- [28] Y.H. Fan, L.L. Wang, Multiplicity of periodic solutions for a delayed ratio-dependent predator-prey model with monotonic functional response and harvesting terms. *Appl. Math. Comput.* 244 (2014) 878-894.
- [29] P.J. Pal, P.K. Mandal, Bifurcation analysis of a modified Leslie-Gower predator-prey model with Beddington-DeAngelis functional response and strong Allee effect. *Math. Comput. Simul.* 97 (2014) 123-146.
- [30] S. Bhattacharya, M. Martcheva, X.Z. Li, A predator-prey-disease model with immune response in infected prey. *J. Math. Anal. Appl.* 411 (1) (2014) 297-313.
- [31] J.N. Wang, W.H. Jiang, Impulsive perturbations in a predator-prey model with dormancy of predators. *Appl. Math. Model.* 38 (9-10) (2014) 2533-2542.
- [32] J. P. Tripathi, S. Abbas, M. Thakur, A density dependent delayed predator-prey model with Beddington-DeAngelis type function response incorporating a prey refuge. *Commun. Nonlinear Sci. Numer. Simul.* 22 (1-3) (2015) 427-450.
- [33] Z.P. Ma, W.T. Li, Bifurcation analysis on a diffusive Holling-Tanner predator-prey model. *Appl. Math. Model.* 37(6) (2013) 4371-4384.
- [34] R. Yuan, W.H. Jiang, Y. Wang, Saddle-node-Hopf bifurcation in a modified Leslie-Gower predator-prey model with time-delay and prey harvesting. *J. Math. Anal. Appl.* 422 (2) (2015) 1072-1090.
- [35] J.D. Flores, E. Gonzalez-Olivares, Dynamics of a predator-prey model with Allee effect on prey and ratio-dependent functional response. *Ecological Complexity* 18 (2014) 59-66.
- [36] Y.M. Chen, F.Q. Zhang, Dynamics of a delayed predator-prey model with predator migration. *Appl. Math. Model.* 37 (3) (2013) 1400-1412.
- [37] Y. Zhang, Q.L. Zhang, X.G. Yan, Complex dynamics in a singular Leslie-Gower predator-prey bioeconomic model with time delay and stochastic fluctuations. *Phys. A: Statistical Mech. Appl.* 404 (2014) 180-191.
- [38] C.J. Xu, Q.M. Zhang, Dynamical behavior of a delayed diffusive predator-prey model with competition and type III functional response. *J. Egyptian Math. Soc.* 22(3) (2014) 379-385.
- [39] O. Makarenkov, Topological degree in the generalized Gause prey-predator model. *J. Math. Anal. Appl.* 410(2) (2014) 525-540.
- [40] A.J. Terry, A predator-prey model with generic birth and death rates for the predator. *Math. Biosci.* 248 (2014) 57-66.
- [41] J. Zhou, Positive steady state solutions of a Leslie-Gower predator-prey model with Holling type II functional response and density-dependent diffusion. *Nonlinear Anal.: TMA* 82 (2013) 47-65.
- [42] Z.J. Du, Z.S. Feng, Periodic solutions of a neutral impulsive predator-prey model with Beddington-DeAngelis functional response with delays. *J. Comput. Appl. Math.* 258(2014) 87-98.
- [43] Z.Z. Ma, F.D. Chen, C.Q. Wu, W.L. Chen, Dynamic behaviors of a Lotka-Volterra predator-prey model incorporating a prey refuge and predator mutual interference. *Appl. Math. Comput.* 219(15) (2013) 7945-7953.
- [44] Y.F. Lv, R. Yuan, Existence of traveling wave solutions for Gause-type models of predator-prey systems. *Appl. Math. Comput.* 229 (2014) 70-84.
- [45] C. Liu, Q.L. Zhang, J.N. Li, W.Q. Yue, Stability analysis in a delayed prey-predator-resource model with harvest effort and stage structure. *Appl. Math. Comput.* 238 (2014) 177-192.
- [46] H.W. Yin, X.Y. Xiao, X.Q. Wen, K. Liu, Pattern analysis of a modified Leslie-Gower predator-prey model with Crowley-Martin functional response and diffusion. *Comput. Math. Appl.* 67(8) (2014) 1607-1621.
- [47] J. Zhou, J.P. Shi, The existence, bifurcation and stability of positive stationary solutions of a diffusive Leslie-Gower predator-prey model with Holling-type II functional responses. *J. Math. Anal. Appl.* 405 (2) (2013) 618-630.
- [48] L.N. Guin, Existence of spatial patterns in a predator-prey model with self-and cross-diffusion. *Appl. Math. Comput.* 226 (2014) 320-335.
- [49] F.D. Chen, Z.Z. Ma, H.Y. Zhang, Global asymptotical stability of the positive equilibrium of the Lotka-Volterra prey-predator model incorporating a constant number of prey refuges. *Nonlinear Anal.: Real World Appl.* 13 (6) (2012) 2790-2793.
- [50] K. Das, M.N. Srinivas, M.A.S. Srinivas, N.H. Gazi, Chaotic dynamics of a three species prey-predator competition model with bionomic harvesting due to delayed environmental noise as external driving force. *CR Biol.* 335 (8) (2012) 503-513.

- [51] S. Jana, M. Chakraborty, K. Chakraborty, T.K. Kar, Global stability and bifurcation of time delayed prey-predator system incorporating prey refuge. *Math. Comput. Simul.* 85 (2012) 57-77.
- [52] X.Q. Ding, B.T. Su, J.M. Hao, Positive periodic solutions for impulsive Gause-type predator-prey systems. *Appl. Math. Comput.* 218 (12) (2012) 6785-6797.
- [53] M.A. Aziz-Alaoui, M.D. Okiye, Boundedness and global stability for a predator-prey model with modified Leslie-Gower and Holling-type II schemes. *Appl. Math. Lett.* 16(7)(2003) 1069-1075.
- [54] Y. Yang, W.C. Chen, Uniformly strong persistence of a nonlinear asymptotically periodic multispecies competition predator-prey system with general functional response. *Appl. Math. Compu.* 183 (1) (2006) 423-426.

**Changjin Xu** graduated from Huaihua University, China, in 1994. He received the M. S. degree from Kunming University of Science and Technology in 2004 and the Ph.D. degree from Central South University, China, in 2010. He is currently a professor at the Guizhou Key Laboratory of Economics System Simulation at Guizhou University of Finance and Economics. He has published about 100 refereed journal papers. He is a Reviewer of *Mathematical Reviews* and *Zentralblatt-Math*. His research interests include nonlinear systems, neural networks, stability and bifurcation theory.

**Yusen Wu** graduated from Liaocheng University, P.R. China, in 2004. He received the M. S. degree from Central South University, P.R. China in 2007 and the Ph.D. degree from Central South University, P.R. China, in 2010. He is currently an associate professor at School of Mathematics and Statistics of Henan University of Science and Technology. He has published about 30 refereed journal papers. He is a Reviewer of *Mathematical Reviews*. His research interests include the qualitative theory of ordinary differential equation and computer symbol calculation.